

CHARACTERIZATION OF SELF-REPORTED SLEEP QUALITY IN MALE AND FEMALE TRAINED AMATEUR TRIATHLETES: A DESCRIPTIVE STUDY

CARACTERIZACIÓN AUTO INFORMADA DE LA CALIDAD DEL SUEÑO EN TRIATLETAS AMATEURS MASCULINO Y FEMENINO ENTRENADOS: UN ESTUDIO DESCRIPTIVO

CARACTERIZAÇÃO AUTO REFERIDA DA QUALIDADE DO SONO EM TRIATLETAS AMADORES TREINADOS MASCULINOS E FEMININOS: UM ESTUDO DESCRITIVO

Luiz Vieira da Silva Neto¹, Tatiana Albuquerque Melo Kramer², Gloria Maria Alves Trajano¹ and Luiz José Frota Solon Júnior³

vieira_luiz@uvanet.br; albuquerqueitati@alu.ufc.br; efgloria.alves@gmail.com;
sluis424@gmail.com

¹Universidade Estadual Vale do Acaraú, Sobral, Brasil

²Universidade Federal do Ceará - UFC, Sobral, Brasil

³Universidade Federal da Paraíba, Joao Pessoa, Brasil

Original submission: 2023-07-21 Resubmitted: 2024-02-05, 2024-03-08

Accepted: 2024-03-11 Published: 2024-03-21

Doi: <https://doi.org/10.15517/pensarmov.v22i1.55925>

ABSTRACT

Sleep quality is a determining factor for the performance and health of athletes in general. The aim of the present study was to characterize the sleep quality in male and female trained amateur triathletes aged 20 to 59 years. A survey was conducted with 151 trained amateur triathletes, 108 male (38.6 ±8.1 years, triathlon training experience 5.8 ±4.3years, training frequency 6.3 ±0.9 days per week) and 43 female (39.3 ±7.6 years, triathlon training experience 4.8 ±3.3years, training frequency 6.5 ±0.6 days per week). The sleep quality was measured using the Pittsburgh Sleep Quality Index-Br (PSQI-Br), total values below 05 points indicate good sleepers, and values equal to or greater than 05 points indicate poor sleepers. Data from the subscales were analyzed using absolute and relative frequency. The other data in order to characterize sleep

quality were analyzed with median, mean, standard deviation, standard error, and 95% Confidence Interval of mean. Male and female triathletes have poor sleep quality (values equal or greater than 05 points), that can have negative effects on health and performance. In conclusion, all triathletes regardless of gender and age group have poor sleep quality.

Keywords: sleep, triathletes, performance, athlete training.

RESUMEN

La calidad del sueño es un factor determinante para el rendimiento y la salud de los deportistas en general, el objetivo del presente estudio fue caracterizar la calidad del sueño en triatletas aficionados masculinos y femeninos entrenados entre 20 y 59 años. Se realizó una encuesta con 151 triatletas aficionados entrenados, 108 hombres (38,6 \pm 8,1 años, experiencia en entrenamiento de triatlón 5,8 \pm 4,3 años, frecuencia de entrenamiento 6,3 \pm 0,9 días por semana) y 43 mujeres (39,3 \pm 7,6 años, experiencia en entrenamiento de triatlón 4,8 \pm 3,3). años, frecuencia de entrenamiento 6,5 \pm 0,6 días por semana). La calidad del sueño se midió mediante el índice de calidad del sueño de Pittsburgh-Br (PSQI-Br), los valores totales por debajo de 05 puntos indican que duermen bien y los valores iguales o superiores a 05 puntos indican que duermen mal. Los datos de las subescalas se analizaron utilizando frecuencias absolutas y relativas. Los demás datos para la caracterización de la calidad del sueño se analizaron con mediana, media, desviación estándar, error estándar e intervalo de confianza del 95% de la media. Los triatletas masculinos y femeninos tienen una mala calidad del sueño (valores iguales o superiores a 05 puntos), lo que puede tener efectos negativos en la salud y el rendimiento. En conclusión, todos los triatletas sin importar el género y el grupo de edad tienen una mala calidad del sueño.

Palabras clave: sueño, triatletas, desempeño, entrenamiento de los atletas.

RESUMO

A qualidade do sono é um fator determinante para o desempenho e a saúde de atletas em geral. O objetivo do presente estudo foi caracterizar a qualidade do sono em triatletas amadores treinados masculinos e femininos com idades de 20 à 59 anos. Uma pesquisa foi realizada com 151 triatletas amadores treinados, 108 homens (38,6 \pm 8,1 anos, experiência em treinamento de triatlo 5,8 \pm 4,3 anos, frequência de treinamento 6,3 \pm 0,9 dias por semana) e 43 mulheres (39,3 \pm 7,6 anos, experiência em treinamento de triatlo 4,8 \pm 3,3 anos, frequência de treino 6,5 \pm 0,6 dias por semana). A qualidade do

sono foi mensurada por meio do Índice de Qualidade do Sono de Pittsburgh-Br (PSQI-Br), valores totais abaixo de 05 pontos indicam bons dormidores e valores iguais ou maiores que 05 pontos indicam maus dormidores. Os dados das subescalas foram analisados por meio de frequência absoluta e relativa. Os demais dados para caracterização da qualidade do sono foram analisados com mediana, média, desvio padrão, erro padrão e intervalo de confiança de 95% da média. Triatletas masculinos e femininos apresentam má qualidade do sono (valores iguais ou superiores a 05 pontos), o que pode acarretar efeitos negativos na saúde e no desempenho. Em conclusão, todos os triatletas independentemente do sexo e faixa etária têm má qualidade do sono.

Palavras-chave: sono, triatletas, desempenho, treinamento atletico.

INTRODUCTION

Triathlon is a sport composed of the stages of swim, cycle, and run, performed in this order and without interruptions (Millet & Vleck, [2000](#)), and is divided by age categories for amateurs. Official Triathlon distances contested in races vary, with the shortest distance being the super sprint (250-300m swim, 05-08 Km cycle, and 1.5-02 Km run) (Quagliarotti et al., [2022](#)), and the Ultraman® Triathlon is a most long-distance race, held over three days (10 Km swim, 420.2 Km cycle e 84.4 Km run) (Smith et al., [2020](#)). In this sense, training involves high volumes to prepare for target distances (Etxebarria et al., [2019](#)) and needs efficient recovery between them.

In general, sleep has a crucial role in the recovery process of athletes (Fullagar et al., [2023](#)). Specifically, in triathlon, the literature presents only two profiles of studies relating sleep to this sport. The first profile comprises studies investigating general aspects of athletes' sleep, including several modalities (Costa & Re, [2023](#); Johnston et al., [2020](#); Vitale et al., [2019](#); Wilson, [2020](#)).

On the other hand, the second profile is composed of specific triathlon studies (Stevens et al., [2018](#); Sinisgalli et al., [2021](#); Kisiolek et al., [2022](#)), who seek to understand more directly the relationship between aspects of sleep and Triathlon. The study by Stevens et al. ([2018](#)) pointed out that a trip to perform a competition impaired the sleep of male master triathletes. Sinisgalli et al. ([2021](#)) indicate insufficient sleep duration in male and female amateur triathletes. Finally, in the study by Kisiolek et al. ([2022](#)), a negative correlation was found between performance and sleep duration. In this sense, although the literature has presented important findings regarding the sleep of triathletes, no study proposed to characterize sleep quality in a generalized way in amateur triathletes.

Characterizing the sleep quality of triathletes allows coaches to pay attention to possible associated variables that may influence their good or poor sleep quality, affecting their recovery. Such findings are important for coaches due to the relationship between sleep quality and the prevalence of injuries (de Sousa Nogueira Freitas et al., 2020), illnesses (Silva et al., 2022) decreased sports performance (Kisiolek et al., 2022), among others. So, the present study aimed to characterize the sleep quality in male and female trained amateur triathletes aged 20 to 59 years.

MATERIAL AND METHODS

Participants

Currently in Brazil there are approximately 25 thousand amateur triathletes (www.cbtri.com). Therefore, considering a confidence interval of 95% for the proportion of the sample and a margin of error of 10%, the sample size required for the present study is 96 triathletes (Serdar et al., 2021). The sample consisted of 151 amateur trained triathletes, 108 male and 43 female triathletes. Amateur trained, consists of participating in local competitions, training at least three times a week, identifying with the sport, training with a competitive purpose and developing sports skills (McKay et al., 2022). The sample characteristics (age, triathlon training experience, and training frequency) are shown in [table 1](#).

Table 1

Participant characteristics (n = 151)

Characteristics	Male (n = 108)	Female (n = 43)
Age (years)	38.6 ±8.1	39,3 ±7.6
Triathlon Training experience (years)	5.8 ±4.3	4.8 ±3.3
Training Frequency (days/week)	6.3 ±0.9	6.5 ±0.6
Training Hours (per day)	3.4 ±1.7	3.2 ±1.5
Training Hours (per week)	20.4 ±5.3	20.8 ±5.8
Height (cm)	175 ±0.10	168 ±0.08
Weight (Kg)	79.3 ±3.2	66 ±2.7
Body Massa Index	25.6 ±2.7	24.5 ±1.0

Source: Own authors.

All male and female volunteers should be healthy, training without interruptions for at least 06 months and composing the three stages of the triathlon weekly in their training. All volunteers were at the beginning of the pre-competitive period, aiming to participate in the respective local championships.

Only triathletes who voluntarily decided to participate, signing the Informed Consent Form, participated in the study. The ethics committee in human research

approved in 06/01/2021 the present study for research with human beings at the State University of Vale Acaraú with the number 4.749.908.

Procedures

All data collection took place remotely, in a virtual environment, via google forms. Volunteers were invited through the social networks of their respective clubs and advisory services where they trained.

After acceptance, a link to the electronic form was sent, containing the consent form on its first page, which, when read in full and accepted, followed up with the instruments for the acquisition of data on the sleep quality of the athletes.

Instruments

Pittsburgh Sleep Quality Index (PSQI-Br)

The PSQI is an instrument in the form of a questionnaire that measures sleep quality in the last 30 days. Developed by Buysse et al. ([1989](#)) and later cross-culturally validated for Brazil by Bertolazi et al. ([2011](#)), then identified as PSQI-Br.

The instrument consists of 19 items that measure seven components: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleep medication, and daytime dysfunction. The PSQI-Br total score is evaluated by adding the points for each component, generating a global score. Total values below 05 points indicate good sleepers, and values equal to or greater than 05 points indicate poor sleepers, and the higher the total score, the worse the sleep quality of those who answered the instrument. In the present study, the reliability of responses to items using the Cronbach's alpha measure was 0.71 (Landis & Koch, [1977](#)).

Statistical analyses

Response data for each instrument component were analyzed using absolute and relative frequency. All data and by category are presented as mean, median, and standard deviation. The standard error of the mean was calculated to provide a measure of the relative reliability of the collected means. In addition to the previous analyses, a 95% confidence interval (CI) for the mean was constructed. All data were analyzed using SPSS 23.0.

RESULTS

Below ([Table 2](#)) are the absolute and relative values separated between males and females of the answers to each item of the seven components of the PSQI-Br. (Vieira da Silva Neto et al., [2024](#)).

Table 2.

Absolute (N) and relative (%) frequencies of the PSQI-Br components for Male and Female.

		PSQI-Br			
		Male		Female	
		N	%	N	%
Subjctive sleep quality	Very good	15	13.9	13	30.2
	Fairly good	65	60.2	28	65.1
	Fairly bad	27	25	2	4.7
	Very bad	1	0.9	0	0
Sleep latency	≤15 min	29	26.9	14	32.6
	16-30 min	44	40.7	20	46.5
	31-60 min	30	27.8	9	20.9
	>60 min	5	4.6	0	0
Sleep duration	≥7 h	20	18.5	14	32.6
	≥6-7 h	67	62.0	23	53.5
	<6 h	19	17.6	6	14
	<5 h	2	1.9	0	0
Habitual sleep efficiency	≥85%	107	99.1	37	86
	75-84%	1	0.9	5	11.6
	65-74%	0	0	1	2.3
	<65%	0	0	0	0
Sleep disturbances	0	2	1.9	2	4.7
	1-9	57	52.8	28	65.1
	10-18	47	43.5	12	27.9
	19-27	2	1.9	1	2.3
Use of sleep medication	None	93	86.1	35	81.4
	Less than once per week	7	6.5	4	9.3
	Once or twice a week	3	2.8	2	4.7
	Three or more times a week	5	4.6	2	4.7
Day time dysfunction	0	15	13.9	13	30.2
	1-2	52	48.1	17	39.5
	3-4	31	30.2	13	30.2
	5-6	10	9.3	0	0

Source: Own authors.

We can characterize the sleep quality of female triathletes in [Table 3](#). In these results, according to the median and mean per age group and overall, all triathletes have poor sleep quality and are classified as poor sleepers, reinforced by the confidence interval present in the table.

Table 3.

Characterization of the PSQI-Br by Age group and Overall of Female Amateur Triathletes.

Age Group	N	Median	Mean	Standard deviation	Standard error	95 % CI of Mean
25-34	14	5	5	2.1	0.57	3.76 – 6.24
35-44	18	5	5.22	2.0	0.48	4.19 – 6.25
45-54	11	6	5.45	2.1	0.65	4.00 – 6.91
Overall	43	5	5.21	2.0	0.31	4.57 – 5.85

Source: Own authors.

When analyzing the results of the male triathletes, we can verify a pattern similar to that of the female group since all age groups and the total male sample have PSQI-Br values above 5 scores, both in the median and in the mean. Furthermore, the confidence intervals are all within the range that classifies the triathletes as poor sleepers and with poor sleep quality, see results in [Table 4](#).

Table 4.

Characterization of the PSQI-Br by Age group and Overall of Male Amateur Triathletes

Age Group	N	Median	Mean	Standard deviation	Standard error	95 % CI of Mean
20-29	16	5.5	5.94	2.62	0.65	4.54 – 7.33
30-39	48	6.5	6.44	2.55	0.36	5.70 – 7.18
40-49	35	6.0	6.03	2.20	0.37	5.27 – 6.79
50-59	9	7	7.44	1.81	0.60	6.05 – 8.84
Overall	108	6	6.31	2.40	0.23	5.86 – 6.77

Source: Own authors.

DISCUSSION

The present study aimed to characterize the sleep quality of male and female trained amateur triathletes, building consultation references for professionals who work directly with the performance and health of athletes involved in triathlon. In summary, all triathletes were classified as poor sleepers, having an overall PSQI score ≥ 5 .

When analyzing the results of the age groups in [Table 3](#), we can consider that all Triathletes are poor sleepers and have a poor sleep quality (Buysse et al., [1989](#); Bertolazi et al., [2011](#)), based on means and medians (Nakagawa & Cuthill, [2007](#); Lee, [2016](#)) of PSQI-Br. By analyzing all the indicators of the general sample, we can say those

female amateur triathletes have poor sleep quality, being classified as poor sleepers (Buysse et al., [1989](#); Bertolazi et al., [2011](#)).

Some factors may explain these results found in women, which may be associated with social and cultural aspects such as working hours, domestic work and family care. (Tang et al., [2017](#); Nordin et al., [2005](#)). Furthermore, although biological factors are associated with changes in sleep, sexual differences have also been noted and discussed in the literature due to chronic hormonal differences in women, over the years (Mallampalli & Carter, [2014](#)), which are relevant to poor sleep quality in female triathletes.

On the other hand, considering the results of male Triathletes, both by age group and overall, we can see that the entire sample has poor sleep quality, being classified as poor sleepers. Indeed, this is a common pattern of sleep quality that has been shown both for general male populations (Matsui et al., [2021](#)) and for the sports population (Swinbourne et al., [2016](#); Biggins et al., [2018](#)).

In summary, the findings presented in [Table 3](#) and [4](#) show indicators that can negatively affect the health and performance of Triathletes. The literature indicates that people who experience reduced sleep (≤ 7 h) or insufficient sleep are more likely to have illnesses, such as upper respiratory tract infections (Robinson et al., [2021](#)), as well as a greater chance of becoming injured (Johnston et al., [2020](#); Huang & Ihm, [2021](#)).

Additionally, some findings support that poor sleep quality also reduces muscle strength (Knowles et al., [2018](#)), impairs decision-making, speed, and accuracy of task performance, post-recovery-exercise (Troynikov et al., [2018](#)), and negatively alters important metabolic aspects (Besedovsky et al., [2019](#)), which are directly associated with the sports performance in triathletes. One factor that may explain poor sleep quality in triathletes is the option to train more, thus impacting the duration and sleep quality (Sargent et al., [2014](#)). Considering the results of the present study for male and female triathletes, actions such as promoting sleep hygiene practices or educational actions to improve sleep quality are necessary.

The main limitation of the present study is that sleep quality was not measured by objective measures such as polysomnography or actigraphy. Thus, the results are subject to subjective biases that the PSQI-Br presents. However, it is worth mentioning that the PSQI is a well-accepted and used instrument in the literature, in addition to being reliable in what it proposes to measure (Ibáñez et al., [2018](#)). The fact that we did not collect data on marital status or number of children also creates a limitation. Another limitation is that when categorizing by age, some groups have a small sample size, thus reducing the explanatory power of the results for these specific cases.

The present study can offer a practical way for professionals to monitor trained amateur triathletes. These indicators can be helpful in the management of sleep and its quality, allowing bold decisions involving the sleep of Triathletes.

CONCLUSION

Thus, we can conclude that male and female trained amateur triathletes have poor sleep quality, in addition to being poor sleepers, which can might potentially lead to several negative outcomes for health and performance, with these findings being more accurate mainly due to the standard error of the measurement presented in this study. With this, our study can provide parameters, so that when analyzing the sleep quality, triathletes or their coaches can see if they are above, within or below the results, thus allowing decision-making to modify the sleep quality.

REFERENCES

- Bertolazi, A. N., Fagondes, S. C., Hoff, L. S., Dartora, E. G., Miozzo, I. C., de Barba, M. E., & Barreto, S. S. (2011). Validation of the Brazilian Portuguese version of the Pittsburgh Sleep Quality Index. *Sleep Medicine*, 12(1), 70–75. <https://doi.org/10.1016/j.sleep.2010.04.020>
- Besedovsky, L., Lange, T., & Haack, M. (2019). The Sleep-Immune Crosstalk in Health and Disease. *Physiological Reviews*, 99(3), 1325–1380. <https://doi.org/10.1152/physrev.00010.2018>
- Biggins, M., Cahalan, R., Comyns, T., Purtill, H., & O'Sullivan, K. (2018). Poor sleep is related to lower general health, increased stress and increased confusion in elite Gaelic athletes. *The Physician and Sports Medicine*, 46(1), 14–20. <https://doi.org/10.1080/00913847.2018.1416258>
- Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R., & Kupfer, D. J. (1989). The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Research*, 28(2), 193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4)
- Costa, M., & Re, M. (2023). Chronotype and quality of sleep in alpine skiers. *Chronobiology International*, 40(5), 673–683. <https://doi.org/10.1080/07420528.2023.2203236>
- de Sousa Nogueira Freitas, L., da Silva, F. R., Andrade, H. A., Guerreiro, R. C., Paulo, F. V., de Mello, M. T., & Silva, A. (2020). Sleep debt induces skeletal muscle injuries in athletes: A promising hypothesis. *Medical Hypotheses*, 142, 109836. <https://doi.org/10.1016/j.mehy.2020.109836>

- Etxebarria, N., Mujika, I., & Pyne, D. B. (2019). Training and Competition Readiness in Triathlon. *Sports*, 7(5), 101. <https://doi.org/10.3390/sports7050101>
- Fullagar, H. H. K., Vincent, G. E., McCullough, M., Halson, S., & Fowler, P. (2023). Sleep and Sport Performance. *Journal of Clinical Neurophysiology*, 40(5), 408–416. <https://doi.org/10.1097/WNP.0000000000000638>
- Huang, K., & Ihm, J. (2021). Sleep and Injury Risk. *Current Sports Medicine Reports*, 20(6), 286–290. <https://doi.org/10.1249/JSR.0000000000000849>
- Ibáñez, V., Silva, J., & Cauli, O. (2018). A survey on sleep questionnaires and diaries. *Sleep Medicine*, 42, 90–96. <https://doi.org/10.1016/j.sleep.2017.08.026>
- Johnston, R., Cahalan, R., Bonnett, L., Maguire, M., Glasgow, P., Madigan, S., O'Sullivan, K., & Comyns, T. (2020). General health complaints and sleep associated with new injury within an endurance sporting population: A prospective study. *Journal of Science and Medicine in Sport*, 23(3), 252–257. <https://doi.org/10.1016/j.jsams.2019.10.013>
- Kisiolek, J. N., Smith, K. A., Baur, D. A., Willingham, B. D., Morrissey, M. C., Leyh, S. M., Saracino, P. G., Mah, C. D., & Ormsbee, M. J. (2022). Sleep Duration Correlates With Performance in Ultra-Endurance Triathlon. *International Journal of Sports Physiology and Performance*, 17(2), 226–233. <https://doi.org/10.1123/ijspp.2021-0111>
- Knowles, O. E., Drinkwater, E. J., Urwin, C. S., Lamon, S., & Aisbett, B. (2018). Inadequate sleep and muscle strength: Implications for resistance training. *Journal of Science and Medicine in Sport*, 21(9), 959–968. <https://doi.org/10.1016/j.jsams.2018.01.012>
- Landis, J. R., & Koch, G. G. (1977). The measurement of observer agreement for categorical data. *Biometrics*, 33(1), 159–174. <https://doi.org/10.2307/2529310>
- Lee, D. K. (2016). Alternatives to P value: confidence interval and effect size. *Korean Journal of Anesthesiology*, 69(6), 555–562. <https://doi.org/10.4097/kjae.2016.69.6.555>
- Mallampalli, M. P., & Carter, C. L. (2014). Exploring sex and gender differences in sleep health: a Society for Women's Health Research Report. *Journal of Women's Health*, 23(7), 553–562. <https://doi.org/10.1089/jwh.2014.4816>
- Matsui, K., Yoshiike, T., Nagao, K., Utsumi, T., Tsuru, A., Otsuki, R., Ayabe, N., Hazumi, M., Suzuki, M., Saitoh, K., Aritake-Okada, S., Inoue, Y., & Kuriyama, K. (2021). Association of Subjective Quality and Quantity of Sleep with Quality of Life among a General Population. *International Journal of Environmental Research and Public Health*, 18(23), 12835. <https://doi.org/10.3390/ijerph182312835>

- McKay, A. K. A., Stellingwerff, T., Smith, E. S., Martin, D. T., Mujika, I., Goosey-Tolfrey, V. L., Sheppard, J., & Burke, L. M. (2022). Defining Training and Performance Caliber: A Participant Classification Framework. *International Journal of Sports Physiology and Performance*, 17(2), 317–331. <https://doi.org/10.1123/ijsp.2021-0451>
- Millet, G. P., & Vleck, V. E. (2000). Physiological and biomechanical adaptations to the cycle to run transition in Olympic triathlon: review and practical recommendations for training. *British Journal of Sports Medicine*, 34(5), 384–390. <https://doi.org/10.1136/bjbm.34.5.384>
- Nakagawa, S., & Cuthill, I. C. (2007). Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biological Reviews of the Cambridge Philosophical Society*, 82(4), 591–605. <https://doi.org/10.1111/j.1469-185X.2007.00027.x>
- Nordin, M., Knutsson, A., Sundbom, E., & Stegmayr, B. (2005). Psychosocial factors, gender, and sleep. *Journal of Occupational Health Psychology*, 10(1), 54–63. <https://doi.org/10.1037/1076-8998.10.1.54>
- Quagliarotti, C., Gaiola, D., Bianchini, L., Vleck, V., & Piacentini, M. F. (2022). How to Form a Successful Team for the Novel Olympic Triathlon Discipline: The Mixed-Team-Relay. *Journal of Functional Morphology and Kinesiology*, 7(2), 46. <https://doi.org/10.3390/jfkm7020046>
- Robinson, C. H., Albury, C., McCartney, D., Fletcher, B., Roberts, N., Jury, I., & Lee, J. (2021). The relationship between duration and quality of sleep and upper respiratory tract infections: a systematic review. *Family Practice*, 38(6), 802–810. <https://doi.org/10.1093/fampra/cmab033>
- Sargent, C., Halson, S., & Roach, G. D. (2014). Sleep or swim? Early-morning training severely restricts the amount of sleep obtained by elite swimmers. *European Journal of Sport Science*, 14(supl.1), S310–S315. <https://doi.org/10.1080/17461391.2012.696711>
- Serdar, C. C., Cihan, M., Yücel, D., & Serdar, M. A. (2021). Sample size, power and effect size revisited: simplified and practical approaches in pre-clinical, clinical and laboratory studies. *Biochemia Medica*, 31(1), 010502. <https://doi.org/10.11613/BM.2021.010502>
- Silva, A., Pinto Pinheiro, L. S., Silva, S., Andrade, H., Pereira, A. G., Rodrigues da Silva, F., Guerreiro, R., Barreto, B., Resende, R., & Túlio de Mello, M. (2022). Sleep in Paralympic athletes and its relationship with injuries and illnesses. *Physical Therapy in Sport*, 56, 24–31. <https://doi.org/10.1016/j.ptsp.2022.06.001>

- Sinisgalli, R., de Lira, C. A. B., Vancini, R. L., Puccinelli, P. J. G., Hill, L., Knechtle, B., Nikolaidis, P. T., & Andrade, M. S. (2021). Impact of training volume and experience on amateur Ironman triathlon performance. *Physiology & Behavior*, 232, 113344. <https://doi.org/10.1016/j.physbeh.2021.113344>
- Smith, K. A., Kisiolek, J. N., Willingham, B. D., Morrissey, M. C., Leyh, S. M., Saracino, P. G., Baur, D. A., Cook, M. D., & Ormsbee, M. J. (2020). Ultra-endurance triathlon performance and markers of whole-body and gut-specific inflammation. *European Journal of Applied Physiology*, 120(2), 349–357. <https://doi.org/10.1007/s00421-019-04279-3>
- Stevens, C. J., Thornton, H. R., Fowler, P. M., Esh, C., & Taylor, L. (2018). Long-Haul Northeast Travel Disrupts Sleep and Induces Perceived Fatigue in Endurance Athletes. *Frontiers in Physiology*, 9, 1826. <https://doi.org/10.3389/fphys.2018.01826>
- Swinbourne, R., Gill, N., Vaile, J., & Smart, D. (2016). Prevalence of poor sleep quality, sleepiness and obstructive sleep apnoea risk factors in athletes. *European Journal of Sport Science*, 16(7), 850–858. <https://doi.org/10.1080/17461391.2015.1120781>
- Tang, J., Liao, Y., Kelly, B. C., Xie, L., Xiang, Y. T., Qi, C., Pan, C., Hao, W., Liu, T., Zhang, F., & Chen, X. (2017). Gender and Regional Differences in Sleep Quality and Insomnia: A General Population-based Study in Hunan Province of China. *Scientific reports*, 7, 43690. <https://doi.org/10.1038/srep43690>
- Troynikov, O., Watson, C. G., & Nawaz, N. (2018). Sleep environments and sleep physiology: A review. *Journal of Thermal Biology*, 78, 192–203. <https://doi.org/10.1016/j.jtherbio.2018.09.012>
- Vieira da Silva Neto, L., Albuquerque Melo Kramer, T., Alves Trajano, G.M., & Frota Solon Júnior, L.J. (2024). Data base of Characterization of self-reported sleep quality in male and female trained amateur triathletes: a descriptive study. *Pensar en Movimiento: Revista de Ciencias del Ejercicio y la Salud*, 22(1), e59280. <https://doi.org/10.15517/pensarmov.v22i1.59280>
- Vitale, J. A., La Torre, A., & Banfi, G. (2019). If RAR's acrophase is influenced by the sport discipline, how actigraphy-based sleep parameters vary in triathlon, volleyball and soccer athletes?. *Chronobiology International*, 36(6), 735–738. <https://doi.org/10.1080/07420528.2019.1594246>
- Wilson, P. B. (2020). Associations between sleep and in-race gastrointestinal symptoms: an observational study of running and triathlon race competitors. *Sleep Science*, 13(4), 293–297. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7856660/>