

## Regional diversity of Amphipoda in the Caribbean Sea

Alberto Martín<sup>1</sup>, Yusbelly Díaz<sup>1</sup>, Patricia Miloslavich<sup>1</sup>, Elva Escobar-Briones<sup>2</sup>, José Manuel Guerra-García<sup>3</sup>, Manuel Ortiz<sup>4</sup>, Bellineth Valencia<sup>5</sup>, Alan Giraldo<sup>5</sup> & Eduardo Klein<sup>1</sup>

1. Departamento de Estudios Ambientales. Centro de Biodiversidad Marina. Universidad Simón Bolívar. Caracas, 89000. Venezuela; amartinz@usb.ve, yusdiaz@usb.ve, pmilos@usb.ve, eklein@usb.ve
2. Instituto de Ciencias del Mar y Limnología. Unidad Académica Ecología y Biodiversidad Acuática. Universidad Nacional Autónoma de México. Ciudad de México, 04510. México; escobri@cmarl.unam.mx
3. Departamento de Zoología. Facultad de Biología. Universidad de Sevilla. Sevilla, 41012. España; jmaguerra@us.es
4. Facultad de Estudios Superiores Iztacala. Laboratorio de Crustáceos. Universidad Nacional Autónoma de México. Los Reyes Iztacala, 54090. México; ortiztouz@yaho.com
5. Departamento de Biología. Facultad de Ciencias Naturales y Exactas. Universidad del Valle. Cali, 10077. Colombia; alan.giraldo@correounivalle.edu.co, bellivalencia@yahoo.com

Received 21-XI-2012. Corrected 05-IV-2013. Accepted 06-V-2013.

**Abstract: Diversidad regional de Amphipoda en el Mar Caribe.** The order Amphipoda is one of the most diverse within Peracarids, and comprises 6 950 described marine species. Amphipod research in the Caribbean Sea began in the late 1 800s, but has increased significantly since 1 980. In this study, we analyzed the amphipod biodiversity (Caprelliidea, Gammaridea, Hyperiidea, and Ingolfiellidea) of the Caribbean Sea. For this, we compiled available data on species diversity of marine amphipods (data bases: WoRMS and OBIS and published species lists) into a comprehensive taxonomic list by country for the ecoregions of the Caribbean. Additionally, we analyzed the relative contribution of each country to regional diversity and the rate of discovery of new species. The Caribbean amphipod fauna is composed of 535 species within 236 genera and 73 families for the higher taxon. The Western Caribbean ecoregion holds the largest diversity (282 species), while the Eastern Caribbean recorded the lowest one (73). Mexico and Venezuela recorded the largest number of species with 266 and 206, respectively. Twelve countries had less than 50 species. The richest suborder is the Gammaridea with 381 species followed by the suborder Hyperiidea with 116. From the total of 535 amphipod species reported for the Caribbean region, 218 have the Caribbean as the holotype locality, and 132 are endemic (about 25% of the total). Areas of higher diversity seem to be concentrated along the Mexican Caribbean, Cuba and the Northern coast of South America (Venezuela-Colombia); however, such pattern is most likely reflecting local collection efforts and taxonomic expertise rather than actual distribution. Knowledge of amphipod species is mostly limited to shallow, near-shore waters, with little information available on the deep sea fauna. Regional research priorities for this group should be focused on completing shallow water coastal inventories of species in Central America and the Greater and Lesser Antilles. In addition, sampling the deep sea ecosystems should follow along with other particular habitats such as anchialine cave systems. It is also necessary to increase ecological research efforts, mainly in some speciose suborders, including the Caprelliidea and Hyperiidea, known to exhibit high diversity in other tropical localities. *Rev. Biol. Trop.* 61 (4): 1681-1720. Epub 2013 December 01.

**Key words:** Amphipoda, diversity, Caribbean Sea, endemism.

The order Amphipoda is one of the most diverse and globally distributed within the superorder Peracarida after order Isopoda. Its members occur worldwide in terrestrial, freshwater, subterranean and marine environments and in virtually all habitats (Foster, LeCroy, Heard & Vargas, 2009; LeCroy, Gasca,

Winfield, Ortiz & Escobar-Briones, 2009). They fulfill an important role in the trophic dynamics of many communities, as they break-down plant material and detritus, and also function as grazers, filter feeders, suspension feeders, commensals, micropredators, scavengers, or as prey for other carnivore species

(Bellan-Santini, 1998; Bellan-Santini & Ruffo, 1998; LeCroy et al., 2009).

Traditionally, the Amphipoda were divided into four suborders: the Gammaridea Latreille, 1802; the Caprellidea Leach, 1814; the Hyperiid Milne-Edwards, 1830; and the Ingolfiellidea Hansen, 1903 (Barnard, 1969; Bousfield, 1973, 2001; Lincoln, 1979; Martin & Davis, 2001). Despite, Myers & Lowry (2003) reestablished the suborder Corophiidea Leach, 1814, and placed the caprellideans *sensu lato* within it, we decided to use the traditional classification system because these categories will likely change with new phylogenetic studies.

Although there is a significant amount of scientific literature regarding Caribbean amphipod species in a variety of books, journals, and reports, many of these publications are restricted to one or a few species and many of them are old and therefore, not easily available. Comprehensive works on amphipods of the region are relatively rare.

The first records and descriptions of amphipods in the Caribbean Sea, began with the collections of gammarideans and corophiideans of the H.M.S. Challenger between 1873-76, when Stebbing described *Ampelisca abyssicola* from Culebra Island, St. Thomas (Lesser Antilles) (Stebbing, 1888). Further descriptions included *Shoemakerella cubensis* (as *Lysianax cubensis*) from Cuba (Stebbing, 1897), and *Talorchestia fritzi* (Stebbing, 1903), *Chelorchestia costaricana* (Stebbing, 1906) and *Pseudorchestoida biolleyi* (Stebbing, 1908) (as *Orchestoidea biolleyi*) from Costa Rica. Other expeditions to the Caribbean, particularly to Puerto Rico and Cuba, provided new amphipod descriptions (Shoemaker, 1933a,b, 1934, 1935, 1948). Hyperiid amphipods are also poorly studied in the Caribbean, and the knowledge we have comes from some historical larger surveys (Shoemaker, 1945, 1948; Fage, 1960; Lewis & Fish, 1969; Shih, 1969; Moore & Sander, 1977, 1979; Gasca & Shih, 2001, 2003; Gasca & Suarez-Morales, 2004; Gasca, 2009a,b) and mainly from the upper layers (0-200m) of the water column. Globally, the ingolfiellideans have a very low biodiversity, with only 44

species. Jan Hendrik Stock explored extensively between 1973 and 1978 more than 25 islands in the Antilles in search of stygobionts, which resulted in a few new descriptions (Stock, 1976, 1977a,b, 1979). More recently, since the late 1960s, several authors have provided important publications on Caribbean amphipods, including descriptions of new species, re-descriptions or additional information, taxonomic lists and inventories, biogeographical and ecological data, and illustrated keys for specific taxa (see Table 1 for a comprehensive reference list by major topics).

In this paper, we analyzed the amphipod biodiversity (Caprellidea, Gammaridea, Hyperiid, and Ingolfiellidea) of the Caribbean Sea. Our goal was to carry out a spatial analysis of the data that allowed us to determine gaps in knowledge and to test if differences/similarities in composition between ecoregions are due to ecological and biogeographic factors or to geographic inequality in sampling efforts. We hope that this paper will contribute to raise awareness about the value of taxonomic inventories and to indicate how much and where scientific sampling is needed, for a better understanding of the Caribbean amphipod biodiversity.

## MATERIAL AND METHODS

**Study area:** The Caribbean Sea is a large sea, closed off to the West and to the South by the Americas, and bordered to the North and to the East by the islands chains of the Greater and Lesser Antilles (Spalding, 2004). It covers about 2754000km<sup>2</sup>, has a volume of nearly 6.5x10<sup>6</sup>km<sup>3</sup>, borders over 13500km of coastline, and is home to 12 island countries, nine continental countries, and 12 island dependencies (to France, The Netherlands, U.K., and U.S.). Towards the East and Northeast, the closely spaced chain of islands, banks, and sills of the Antilles Islands Arc separate the Caribbean from the Atlantic Ocean, and act as a sieve for the inflow of Atlantic water (Andrade & Barton, 2000), whereas towards the Northwest it is linked to the Gulf of Mexico by the Yucatan Channel (Miloslavich, Diaz, Klein,

TABLE 1  
Main topics developed in amphipod research in the Caribbean and bibliographical sources

CUADRO 1  
Principales tópicos de investigación sobre anfipodos en el Mar Caribe con sus fuentes bibliográficas

Topic	Taxonomic group (Suborder)	References
Species description	Caprelliidea	Ortiz, Guerra-García & Lalana (1998); Guerra-García (2003a); Guerra-García et al. (2006).
	Gammaridea	Stebbing (1888, 1897, 1903, 1906, 1908); Chevreux (1901); Shoemaker (1926, 1933a,b, 1934, 1948); Stephensen (1933, 1947, 1948); Ruffo (1950, 1954); Barnard (1954, 1960, 1961, 1962, 1964, 1972, 1973); Mateus & Mateus (1966); Holsinger & Peck (1968); Myers (1968, 1977, 1978, 1979); Holsinger (1977, 1990, 1992); Ortiz (1976a,b,c,d,e, 1980,1991); Just (1977, 1983, 1984); Stock (1977a,b, 1978, 1980, 1981, 1983, 1985); McKinney (1978, 1979, 1980); McKinney et al. (1978); Ortiz & Gómez (1979); Ortiz & Lalana (1980, 1989, 1994a,b, 1995, 2002a,b,c); Thomas (1983, 1997); Thomas & Barnard (1983, 1985, 1986, 1989, 1990, 1991a,b); Ortiz & Nazábal (1984a,b, 1988); Ortiz & Veleo (1985); Barnard & Agard (1986); Barnard & Thomas (1987, 1990); Vonk & Stock (1987); Vonk (1988, 1989, 1990, 1991); Ciavatti (1989); Bellan-Santini (1990, 1997); Ortiz, Lalana & López (1992); Ortiz, Lalana & Beltran (1993); LeCroy (1995); Ortiz & Pérez (1995); Lowry & Stoddart (1997); Ortiz & Lemaitre (1997); Jaume & Wagner (1998); Ortiz, Lalana & Lio (1999); Krapp-Schickel & Ruffo (2000, 2001); Martín, Ortiz & Atienza (2000, 2001); Ortiz, Lalana & Sánchez (2000); Ortiz, Martín & Atienza (2000); Ruffo et al. (2000); Berge et al. (2001); Lazo-Wasem & Gable (2001); Ortiz, Cházaro-Olvera & Winfield (2001); Ortiz, García-Debras & Lalana (2002, 2003); Ortiz, Lalana & Varela (2002, 2004, 2006, 2007a,b,c, 2008); Van der Ham & Vonk (2003); Thomas & Klebba (2006, 2007); Ortiz & Thomas (2007); Coleman (2011); Ortiz, Varela & Lalana (2011).
	Ingolfiellidea	Stock (1976, 1979).
Inventories, taxonomic lists and reports	Caprelliidea	Díaz et al. (2005); Guerra-García et al. (2006).
	Gammaridea	Shoemaker (1935); McCain (1968); McCain & Steinberg (1970); McKinney (1977); Ortiz (1978, 1979a,b, 1983); Galan (1983, 1984); Stoner & Lewis (1985); Lagarde (1987); Ortiz & Lalana (1993, 1996, 1998, 2002a, 2006); Ortiz & Lemaitre (1994); Villaroel & Graciani (1997); Díaz & Martín (2000a,b,c,d, 2001a,b, 2003, 2005); Martín, Atienza & Díaz (2000); Atienza & Martín (2001); Díaz (2001); Lazo-Wasem & Gable (2001); Martín (2001a,b, 2003, 2007); Ortiz, Lalana & Lio (2001); Escobar-Briones et al. (2002); Martín, Ortiz & Díaz (2002); Ayala & Martín (2003); Martín & Díaz (2003); Van der Ham & Vonk (2003); Varela et al. (2003); Jara (2007); Ortiz, Lalana & Varela (2007c, d,e, 2008); Ortiz, Martín & Díaz (2007); LeCroy et al. (2009); López et al. (2009).
	Hypერიidea	Shoemaker (1945, 1948); Fage (1960); Lewis & Fish (1969); Shih (1969); Moore & Sander (1977, 1979); Ortiz, Lalana & Guevara (1990); Gasca & Shih (2001, 2003); Ortiz, Lalana, Varela & Leal (2003); Gasca & Suárez-Morales (2004); Ortiz, Lopeztegui, Lalana & Varela (2006); Gasca (2009a,b); LeCroy et al. (2009).
Biology, ecology and biogeography	Ingolfiellidea	Stock (1977b, 1979); Vonk & Schram (2003).
	Caprelliidea	Guerra-García et al. (2006).
	Gammaridea	Galan (1983, 1984); Sánchez (1985); Thomas (1997); Martín & Díaz (2003); Díaz & Martín (2005); Jara (2007).
	Hypერიidea	Lewis & Fish (1969); Moore & Sander (1977, 1979).
	Ingolfiellidea	Stock (1977b, 1979); Vonk & Schram (2003).
Taxonomic keys	Caprelliidea	Díaz et al. (2005); Guerra-García et al. (2006).
	Gammaridea	Galan (1983,1984); LeCroy (1995); Thomas (1997); Ortiz, Alvarez & Winfield (2002); Ortiz, Martín, Winfield, Díaz & Atienza (2004).

Alvarado, Díaz, Gobin, Escobar-Briones, Cortés, Weil, Bastidas, Robertson, Zapata, Martín, Kazandjan & Ortiz, 2010).

We compiled and analyzed the amphipod species composition of the Caribbean region from the Southeastern coast of the Yucatan Peninsula in Mexico (Quintana Roo State) to Belize, Guatemala, Honduras, Nicaragua, Costa Rica, and Panama down to Colombia and Venezuela in South America, and also along the Greater Antilles (Cuba, the Cayman Islands, Jamaica, Hispaniola, Navassa and Puerto Rico), the Lesser Antilles (St. Thomas, Tortola, Anegada, Jost Van Dyke, Virgin Gorda, St. Croix, Anguilla, St. Martin, St. Barthelemy, Saba, St. Eustatius, Barbuda, St. Kitts, Nevis, Antigua, Monserrat, Guadeloupe, Dominica, Martinique, St. Lucia, St. Vincent, Grenadines and Grenada), down to the Southern islands of Barbados, Tobago, Trinidad and the Netherlands Antilles (Bonaire, Curaçao, Aruba).

**Diversity inventories:** To compile the available data on marine Amphipoda species diversity in the Caribbean, we used two approaches. The first approach summarized the number of marine amphipod species for Caprellidea, Gammaridea, Hyperidea and Ingolfiellidea using georeferenced species records from the open-access databases (WoRMS and OBIS) and from local, country, and regional checklists. The second approach was to produce revised species lists for the group by country (where information was available).

**Data analysis:** The presence/absence data for each Amphipoda taxon was used to calculate a Sørensen distance matrix. This symmetrical index excludes double absences in the calculation. Only countries with 10 or more amphipod species were included in the matrix. To analyze the relative contribution of each country to the Caribbean regional diversity, we used the contribution partition analysis proposed by Lu, Wagner & Chen (2007), using species richness as a measure of diversity. The gamma (regional) diversity is the sum of the alpha (local) and beta (interlocal) diversities.

Using these estimates, we calculated the index of relative contribution for each country where the larger the number of species listed, the higher the alpha diversity value. Based on the number of endemic species by country, the relevance of this country to the relative contribution to the gamma diversity changed.

Additionally, in order to determine the rate of description of new species, we arranged the number of species records per year and produced cumulative curves for all suborders. Based on the fact that the quality of taxonomic inventories depends on the availability of identification guides and taxonomic experts, our review also related these sources to each higher taxon.

## RESULTS

**Diversity and distribution:** There are 535 species identified within 236 genera and 73 families that have been reported for the Caribbean Sea. The most speciose taxon are the Gammaridea with 387 species (72.34%), followed by the Hyperidea with 116 species (21.68%), the Caprellidea with 25 (4.67%) and the Ingolfiellidea with 7 (1.31%) species. A complete taxonomic listing of these species and their distributions within the Caribbean is presented in a supplementary table (Appendix 1).

Regarding the regional distribution of these species, each taxon apparently exhibits its own spatial pattern of species richness, as no common distribution pattern was evident (Table 1 and 2). The Western Caribbean ecoregion had the largest diversity (282 species), followed by the Southern Caribbean (230 species), while the Eastern Caribbean ecoregion had the lowest diversity (73 species). When examined by country, Mexico and Venezuela were the most species-rich with 266 and 206 species respectively, followed by Cuba with 168 species. The species number of the three countries combined accounted for 49.72% of the total amphipod species reported for the Caribbean. Twelve countries had less than 50 species (Table 2). When the richness was standardized by coastline length, the average number of

species per 100 kilometers of country-coastline was  $5 \pm 8$  species/100km, with Mexico having the highest richness (29.4 species/100km), and Nicaragua the lowest (0.2 species/100km). When standardized by the Economic Exclusive Zone (EEZ) coverage, the average number of species was  $0.05 \pm 0.08$  species/100km<sup>2</sup> of EEZ, with Mexico having the highest number of species (0.30 species/100km<sup>2</sup>) of EEZ, followed by Costa Rica (0.12 species/100km<sup>2</sup>), and Nicaragua having the lowest (0.001 species/100km<sup>2</sup>).

The Gammaridea were clearly the most speciose taxon in all countries and ecoregions. The Southern Caribbean ecoregion (represented by Venezuela, Aruba, Curaçao, Bonaire, Trinidad and Tobago) encompassed the largest gammaridean diversity (188 species), while the Eastern Caribbean ecoregion (Lesser Antilles) had the lowest diversity (54 species) (Table 2). When examined by country, Venezuela had the highest diversity with 169 species, followed by Mexico and Cuba, with 140 and

TABLE 2  
Number of Caribbean Amphipoda species per country within the five ecoregions

CUADRO 2  
Riqueza de especies del orden Amphipoda por país en cada ecorregión del Mar Caribe

Ecoregion	Amphipoda Suborders				Total species	Coastline (km)*	No. spp./100km of coastline	EEZ (km <sup>2</sup> )*	spp./100km <sup>2</sup> EEZ
	Caprellidea	Gammaridea	Hyperidea	Ingolfiellidea					
WESTERN CARIBBEAN	13	156	112	1	282				
México (Yucatán)	13	140	112	1	266	905	29.4	88 874	0.30
Belize	0	37	0	0	37	516	7.2	35 351	0.10
Guatemala	0	1	0	0	1	150	0.7	1 642	0.06
Honduras	0	2	0	0	2	644	0.3	18 151	0.01
SOUTH-WESTERN CARIBBEAN	9	111	0	0	120				
Nicaragua	0	1	0	0	1	503	0.2	87 930	0.001
Costa Rica	0	31	0	0	31	212	15	25 090	0.12
Panama	0	33	0	0	33	1 295	2.5	143 442	0.02
Colombia	9	67	0	0	67	1 760	3.8	472 891	0.01
SOUTHERN CARIBBEAN	12	188	24	6	230				
ABC*	0	22	0	5	27				
Venezuela	12	169	24	1	206	2 800	7.4	470 666	0.04
Trinidad & Tobago	0	8	0	0	8	362	2.2	74 199	0.01
GREATER ANTILLES	11	177	28	0	216				
Cuba	10	131	27	0	168	3 755	4.5	350 751	0.05
Cayman Islands	0	4	0	0	4	160	2.5		
Jamaica	0	5	3	0	8	1 022	0.8	258 137	0.003
Hispaniola	0	18	0	0	18	3 051	0.6	382 658	0.004
Puerto Rico	1	30	0	0	31	501	6.2	205 529	0.02
EASTERN CARIBBEAN	3	54	16	0	73				
Lesser Antilles	3	54	16	0	73				

(ABC\*=Aruba, Bonaire and Curacao and\*=Coastline information taken from The world fact book (<http://www.cia.gov/library/publications/the-world-factbook/geos>) and Economic Exclusive Zone (EEZ) information was taken from [www.seaaroundus.org/eez/eez.aspx](http://www.seaaroundus.org/eez/eez.aspx))



131 species respectively, while Nicaragua and Guatemala had the lowest with only one species each. Gammaridea are composed of 49 families (Table 3), of which the Melitidae and Hadziidae were the most diverse, with 57 and 31 species, respectively, while 10 families were only represented by one species (Appendix 1). The most eurytopic species within the region were *Parahyale hawaiiensis* (eight countries), *Ampelisca lobata* and *P. fascigera* (seven countries); in contrast 208 of the 387 species of Gammaridea were reported as unique to one country. The genus *Ampelisca* was the most speciose, with 23 species.

The highest diversity was recorded in Caprellidea for the Western Caribbean ecoregion (13 species), the lowest occurred in the Eastern Caribbean (three species) (Table 2). Mexico displayed the highest diversity values, with 13 species, representing 52.0% of reported for the Caribbean, followed by Venezuela with a total of 12 species and Cuba with 10 species. This taxon, composed of five families (Table 3), includes the Caprellidae and Pariambidae, the most diverse taxa, with ten and seven species respectively (Appendix 1). In the Caribbean, the genus *Caprella* and *Deutella* were the most speciose with five species reported, respectively. The most eurytopic species were *Paracaprella pusilla*, *Deutella incerta* and *Hemiaegina minuta*, represented in four countries, respectively.

The highest species richness (112 species) in the Hyperidea occurred in the Western Caribbean ecoregion, mainly from Yucatan in

the Mexican Caribbean where pelagic samples have frequently been obtained (Table 2). Species of this taxon were reported only in five of the 17 countries analyzed herein. In the Caribbean Sea, the hyperiids are represented by 19 families (Table 3), of which the Scinidae holds 17 species, followed by the Lestrigonidae and Platyscelidae with 14 species each (Appendix 1). The genus *Scina* is the most diverse with 16 species. The most eurytopic species is *Lestrigonus bengalensis*, occurring in five countries, in contrast to 70 of the 116 species in this suborder reported in one country.

For Ingolfiellidea, only seven species within one genus and one family were reported for the Caribbean Sea (Table 3, Appendix 1). Of these species, six were recorded in the Southern Caribbean ecoregion (Aruba, Curaçao, Bonaire and Venezuela) and the remaining species in the Western Caribbean (Mexico, Yucatan) (Table 2), and all were reported only for one country.

**Advances in regional knowledge:** The first amphipod descriptions were made by Linnaeus in 1758 (*Gammarus locusta*, *Gammarus pulex*, and *Cyamus ceti*). Between 1800 and 1850, an average of 6.28 species were described per year and then, between 1850 and 1900, this number increased to 44.64 new descriptions per year, and to 51.04 between 1850 and 1900. Between 1900 and 1950, the number of descriptions almost doubled to 97.70 species per year, and since 1950, about 103.18 species are described per year (Fig. 1A). The accumulation curve of world species discoveries (Fig. 1B) shows no sign of leveling off, indicating that a full inventory of these taxa is still far from being completed despite two centuries of efforts. In the Caribbean region, the description of species began in 1888 with *Ampelisca abyssicola* by Stebbing and increased at a slow rate changing in trend with a substantial accumulation of species in the 1980s (Fig. 2). As in the global graph, species description in the Caribbean region is far from reaching an asymptote, and at present, 218 species of amphipods (40.75% of the Caribbean

TABLE 3  
Species, genera and families of Caribbean amphipods


CUADRO 3  
Número de especies, generos y familias de los anfipodos presentes en el Mar Caribe

Suborder	Family	Genus	Species
Caprellidea	5	2	25
Gammaridea	49	179	387
Hyperidea	18	44	116
Ingolfiellidea	1	1	7
Total	73	236	535


total) have been described from this region. Of these, 60.55% (132 species) are endemic to the Caribbean region, while the remaining 39.45% (86 species) have been reported elsewhere, mainly for the Gulf of Mexico and Florida.

**Similarity analysis:** In the cluster tree generated with the Sorensen Distance Index, the amphipod fauna of Cuba, Mexico and

Venezuela was aggregated into one group (Fig. 3). The similarity between Mexico and Venezuela is based on 119 common species (22.24% of the total species in the region), while the similarity between Cuba and Mexico is based in 99 common species (18.50% of the total species in the region; Table 4). Mexico, Cuba and Venezuela share species with other countries in the region (14, 13 and 10 countries,



**Fig. 1.** (A) Amphipoda species discoveries around the world; (B) Cumulative Amphipoda species discoveries around the world.



**Fig. 2.** (A) Amphipoda species discoveries in the Caribbean Sea; (B). Cumulative Amphipoda species discoveries in the Caribbean Sea.

respectively). Countries with the highest alpha diversity were Mexico, Venezuela, and Cuba and those contributed proportionally to the gamma diversity. The Lesser Antilles contributed to regional diversity, despite having fewer species than i.e. Colombia. Aruba, Curaçao and Bonaire contributed more to the regional diversity than countries with higher alpha

diversity values, i.e. Puerto Rico, Costa Rica, and Belize (Fig. 4).

## DISCUSSION

The crustaceans, along with the mollusks, are the most speciose taxonomic groups among the marine fauna, with 44 950 reported species




(Bouchet, 2006). Despite the lack of a larval stage, and therefore, with limited dispersal capability, the order Amphipoda is a relatively species-rich group within the crustaceans, representing 15% of the total if we consider that the number of marine amphipod species is 6 950 (Vader, 2005), or up to 26% if we consider that the number of species is 11 780 (WoRMS, 2012).

In this sense, the 535 amphipod species reported in the Caribbean region represent a small contribution to global diversity. Despite this apparent low richness, large percentage of unique species, represent an important contribution to the endemic regional diversity. When comparing this amphipod diversity across geographical regions, the Caribbean has a larger diversity than the Patagonian Shelf (98 species) and the Tropical West Atlantic (107 species) (Miloslavich, Klein, Díaz, Hernández, Bigatti, Campos, Artigas, Castillo, Penchaszadeh, Neill, Carranza, Retana, Diaz de Astarloa, Lewis, Yorio, Piriz, Rodríguez, Yoneshigue-Valentin, Gamboa & Martín, 2011), but lower than Australia (888 species; Butler, Rees, Beesley & Bax, 2010), the South American Humboldt Current region (860 species; Miloslavich et al., 2011), the Southern Ocean (797 species;

De Broyer, Lowry, Jazdzewski & Robert, 2007; Zeidler & De Broyer, 2009), and Japan (544 species; Fujikura, Lindsay, Kitazato, Nishida & Shirayama, 2010). In order to establish management and conservation plans, hotspots of endemism should be identified, the species in need of protection prioritized, and their population size estimated. This should also lead to recommendations to upgrade their current conservation status. There is currently little published recognition that the diversity associated with the islands exists at all and as a result, not only its conservation has received little attention, but these species have often been affected by diverse, ongoing anthropogenic activities.

Gammaridea are the most diverse taxon among the Amphipoda, with at least 8 300 species according to WoRMS (2012). In the Caribbean, gammaridean amphipods were also the most diverse group (387 species), representing about 72% of the regional biodiversity. One of the most comprehensive reviews in marine gammaridean was carried out by Barnard & Karaman (1991) who provided not only keys, diagnoses and species lists for the group, but also discussed their geographic distribution based on thermal zones and depth, and some aspects of endemism. According to these authors, in terms of generic diversity, the Gammaridea are mainly a cold water group, whereas tropical regions seem to be poor in tropical endemic genera. A higher richness in cool and temperate environments, contrasting with poor diversity in the tropics, is also a general trend in freshwater amphipods (Vainola, Witt, Grabowski, Bradbury, Jazdzewski & Sket, 2008). In the tropics, even if many new species are expected to be described, it is unlikely that any of these ones will be associated with a new genus (Barnard & Karaman, 1991). In the Caribbean region, four new genera have been described since Barnard's and Karaman's work in 1991: *Ottenwalderia* (Jaume & Wagner, 1998), *Miramarassa* (Ortiz, Lalana & Lio, 1999) *Pleusiroides* and *Tantena* (Ortiz, Lalana & Varela, 2007a,b respectively). Other important components of this taxon in the Caribbean are the subterranean and cave



**Fig. 3.** Cluster tree based on species presence/absence matrix and generated with the Sørensen distance index. PRC Puerto Rico, LAN Lesser Antilles, HIS Hispaniola, ABC Aruba, Bonaire and Curacao, CRC Costa Rica, PAN Panama, BEL Belize, COL Colombia, CUB Cuba, MEX Mexico, VEN Venezuela.

taxa, whose present distribution and biogeographic pattern reflect their origin. In the case of marine and brackish species, it has been suggested that they may have derived from epigeal marine or brackish water ancestors as they adapted to fluctuating sea levels, and even some freshwater species may have derived from marine and brackish species that were caught inland during regression of seawater (Holsinger, 1994; Biernbaum, 1996), in both cases leading to speciation by isolation.

Caprellidea are a much less diverse taxon with 451 species worldwide and 25 species (4.72%) within 12 genera in the Caribbean (Table 1). Of these, more than 30% could be considered endemic for the region (*Pseudaeiginella colombiensis*, *Deutella caribensis*, *D. margaritae*, *D. mayeri*, *Mayerella redunca*,


*Metaprotella hummelincki*, *Tritella chibcha*, and *Cubadeutella cavernicola*), while about 40% of the species have an eurytopic distribution either in the tropical and the temperate marine ecosystems (*Caprella andreae*, *C. danilevskii*, *C. equilibra*, *C. penantis*, *C. scaura*, *Pseudaeiginella biscaynensis*, *Hemiaeigina minuta*, *Paracaprella pusilla* and *Phtisica marina*; McCain & Steinberg, 1970). In the case of *D. californica* and *Hemiproto wigleyi*, these species are distributed also in the East Pacific (Guerra-García, 2003a; Díaz, Guerra-García & Martín, 2005), while *D. incerta* is distributed along the East coast of the United States and Canada, and *Paracaprella digitimanus* has been also found on the Brazilian coast (Díaz et al., 2005). The Caribbean caprellid diversity could be considered high when

TABLE 4  
Species number of amphipods in common between countries in the Caribbean Sea

CUADRO 4  
Número de especies de anfípodos comunes entre cada uno de los países del Mar Caribe

	México	Puerto Rico	Hispaniola	Jamaica	Belize	Honduras	Guatemala	Nicaragua	Costa Rica	Panamá	Colombia	Venezuela	ABC*	Lesser Antilles	Trinidad & Tobago	Cayman Islands
Cuba	99	6	2	3	8	0	0	0	13	6	38	87	5	16	2	1
México		13	3	4	21	2	1	0	17	10	41	119	6	35	3	1
Puerto Rico			2	0	2	1	0	0	3	1	5	8	2	9	1	0
Hispaniola				1	0	0	0	0	0	0	2	3	2	3	0	0
Jamaica					0	0	0	0	0	0	0	5	0	2	0	0
Belize						1	0	0	7	7	7	16	0	3	2	0
Honduras							0	0	1	1	1	1	0	0	0	0
Guatemala								0	1	1	1	0	0	0	0	0
Nicaragua									0	0	0	0	0	0	0	0
Costa Rica										7	12	13	3	4	2	0
Panamá											5	7	2	0	0	0
Colombia												45	7	6	2	1
Venezuela													8	27	3	3
ABC*														4	1	1
Lesser Antilles															3	2
Trinidad & Tobago																0

(ABC\*=Aruba, Bonaire and Curacao).



**Fig. 4.** Contributions to the Amphipoda Caribbean gamma diversity by country. All countries are ordered by their alpha diversity value, from lowest to highest. HIS Hispaniola, PAN Panama, ABC Aruba, Bonaire and Curacao, BEL Belize, CRC Costa Rica, PRC Puerto Rico, LAN Lesser Antilles, COL Colombia, CUB Cuba, MEX Mexico, VEN Venezuela.

compared to other tropical areas [Brazil: 12 species (Wakabara, Tararam, Valério-Berardo, Duleba & Pereira-Leite, 1991; Guerra-García, 2003b); Tanzania: 13 species (Guerra-García, 2001); Mauritius: seven species (Guerra-García, 2003c); India: 11 species (Guerra-García, Ganesh, Jaikumar & Raman, 2010); Phuket, Thailand: 12 species (Guerra-García, 2004); Indonesia: seven species (Krapp-Schickel & Guerra-García, 2005); Philippines: three species (Guerra-García, 2002); Papua New-Guinea: seven species (Guerra-García, 2003d); Great Barrier Reef: 17 species (Guerra-García & Lowry, 2009), but still significantly lower when compared to temperate waters (Thiel, Guerra-García, Lancellotti & Vásquez, 2003). For example, 105 species have been reported for Japan (Takeuchi, 1999) and 41 species for the Mediterranean (Guerra-García & Takeuchi, 2002; Sturaro & Guerra-García, 2012).

The Hyperiidea have an overall number of valid species of 304 according to the WoRMS. There are 222 species known to occur in the Pacific Ocean and 175 species in the Atlantic (Gasca, 2009a,b). This review recorded 116 species in the Caribbean region, representing 38.16% of the global hyperiid fauna. The few inventories and species lists in the region have been provided not only for shallow water species (Gasca & Shih, 2001, 2003), but also for the deep water masses (Gasca, 2009b). One of the most complete studies on the Western Atlantic Hyperiidea, although restricted to the Southwestern Atlantic, reports 151 species in that region (Vinogradov, 1999). In comparison, the known species richness of hyperiid amphipods from the Caribbean region is larger than in the other adjacent tropical areas, such as the Gulf of Mexico (100 species; LeCroy et al., 2009), the Sargasso Sea (88 species;

Gasca, 2007), and Brazil (90 species; Montu, 1998; Lima & Valentin, 2001). We suggest that the number of hyperiid species inhabiting the Caribbean region is higher, mainly because the hyperiids can be divided roughly into warm-water and cold-water species, and most of the warm-water species are circumtropical (including the tropical and subtropical belts). Most of the hyperiid specimens contained in zooplankton collections have been identified at the suborder level, which leads us to recognize that new species and new records will occur as the taxonomic effort to identify these samples increases.

The ingolfiellidean amphipods are not very diverse in the Caribbean Sea (seven species), but they still represent an important percentage (15.91%) of the global total of 44 valid species. Such biodiversity is low, considering the wide range of ecological conditions in which they occur, which varies from deep sea soft muds, to high altitude freshwater rivers in the continents, to subterranean fresh, brackish and marine interstitial waters (Vonk & Schram, 2003), but comparable to other families of amphipods with partly overlapping habitat requirements such as hadziids (30 species) and bogdiellids (seven species). Seven ingolfiellidean amphipods are recorded for the Caribbean region, six of them are endemic and one, *Ingolfiella fuscina* is also recorded for the Gulf of Mexico and Florida (Felder & Camp, 2009). More exploration and collecting efforts within anchialine cave systems in the Caribbean islands will very likely lead to the discovering of a larger number of species in this suborder.

In general terms, Mexico, Venezuela and Cuba exhibit the highest species diversity, due in part to the higher intensity of research in these areas. In relation to species descriptions and taxonomic publications, Cuba has taken the lead, particularly in coastal and shallow waters. The least studied coasts in the Caribbean are those of Guatemala, Honduras, Nicaragua, Trinidad and Tobago, Jamaica and the Cayman Islands. The lack of knowledge in these countries is probably due to limitations in funding, research institutions, and/or

taxonomic expertise. Also, countries having a low reported biodiversity are probably undersampled, especially if they are in close proximity to highly diverse countries. For example, it is likely that the biodiversity of Belize, Colombia, Aruba, Bonaire, Curaçao and Trinidad and Tobago would resemble the biodiversity of their neighbouring countries Mexico and Venezuela. Another important aspect to consider is that a low number of species could also be related to a small size of the coastal zone, to the lack of habitat diversity, and/or to the level of degradation of the coast due to natural or anthropogenic disturbances (Dauvin & Ruellet, 2009; De la Ossa-Carretero, Del-Pilar-Ruso, Giménez-Casaldueiro, Sánchez-Lizaso & Dauvin, 2012). In addition, the number of species can be also related to the extent of the shelf area, the hydrographic isolation of bays, coves, among others.

In conclusion, available information on amphipod taxonomy in the Caribbean region is heterogeneous, restricted for the most part to Cuba, Mexico and Venezuela, and in these countries, mainly focused on Gammaridea and Caprellidea. Given this heterogeneity in the data, knowledge on the distributions of the species is also limited. There is a need to compare between habitats throughout the region, i.e. compare seagrass habitat diversity along a latitudinal gradient, or between depth zones, i.e. shelf, slope and abyssal marine ecosystems. This would lead us to understand distribution patterns and recognize hotspots that will allow for recommendations on management plans for each of them. On the other hand, knowledge of ecology and behaviour is virtually non-existent for the majority of the species. More significantly, at this time of rapidly vanishing natural habitats, we have not yet estimated how many species are relatively conserved within the boundaries of national parks and other protected areas. In addition to climate change which is related to higher temperatures, sealevel rise, erosion of the coastal habitats and the acidification and disappearance of important habitats, many intertidal zones are currently subject to disturbance from

industrialization and infrastructure development from the increasingly growing tourism industry. The fauna and flora of the intertidal zone are likely to be affected by such anthropogenic changes in environmental conditions; therefore there is a need for monitoring programmes, which could help document such environmental changes. Thomas (1993) stressed that gammaridean amphipods are ideal candidates for monitoring studies because they are ecologically and trophically important, numerically dominant, and have relative low dispersal capabilities. However, the use of these organisms as biomonitors in the tropics is only possible if their taxonomic and ecological information is available.

Besides increasing sampling efforts in coastal zones of poorly sampled countries in Central America and the Lesser Antilles, several ecosystems require additional sampling efforts like the anchialine cave systems, the continental margins, and the deep sea, including the escarpments, canyons, slopes, seamounts and diapirs, the abyssal plains, and geologically active ecosystems such as vents and seeps. These collecting initiatives should be combined with an increase in taxonomic efforts, both in classical taxonomy and in the use of molecular tools like barcoding, for which regional capacity building is required. All of these efforts should be carried out through regional coordinated collaboration.

#### ACKNOWLEDGMENTS

The authors thank the countless students who have worked at the Laboratory of Peracaridean Crustaceans, as well as in the amphipod collection of the Natural Science Museum at Simon Bolivar University.

#### RESUMEN

El orden Amphipoda es uno de los más diversos dentro de los Peracáridos. Las investigaciones sobre este orden en el Mar Caribe se iniciaron a finales de los años 1800, incrementándose significativamente a partir de 1980. En este estudio se analizó la biodiversidad de los anfípodos

(Caprellídea, Gammarídea, Hyperídea e Ingolfiellídea) en el Mar Caribe, compilándose los registros de especies citadas en las diversas ecorregiones del área, tanto en publicaciones como a través de bases de datos (WoRMS y OBIS). Se listan un total de 535 especies pertenecientes a 236 géneros y 73 familias. La ecorregión del Caribe Occidental presentó la mayor riqueza específica (282), mientras que el Caribe Oriental presentó el menor valor (73). México y Venezuela presentaron el mayor número de especies registradas (266 y 206, respectivamente), mientras que doce países presentaron menos de 50. Los Gammarídea fueron el grupo más diverso con 387 especies, seguido de los Hyperídea (116 especies). Hasta el presente, se han descrito 218 especies nuevas en aguas del Caribe, de las cuales 132 son endémicas. Las áreas con mayor diversidad de anfípodos se concentraron a lo largo del Caribe Mexicano, Cuba y la costa norte de Sur-América (Venezuela-Colombia); sin embargo, este patrón refleja más bien un mayor esfuerzo en la recolección de estos organismos y una mayor experiencia taxonómica, que un patrón específico de distribución del grupo. En general, el conocimiento del grupo se limita a las aguas someras, con muy poca información sobre zonas profundas. En este sentido, las prioridades de investigación regional en este grupo, debe enfocar un mayor esfuerzo en la costa de Centro América y las Antillas Mayores y Menores. Así mismo, es necesario aumentar las recolecciones en aguas profundas y otros ecosistemas particulares como los anquialinos. Por último, es importante aumentar las investigaciones taxonómicas y ecológicas de los taxones que presentan mayor diversidad.

**Palabras clave:** Amphipoda, diversidad, riqueza específica, Mar Caribe, endemismo.

#### REFERENCES

- Andrade, C. A. & Barton, E. D. (2000). Eddy development and motion in the Caribbean Sea. *Journal of Geophysical Research*, 105(C11), 26191-26201.
- Atienza, A. & Martín, A. (2001). *Tiburonnella viscana* (Amphipoda: Plastynopidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(3), 1270.
- Ayala, Y. & Martín, A. (2003). Ampliación del ámbito de *Podocerus kleidus* (Amphipoda: Podoceridae) al Mar Caribe. *Revista de Biología Tropical*, 51(1), 267.
- Barnard, J. L. (1954). Amphipoda of the family Ampeliscidae collected by the *Velero III* in the Caribbean Sea. *Allan Hancock Atlantic Expeditions Reports*, 7, 1-13.
- Barnard, J. L. (1960). The amphipod family Phoxocephalidae in the Eastern Pacific Ocean, with analysis of other species and notes for a revision of the family. *Allan Hancock Pacific Expeditions*, 18, 174-368.
- Barnard, J. L. (1961). Gammaridean Amphipoda from depths of 400 to 6000 meters. *Galathea Reports*, 5, 23-128.



- Barnard, J. L. (1962). South Atlantic abyssal amphipods collected by R.V. Vema. *Abyssal Crustacea. Vema Research Series, 1*, 1-78.
- Barnard, J. L. (1964). Deep-sea Amphipoda (Crustacea) collected by the R/V "Vema" in the eastern Pacific Ocean and the Caribbean and Mediterranean seas. *Bulletin of the American Museum of Natural History, 127*(1), 1-46.
- Barnard, J. L. (1969). The families and genera of marine Gammaridean Amphipoda. Smithsonian Institution. *Bulletin United State National Museum, 271*, 1-535.
- Barnard, J. L. (1972). A review of the family Synopiidae (=Tironidae), mainly distributed in the deep sea (Crustacea: Amphipoda). *Smithsonian Contributions to Zoology, 124*, 1-94.
- Barnard, J. L. (1973). Deep-sea Amphipoda of the genus *Lepechinella* (Crustacea). *Smithsonian Contributions to Zoology, 133*, 1-31.
- Barnard, J. L. & Agard, J. B. R. (1986). A new species of *Ampelisca* (Crustacea, Amphipoda) from Trinidad. *Bulletin of Marine Science, 39*(3), 630-639.
- Barnard, J. L. & Karaman, G. S. (1991). The Families and Genera of Marine Gammaridean Amphipoda (Except Marine Gammaroids). *Records of Australian Museum, 13* (Parts 1&2), 1-866.
- Barnard, J. L. & Thomas, J. D. (1987). A new species of *Chevalia* from the Caribbean Sea (Crustacea: Amphipoda). *Proceedings of Biological Society of Washington, 100*(3), 532-542.
- Barnard, J. L. & Thomas, J. D. (1990). *Ensayara jumane*, a new species from Belize, Caribbean Sea (Amphipoda, Lysianassidae). *Proceedings of Biological Society of Washington, 103*(1), 120-126.
- Bellan-Santini, D. (1990). New species of *Orchomene* s.l. (Crustacea, Amphipoda) of the abyssal benthos. Affinity with other deepwater *Orchomene* spp. *Beaufortia, 41*(3), 15-24.
- Bellan-Santini, D. (1997). Amphipods of the cold seep community on the South Barbados accretionary prism. *Crustaceana, 70*(1), 1-30.
- Bellan-Santini, D. (1998). Ecology. In S. Ruffo (Ed.), *The Amphipoda of the Mediterranean. Part 4. Localities and Map, Addenda to Parts 1-3, Key to Families, Ecology, Faunistics and Zoogeography, Bibliography, Index.* (815-959). Mónaco: Memoirs de L'Institut Oceanographique.
- Bellan-Santini, D. & Ruffo, S. (1998). Faunistics and Zoogeography. In S. Ruffo (Ed.), *The Amphipoda of the Mediterranean. Part 4. Localities and Map, Addenda to Pars 1-3, Key to Families, Ecology, Faunistics and Zoogeography, Bibliography, Index* (895-911). Mónaco, Memoirs de L'Institut Oceanographique.
- Berge, J., Vader, V. & Galan, A. (2001). Type material of Stegocephalidae Dana, 1855 (Crustacea, Amphipoda) in the collections of The Natural History Museum, London, including the description of seven new species. *Bulletin of Natural History Museum London (Zoology), 67*(2), 109-136.
- Biernbaum, C. K. (1996). Biogeography of coastal and anchialine amphipods of Ascension Island, South Atlantic Ocean. *Journal of Natural History, 30*(11), 1597-1615.
- Bouchet, P. (2006). The magnitude of marine biodiversity. In C. M. Duarte (Ed.), *The Exploration of Marine Biodiversity: Scientific and Technological Challenges.* (3162). Madrid, Spain: Fundación BBVA.
- Bousfield, E. L. (1973). *Shallow-water gammaridean amphipoda of New England.* London: Comstock Publishing Associates. Cornell University Press.
- Bousfield, E. L. (2001). An updated commentary on phylogenetic classification of the amphipod Crustacea and its application to the North American fauna. *Amphipacifica, 3*(1), 49-119.
- Butler, A. J., Rees, T., Beesley, P. & Bax, N. J. (2010). Marine Biodiversity in the Australian Region. *PLoS ONE, 5*(8), 1-15. doi:10.1371/journal.pone.0011831.
- Chevreaux, E. (1901). Crustaces amphipodes. Mission scientifique de M Ch. Alluaud aux Iles Seychelles (Mars, Avril, Mai, 1892). *Memoirs Societe Zoologique de France, 14*, 388-438.
- Ciavatti, G. (1989). Les talitrides (Crustacea, Amphipoda) des plages de la Guadeloupe, description de deux especes nouvelles. *Annales de l'Institut Oceanographique, Paris, 65*, 127-146.
- Coleman, C. O. (2011). *Batea aldebaranae* sp. nov. (Crustacea: Amphipoda: Bateidae) from the coast of Belize with redescrptions of two related species. *Journal of Marine Biology Association of the United Kingdom, 91*(2), 455-469. doi: 10.1017/S0025315410001207
- Dauvin, J. C. & Ruellet, T. (2009). The estuarine quality paradox: is it possible to define an ecological quality status for specific modified and naturally stressed estuarine ecosystems? *Marine Pollution Bulletin, 59*(1-3), 38-47.
- De-la-Ossa-Carretero, J. A., Del-Pilar-Ruso, Y., Giménez-Casaldueo, F., Sánchez-Lizaso, J. L. & Dauvin, J. C. (2012). Sensivity of amphipods to sewage pollution. *Estuarine Coastal and Shelf Science, 96*(1), 129-138.
- De Broyer, C., Lowry, J. K., Jazdzewski, K. & Robert, H. (2007). Part 1. Catalogue of the Gammaridean and Corophiidean Amphipoda (Crustacea) of the Southern Ocean, with distribution and ecological data. *Bulletin of Institute Royale of Science Nature, Belgium, Biology, 77*(1), 1-325.
- Díaz, Y. J. (2001). *Contribución al conocimiento del orden Amphipoda de la costa Centro-Occidental de Venezuela.* (Tesis inédita de maestría). Universidad Simón Bolívar, Venezuela.

- Díaz, Y. J. & Martín, A. (2000a). *Audulla chelifera* (Amphipoda: Corophiidae) en las costas de Venezuela. *Revista de Biología Tropical*, 48(2-3), 722.
- Díaz, Y. J. & Martín, A. (2000b). *Nasageneia yucatanensis* (Amphipoda: Eusiridae) en las costas de Venezuela. *Revista de Biología Tropical*, 48(2-3), 724.
- Díaz, Y. J. & Martín, A. (2000c). *Eusiroides yucatanensis* (Amphipoda: Eusiridae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(2), 767.
- Díaz, Y. J. & Martín, A. (2000d). *Listriella carinata* (Amphipoda: Liljeborgiidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(2), 767.
- Díaz, Y. J. & Martín, A. (2001a). *Amphilochus casahoya* (Amphipoda: Amphilochidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(3-4), 1269.
- Díaz, Y. J. & Martín, A. (2001b). New records of amphipods (Crustacea: Amphipoda) from shallow water on the Caribbean coast of Venezuela. *Revista de Biología Tropical*, 49(3-4), 1271-1276.
- Díaz, Y. J. & Martín, A. (2003). Primer registro de *Tropichelura gomezi* (Amphipoda: Cheluridae) para el Caribe Sur. *Revista de Biología Tropical*, 51(1), 266.
- Díaz, Y. J. & Martín, A. (2005). Crustáceos anfípodos asociados a praderas de *Thalassia testudinum* del Parque Nacional Morrocoy, Venezuela. In D. Bone (Ed.), *Estudio integral del sistema Parque Nacional Morrocoy con vías al desarrollo de planes de uso y gestión para su conservación. Informe Final*. (572-612). Caracas: USB-UCV-UNEFM-INIA-FONACIT.
- Díaz, Y. J., Guerra-García, J. M. & Martín, A. (2005). Caprellids (Crustacea: Amphipoda: Caprellidae) from shallow waters of the Caribbean coast of Venezuela. *Organisms Diversity & Evolution*, 5(10), 1-25.
- Escobar-Briones, E., Winfield, I., Ortíz, M., Gasca, R. & Suárez, E. (2002). Amphipoda. In J. Llorente & J. J. Morrone (Eds.), *Biodiversidad, taxonomía y biogeografía de artrópodos de México: Hacia una síntesis de su conocimiento. Volumen III*. (341-371). México: Comisión Nacional para el Conocimiento y uso de la Biodiversidad. Universidad Nacional Autónoma de México.
- Fage, L. (1960). Oxycephalidae. Amphipodes pélagiques. *Dana Reports*, 52, 1-145.
- Felder, D. L. & Camp, D. K. (2009). *Gulf of Mexico Origin, waters, and biota. Volume 1, Biodiversity*. USA: Texas A&M University Press.
- Foster, J. M., LeCroy, S. E., Heard, R. W. & Vargas, R. (2009). Gammariden amphipods. En Wehrmann, I.S. & Cortés, J. (Eds.), *Marine Biodiversity of Costa Rica, Central America. Series: Monographiae Biologicae*. (265-274). Costa Rica: Springer Netherlands.
- Fujikura, K., Lindsay, D., Kitazato, H., Nishida, S. & Shirayama, Y. (2010). Marine Biodiversity in Japanese Waters. *PLoS ONE*, 5(8), 1-13. doi: 10.1371/journal.pone.0011836
- Galan, A. (1983). Systematic studies on *Erichtonius brasiliensis* (Crustacea, Amphipoda, Ischyroceridae) from the Caribbean Sea of Venezuela. *Boletín del Instituto Oceanográfico Universidad de Oriente*, 22, 65-69.
- Galan, A. (1984). *A systematic study of Amphipoda (Crustacea) of the Caribbean coast of Venezuela*. (Doctoral thesis). Imperial College of Science and Technology, London University.
- Gasca, R. (2007). Hyperiid amphipods of the Sargasso Sea. *Bulletin of Marine Science*, 81(1), 115-125.
- Gasca, R. (2009a). Hyperiid Amphipods. In I. S. Wehrmann, & J. Cortés (Eds.), *Marine Biodiversity of Costa Rica, Central America. Series: Monographiae Biologicae* (275-282). Netherlands, Springer.
- Gasca, R. (2009b). Diversity of Hyperiid Amphipods (Crustacea: Peracarida) in the Western Caribbean Sea: news from the deep. *Zoological Studies*, 48(1), 63-70.
- Gasca, R. & Shih, C-t. (2001). Hyperiid amphipods of surface waters of the western Caribbean Sea. *Crustaceana*, 74(5), 489-499.
- Gasca, R. & Shih, C-t. (2003). Hyperiid Amphipods of Banco Chinchorro. *Bulletin of Marine Science*, 73, 91-98.
- Gasca, R. & Suárez-Morales, E. (2004). Distribution and abundance of Hyperiid Amphipods (Crustacea: Peracarida) of the Mexican Caribbean Sea, (August 1986). *Caribbean Journal of Science*, 40(1), 23-30.
- Guerra-García, J. M. (2001). The Caprellidea (Crustacea: Amphipoda) collected by the expedition of "Grigore Antipa" National Museum of Natural History from Tanzania, with the description of a new genus and two new species. *Travaux Du Museum d'Histoire Naturelle "Grigore Antipa"*, 43, 23-45.
- Guerra-García, J. M. (2002). Littoral caprellids (Crustacea: Amphipoda: Caprellidea) from Philippines, with the description of a new species. *The Raffles Bulletin of Zoology*, 50, 395-406.
- Guerra-García, J. M. (2003a). Revision of the genus *Deutella* (Crustacea: Amphipoda: Caprellidea) with description of a new species, redescription of *Deutella venenosa* Mayer, 1890 and a key to the species of *Deutella*. *Journal of Natural History*, 37, 1059-1084.
- Guerra-García, J. M. (2003b). Two new species of deep-water caprellids (Crustacea: Amphipoda) from north-eastern Brazil. *Cahiers de Biologie Marine*, 44, 171-184.
- Guerra-García, J. M. (2003c). The Caprellidea (Crustacea: Amphipoda) from Mauritius Island, Western Indian Ocean. *Zootaxa*, 232, 1-24.



- Guerra-García, J. M. (2003d). Caprellids (Crustacea: Amphipoda) from Papua New Guinea, with the description of a new species. *Helgolaender Marine Research*, 57, 100-109.
- Guerra-García, J. M. (2004). Littoral Caprellidea (Crustacea: Amphipoda) from Phuket, Thailand. *Steenstrupia*, 28(2), 159-175.
- Guerra-García, J. M. & Takeuchi, I. (2002). The Caprellidea (Crustacea: Amphipoda) from Ceuta, North Africa, with the description of three species of *Caprella*, a key to the species of *Caprella*, and biogeographical discussion. *Journal of Natural History*, 36(6), 675-713.
- Guerra-García, J. M. & Lowry, J. (2009). Caprellidae. *Zootaxa*, 2260, 290-327.
- Guerra-García, J. M., Krapp-Schickel, T. & Müller, H. G. (2006). Caprellids from the Caribbean coast of Colombia, with description of three new species and a key for species identification. *Boletín de Investigaciones Marino Costeras*, 35(1), 149-194.
- Guerra-García, J. M., Ganesh, T., Jaikumar, M. & Raman, A. V. (2010). Caprellids (Crustacea: Amphipoda) from India. *Helgolaender Marine Research*, 64(4), 297-310.
- Holsinger, J. R. (1977). A new genus and two new species of subterranean amphipod crustaceans (Gammaridae S. Lat.) from Yucatan Peninsula in Mexico. *Association for Mexican Cave Studies Bulletin*, 6, 15-25.
- Holsinger, J. R. (1990). *Tuluweckelia cernua*, a new genus and species of stygobiont amphipod crustaceans (Hadziidae) from anchialine caves on the Yucatan peninsula in Mexico. *Beaufortia*, 41(14), 97-107.
- Holsinger, J. R. (1992). Two new species of the subterranean amphipod genus *Bahadzia* (Hadziidae) from the Yucatan Peninsula region of southern Mexico, with an analysis of phylogeny and biogeography. *Stygologia*, 7, 85-105.
- Holsinger, J. R. (1994). Pattern and process in the biogeography of subterranean amphipods. *Hydrobiologia*, 287(1), 131-145. doi: 10.1007/BF00006902.
- Holsinger, J. R. & Peck, S. B. (1968). A new genus and species of subterranean amphipod (Gammaridea) from Puerto Rico, with notes on its ecology, evolution and relationship to other Caribbean amphipods. *Crustaceana*, 15(3), 249-262.
- Jara, A. C. (2007). *Estructura de la comunidad de macrocrustáceos bentónicos de las áreas someras (0 - 20 m) y profundas (80-170 m) de las islas de Providencia y Santa Catalina presentes durante la época seca del 2005*. (Tesis de maestría). Universidad de Bogotá Jorge Tadeo Lozano, Bogotá, Colombia.
- Jaume, D. & Wagner, H. P. (1998). New cave-dwelling amphipods (Lysianassidae, Hadziidae) from the Dominican Republic (Hispaniola). *Contributions to Zoology*, 68(1), 37-66.
- Just, J. (1977). *Amphylodomus incurvaria* gen. et sp.n. (Crustacea, Amphipoda), a remarkable leaf-cutting amphitoid from the marine shallows of Barbados. *Zoologica Scripta*, 6(3), 229-232.
- Just, J. (1983). Siphonoecetinae subfam. n. (Crustacea, Amphipoda, Corophiidae) 1: Classification. *Steenstrupia*, 9, 117-135.
- Just, J. (1984). Siphonoecetinae (Crustacea, Amphipoda, Corophiidae) 2: *Caribboecetes* Just, 1983, with description of six new species. *Steenstrupia*, 10, 37-64.
- Krapp-Schickel, T. & Ruffo, S. (2000). The *Maera quadrimana* – complex (Crustacea Amphipoda, Melitidae) demands a new concept: *Quadrimaera* n. gen. (with description of three new species from Western Atlantic). *Bollettino del Museo Civico di Storia Naturale di Verona*, 24, 193-214.
- Krapp-Schickel, T. & Ruffo, S. (2001). A new *Maeracoota* from the Caribbean Sea: *Maeracoota galani* n. sp. (Crustacea Amphipoda, Melitidae). *Bollettino del Museo Civico di Storia Naturale di Verona*, 25, 3-8.
- Krapp-Schickel, T. & Guerra-García, J. M. (2005). Littoral Caprellidae (Crustacea: Amphipoda) from Indonesia, with the description of a new species. *Bollettino del Museo Civico di Storia Naturale di Verona*, 29, 47-62.
- Lagarde, G. (1987). Anfipodos gammaridea del litoral de Golfo Triste y áreas adyacentes. *Caribbean Journal of Science*, 23(2), 260-277.
- Lazo-Wasem, E. A. & Gable, M. F. (2001). A revisión of *Parhyalella* Kunkel (Crustacea: Amphipoda: Gammaridea). *Bulletin of Peabody Museum of Natural History*, 46, 1-80.
- LeCroy, S.E. (1995). Amphipoda Crustacea III. Family Colomastigidae. *Memoirs of Hourglass Cruises*, IX (Part II), 1-139.
- LeCroy, S. E., Gasca, R., Winfield, I., Ortiz, M. & Escobar-Briones, E. (2009). Amphipoda (Crustacea) of the Gulf of Mexico. In D. L. Felder & D. K. Camp (Eds.), *Gulf of Mexico. Origin, Waters, and Biota. Vol. 1, Biodiversity*. (941-972). Texas, A&M University Press, College Station.
- Lewis, J. B. & Fish, A. G. (1969). Seasonal variation of the zooplankton fauna of surface waters entering the Caribbean Sea at Barbados. *Caribbean Journal of Science*, 9(1-2), 1-24.
- Lima, M. C. G. & Valentin, J. L. (2001). Preliminary results to the holistic knowledge of the Amphipoda Hyperidea faunal composition off the Brazilian coast. *Journal of Plankton Research*, 23(5), 469-480.
- Lincoln, R. J. (1979). *British Marine Amphipoda: Gammaridea*. London: British Museum (Natural History).

- Linnaeus, C. (1758). *Systema Naturae*. Editio Decima. Tomus I. Holmiae [Stockholm]: Laurentii Salvii.
- López, S., Díaz, Y. J. & Martín, A. (2009). Anfipodos (Crustacea: Gammaridea) de la isla de Cubagua, Venezuela. *Boletín del Instituto Oceanográfico de Venezuela*, 48(1), 43-51.
- Lowry, J. K. & Stoddart, H. E. (1997). Amphipoda Crustacea IV. Families Aristiidae, Cyphocarididae, Endeavouridae, Lysianassidae, Scopelocheiridae, Uristidae. *Memoirs of Hourglass Cruises*, 10(Part I), 1-148.
- Lu, H. P., Wagner, H. H. & Chen, X. Y. (2007). A contribution diversity approach to evaluate species diversity. *Basic and Applied Ecology*, 8(1), 1-12.
- Martín, A. (2001a). *Westwoodilla longimana* (Amphipoda: Oedicerotidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(3-4), 1270.
- Martín, A. (2001b). Ampliación del ámbito de *Seba tropica* (Amphipoda: Sebidae) en las costas de Venezuela. *Revista de Biología Tropical*, 49(3-4), 1270.
- Martín, A. (2003). Ampliación del ámbito de *Bemlos kunkelae* (Amphipoda: Aoridae) en el Mar Caribe. *Revista de Biología Tropical*, 51(1), 267.
- Martín, A. (2007). *Catálogo de los anfipodos (Crustacea: Amphipoda: Gammaridea y Caprellidae) de Venezuela*. Tomos I y II. (Trabajo de ascenso Titular). Universidad Simón Bolívar, Caracas, Venezuela.
- Martín, A. & Díaz, Y. J. (2003). La fauna de anfipodos (Crustacea: Amphipoda) de las aguas costeras de la región oriental de Venezuela. *Boletín del Instituto Español de Oceanografía*, 19(1-4), 327-344.
- Martín, A., Atienza, D. & Díaz, Y. (2000). Ampliación de ámbito de *Melita persona* (Amphipoda: Melitidae) en el Caribe Sur. *Revista de Biología Tropical*, 49(1), 397.
- Martín, A., Ortiz, M. & Atienza, D. (2000). Una nueva especie de anfipodo del género *Listriella* (Crustacea, Amphipoda, Liljeborgiidae) de Venezuela. *Boletín del Centro de Investigaciones Biológicas, Universidad del Zulia*, 34(3), 399-409.
- Martín, A., Ortiz, M. & Atienza, D. (2001). Una nueva especie de anfipodo del género *Amphilochus* (Crustacea, Amphipoda, Amphilochidae) de Venezuela. *Acta Científica Venezolana*, 52, 34-39.
- Martín, A., Ortiz, M. & Díaz, Y. (2002). Nuevos registros de anfipodos (Crustacea: Peracarida) para Venezuela. *Boletín de Investigaciones Marino Costeras*, 31, 15-24.
- Martin, J. W. & Davis, G. E. (2001). *An updated classification of the recent Crustacea*. Science Series No. 39. Los Angeles: Natural History Museum of Los Angeles County.
- Mateus, A. & Mateus, E. de O. (1966). Amphipodes littoraux de Principe et de Sao Tome. *Annales de l'Institut Océanographique, Paris*, 44, 173-198.
- McCain, J. C. (1968). The Caprellidea (Crustacea, Amphipoda) of the Western North Atlantic. *Bulletin of the United States Natural Museum*, 278, 1-147.
- McCain, J. C. & Steinberg, J. E. (1970). Amphipoda-I, Caprellidea-I. In H. E. Gruner & L. B. Holthuis (Eds.). *Crustaceorum Catalogus 2* (1-78). Den Haag, W. Jung.
- McKinney, L. D. (1977). *The origin and distribution of shallow water gammaridean Amphipoda in the Gulf of Mexico and Caribbean Sea with notes on their ecology*. (Ph.D. Thesis). Texas A&M University, College Station, U.S.A.
- McKinney, L. D. (1978). Amphilochidae (Crustacea: Amphipoda) from the western Gulf of Mexico and Caribbean Sea. *Gulf Research Reports*, 6(2), 137-143.
- McKinney, L. D. (1979). Liljeborgiid amphipods from the Gulf of Mexico and Caribbean Sea. *Bulletin of Marine Science*, 29(2), 140-154.
- McKinney, L. D. (1980). Four new and unusual amphipods from the Gulf of Mexico and Caribbean Sea. *Proceeding of the Biological Society of Washington*, 93(1), 83-103.
- McKinney, L. D., Kalke, R. D. & Holland, J. S. (1978). New species of amphipods from the western Gulf of Mexico. *Contributions to Marine Science*, 21, 133-159.
- Miloslavich, P., Díaz, J. M., Klein, E., Alvarado, J. J., Díaz, C., Gobin, J., Escobar-Briones, E., Cortés, J., Weil, E., Bastidas, A. C., Robertson, R., Zapata, F. A., Martín, A., Kazandjan, A. & Ortiz, M. (2010). Marine biodiversity in the Caribbean: regional estimates and distribution patterns. *PLoS ONE*, 5(8), 1-25. doi: 10.1371/journal.pone.0011916.
- Miloslavich, P., Klein, E., Díaz, J. M., Hernández, C. E., Bigatti, G., Campos, L., Artigas, F., Castillo, J., Penchaszadeh, P. E., Neill, P. E., Carranza, A., Retana, M. V., Díaz de Astarloa, J. M., Lewis, M., Yorio, P., Piriz, M. L., Rodríguez, D., Yoneshigue-Valentin, Y., Gamboa, L., & Martín, A. (2011). Marine Biodiversity in the Atlantic and Pacific Coasts of South America: Knowledge and Gaps. *PLoS ONE*, 6(1), 1-43. doi: 10.1371/journal.pone.0014631.
- Montu, M. (1998). Malacostraca-Peracarida. Amphipoda-Hyperiidea. In P. S. Young (Ed.), *Catalogue of Crustacea of Brazil*. Série Livros 6. (595-603). Rio de Janeiro: Museu Nacional.
- Moore, E. A. & Sander, F. (1977). A study of the offshore zooplankton of the tropical western Atlantic near Barbados. *Ophelia*, 16(1), 77-96.

- Moore, E. A. & Sander, F. (1979). A comparative study of zooplankton from oceanic, shelf and harbor waters of Jamaica. *Biotropica*, 11(3), 196-206.
- Myers, A. A. (1968). Two Aoridae (Amphipoda, Gammaridea) including a new species of *Amphideutopus* Barnard from Venezuelan waters. *Crustaceana*, 14(2), 127-130.
- Myers, A. A. (1977). Studies on the genus *Lembos* Bate VI. Atlantic species: *L. dentischium* sp. nov., *L. kunkelae* sp. nov., *L. rectangulatus* sp. nov., *L. unicornis* Bynum & Fox. *Bollettino del Museo Civico di Storia Naturale di Verona*, 4, 125-154.
- Myers, A. A. (1978). Studies on the genus *Lembos* Bate VII. Atlantic species 4. *L. setosus* sp. nov., *L. brunneomaculatus* Myers ssp. *longicornis* nov. and ssp. *mackinneyi* nov., *L. foresti* Mateus and Mateus, *L. longicarpus* sp. nov. *Bollettino del Museo Civico di Storia Naturale di Verona*, 5, 183-209.
- Myers, A. A. (1979). Studies on the genus *Lembos* Bate VIII. Atlantic species 5. *L. tigrinus* sp. nov., *L. tempus* sp. nov., *L. spinicarpus* (Pearse) comb. nov. with ssp. *inermis* nov., *L. ovalipes* sp. nov., *L. unifasciatus* Myers ssp. *reductus* nov. *Bollettino del Museo Civico di Storia Naturale di Verona*, 6, 221-248.
- Myers, A. A. & Lowry, J. K. (2003). A phylogeny and a new classification of the Corophiidea Leach, 1814 (Amphipoda). *Journal of Crustacean Biology*, 23(2), 443-485.
- Ortiz, M. (1976a). Un nuevo anfípodo de aguas cubanas (Amphipoda, Gammaridea, Phliantidae). *Investigaciones Marinas*, 8(25), 21-35.
- Ortiz, M. (1976b). Un nuevo género y una nueva especie de anfípodo de aguas cubanas (Amphipoda, Gammaridea, Amphithoidae). *Ciencias (La Habana). Investigaciones Marinas*, 8(27), 3-12.
- Ortiz, M. (1976c). Un nuevo anfípodo (Amphipoda, Gammaridea, Gammaridea) colectado del contenido estomacal del ronco amarillo *Haemulon sciurus*, de aguas cubanas. *Investigaciones Marinas*, 8(27), 13-20.
- Ortiz, M. (1976d). Un nuevo anfípodo perforador de madera (Amphipoda, Gammaridea, Cheluridae) de aguas cubanas. *Investigaciones Marinas*, 8(27), 21-26.
- Ortiz, M. (1976e). A new crustacean amphipod, *Mallacoota carausui*, from the Cuban waters. *Revue Roumaine de Biologie, Série de Biologie Animale*, 21(2), 93-95.
- Ortiz, M. (1978). Invertebrados marinos bentónicos de Cuba. I. Crustacea, Amphipoda, Gammaridea. *Investigaciones Marinas*, 38, 3-10.
- Ortiz, M. (1979a). Lista de especies y bibliografía de los anfípodos (Crustacea: Amphipoda) del Mediterráneo Americano. *Investigaciones Marinas*, 43, 1-40.
- Ortiz, M. (1979b). Contribución al estudio de los anfípodos (Gammaridea) del Mediterráneo americano. *Investigaciones Marinas*, 45, 1-16.
- Ortiz, M. (1980). Una nueva especie de anfípodo bentónico (Amphipoda, Gammaridea) de aguas cubanas. *Revista de Investigaciones Marinas*, 1, 92-103.
- Ortiz, M. (1983). Los anfípodos (Gammaridea) de las costas del mar Caribe de la República de Colombia. *Revista de Investigaciones Marinas*, 4, 23-31.
- Ortiz, M. (1991). Amphipod Crustacea. II. Family Bateidae. *Memoirs of Hourglass Cruises*, VII, 1-31.
- Ortiz, M., Alvarez, F. & Winfield, I. (2002). *Caprellid Amphipods: Illustrated key for the genera and species from the Gulf of Mexico and the Caribbean Sea*. Mexico. Ediciones de la Universidad Nacional Autónoma de México.
- Ortiz, M., Cházaro-Olvera, S. & Winfield, I. (2001). A new amphipod crustacean of the genus *Haustorius* (Gammaridea, Haustoriidae), from coast of Mexico. *Avicennia*, 14, 53-59.
- Ortiz, M., García-Debras, A. & Lalana, R. (2002). Una nueva especie de anfípodo anquialina del género *Melita* (Gammaridea: Melitidae), de la Isla de Cuba. *Avicennia*, 15, 43-52.
- Ortiz, M., García-Debras, A. & Lalana, R. (2003). Una nueva especie de anfípodo anquialino del género *Socarnopsis* (Amphipoda, Lysianassidae) del sistema cavernario de Playa Girón, Cuba. *Avicennia*, 16, 71-77.
- Ortiz, M. & Gómez, O. (1979). Una nueva especie de anfípodo (Amphipoda, Gammaridea) de aguas profundas del sur de Cuba. *Investigaciones Marinas*, 40, 23-30.
- Ortiz, M., Guerra-García, J. M. & Lalana, R. (1998). *Cuba-deutella cavernicola*, a new genus and new species of Caprellidae (Crustacea: Amphipoda) from Cuba. *Zootaxa*, 2130, 60-68.
- Ortiz, M. & Lalana, R. (1980). Un nuevo anfípodo del género *Leptocheirus* (Amphipoda, Gammaridea) de aguas cubanas. *Revista de Investigaciones Marinas*, 1, 58-73.
- Ortiz, M. & Lalana, R. (1989). Una nueva especie de anfípodo del complejo *Eriopisa* (Amphipoda, Gammaridea), de aguas cubanas. *Revista de Investigaciones Marinas*, 10, 233-237.
- Ortiz, M. & Lalana, R. (1993). Adición a la lista de especies y bibliografía de los anfípodos (Crustacea, Amphipoda) del Mediterráneo Americano. *Revista de Investigaciones Marinas*, 14, 16-37.
- Ortiz, M. & Lalana, R. (1994a). A new amphipod species of the genus *Nasageneia* (Gammaridea: Eusiridae) from the south-western cuban marine waters. *Travaux Du Museum d'Histoire Naturelle "Grigore Antipa"*, 34, 285-292.

- Ortiz, M. & Lalana, R. (1994b). Two new species of the genus *Elasmopus* (Amphipoda, Gammaridea), from the Cuban marine waters. *Travaux Du Museum d'Histoire Naturelle "Grigore Antipa"*, 34, 293-302.
- Ortiz, M. & Lalana, R. (1995). *Gitanopsis templadoi*, una nueva especie de anfípodo (Gammaridea, Amphilo-chidae), de la plataforma suroccidental de Cuba. *Anales del Instituto de Ciencias Marinas y Limnología*, 22, 121-124.
- Ortiz, M. & Lalana, R. (1996). Los anfípodos de la primera expedición conjunta Cuba-US, a bordo del B/I "Ulises", a las aguas del Archipiélago Sabana-Camagüey, Cuba, en 1994. *Anales del Instituto de Biología, Universidad Nacional Autónoma de México Serie Zoológica*, 67, 89-101.
- Ortiz, M. & Lalana, R. (1998). Lista actualizada de los crustáceos no decápodos de Cuba. *Revista de Investigaciones Marinas*, 19(2-3), 92-99.
- Ortiz, M. & Lalana, R. (2002a). Una nueva especie de anfípodo del género *Spathiopus* (Gammaridea, Melitidae), de la plataforma noroccidental de Cuba. *Avicennia*, 15, 31-36.
- Ortiz, M. & Lalana, R. (2002b). Primer registro del género *Neoischyrocerus* (Amphipoda, Ischyroceridae) para el Mar Caribe y el archipiélago cubano, con la descripción de nueva especie de Cuba. *Avicennia*, 15, 37-51.
- Ortiz, M. & Lalana, R. (2002c). Una nueva especie de anfípodo del género *Bemlos* (Amphipoda, Gammaridea, Aoridae), del archipiélago cubano. *Revista de Investigaciones Marinas*, 23(1), 69-76.
- Ortiz, M. & Lalana, R. (2006). Nuevos datos sobre crustáceos peracáridos (Crustacea, Peracarida) de las costas del Caribe de Costa Rica. *Cocuyo*, 16, 16-18.
- Ortiz, M., Lalana, R. & Beltrán, J. (1993). Una nueva especie de anfípodo hadzioido (Amphipoda: Gammaridea) del Caribe de Nicaragua. *Revista de Investigaciones Marinas*, 14, 103-109.
- Ortiz, M., Lalana, R. & Guevara, E. (1990). *Platyscelus ovooides* (Amphipoda: Hyperiididae) en el contenido estomacal del bonito (*Katsuwonus pelamis*). *Revista de Investigaciones Marinas*, 11, 174-175.
- Ortiz, M., Lalana, R. & Lio, V. (1999). Un nuevo género y una especie de anfípodo marino (Amphipoda: Aristiidae), de Cuba. *Avicennia*, 10/11, 137-142.
- Ortiz, M., Lalana, R. & Lio, V. (2001). Primera consignación del género *Protohadzia* (Amphipoda, Gammaridea), para el archipiélago cubano. *Avicennia*, 14, 129-132.
- Ortiz, M., Lalana, R. & López, M. (1992). Nueva especie de anfípodo del género *Bemlos* (Gammaridea, Corophiidae), de Cuba. *Anales del Instituto de Ciencias Marinas y Limnología, Universidad Nacional Autónoma de México*, 19, 163-166.
- Ortiz, M., Lalana, R. & Sánchez, A. (2000). Una nueva especie de anfípodo espongiícola del género *Hoplopheonoides* Shoemaker, 1956 (Gammaridea, Cyproideidae) de Cuba. *Avicennia*, 12/13, 63-68.
- Ortiz, M., Lalana, R. & Varela, C. (2002). Descripción de la hembra de *Spathiopus cojimarensis* (Amphipoda: Melitidae). *Solenodon*, 2, 17-20.
- Ortiz, M., Lalana, R. & Varela, C. (2004). Una nueva especie de anfípodo marino del género *Elasmopus* (Amphipoda, Melitidae) del Archipiélago cubano. *Avicennia*, 17, 35-40.
- Ortiz, M., Lalana, R. & Varela, C. (2006). A new species of the genus *Elasmopus* (Gammaridea, Melitidae) from Cuban waters. *Avicennia*, 17, 35-40.
- Ortiz, M., Lalana, R. & Varela, C. (2007a). Un género nuevo y una especie nueva de anfípodo espongiícola (Gammaridea: Eusiridae) de las aguas profundas del sur de Cuba. *Solenodon*, 6, 8-14.
- Ortiz, M., Lalana, R. & Varela, C. (2007b). *Tantena*, género nuevo y especie nueva de anfípodo marino (Lysianasidae) y primera consignación de la familia Ochlesidae y del género *Curidia*, con la descripción de una especie nueva para Cuba (Amphipoda, Gammaridea). *Solenodon*, 6, 20-32.
- Ortiz, M., Lalana, R. & Varela, C. (2007c). Primer registro del género *Aristias* (Amphipoda: Aristiidae) para el Mar Caribe, con la descripción de una especie nueva. *Solenodon*, 6, 33-40.
- Ortiz, M., Lalana, R. & Varela, C. (2007d). Crustáceos no Decápodos - Filo Arthropoda, sub-filo Crustacea. Lista de especies registradas en Cuba (diciembre de 2006). In R. Claro, (Ed.), *La Biodiversidad marina de Cuba*. (CD-ROM), La Habana: Instituto de Oceanología, Ministerio de Ciencia, Tecnología y Medio Ambiente.
- Ortiz, M., Lalana, R. & Varela, C. (2007e). First record of the genus *Rhachotropis* (Crustacea: Amphipoda: Gammaridea: Eusiridae) for the Cuban marine waters, with the description of a new species. *Travaux Du Museum d'Histoire Naturelle "Grigore Antipa"*, 50, 25-30.
- Ortiz, M., Lalana, R. & Varela, C. (2008). Primer registro de la familia Melphidippidae y del género *Hornellia* (Amphipoda: Gammaridea) para aguas cubanas, con la descripción de una especie nueva. *Solenodon*, 7, 7-13.
- Ortiz, M., Lalana, R., Varela, C. & Leal, S. (2003). Crustáceos marinos planctónicos (Malacostraca), de Cuba. *Revista de Investigaciones Marinas*, 24, 77-79.
- Ortiz, M. & Lemaitre, R. (1994). Crustáceos anfípodos (Gammaridea) colectados en las costas del Caribe colombiano, al sur de Cartagena. *Anales de Investigaciones Marinas de Punta Betín*, 23, 119-127.
- Ortiz, M. & Lemaitre, R. (1997). Seven new amphipods (Crustacea: Peracarida: Gammaridea) from the



- Caribbean coast of South America. *Anales del Instituto de Investigaciones Marinas de Punta Betín*, 26, 71-104.
- Ortiz, M., Lopeztegui, A., Lalana, R. & Varela, C. (2006). Registros nuevos de anfipodos planctónicos (Crustacea, Amphipoda, Hyperideae) de la costa Sur de la Isla de la Juventud, suroeste de Cuba. *Cocuyo*, 16, 15.
- Ortiz, M., Martín, A. & Atienza, D. (2000). Una nueva especie de anfipodo del género *Tiburonnella* (Crustacea, Amphipoda, Platyischnopidae) de Venezuela. *Acta Biologica Venezuelica*, 20(2), 29-36.
- Ortiz, M., Martín, A. & Díaz, Y. J. (2007). Lista y referencias de los crustáceos anfipodos (Amphipoda: Gammaridea) del Atlántico occidental tropical. *Revista de Biología Tropical*, 55(2), 479-498.
- Ortiz, M., Martín, A., Winfield, I., Díaz, Y. & Atienza, D. (2004). *Anfipodos (Crustacea: Gammaridea). Clave gráfica para la identificación de las familias, géneros y especies marinas y estuarinas del Atlántico occidental tropical*. México: Universidad Nacional Autónoma de México, Facultad de Estudios Superiores Iztacala.
- Ortiz, M. & Nazábal, J. (1984a). A new amphipod crustacean of the genus *Lembos* (Gammaridea, Aoridae) from the Cuban marine waters. *Travaux Du Museum d'Histoire Naturelle "Grigore Antipa"*, 26, 11-13.
- Ortiz, M. & Nazábal, J. (1984b). *Corocubanus*, un nuevo género de anfipodo (Amphipoda, Gammaridea, Corophiidae), de aguas Cubanas. *Revista de Investigaciones Marinas*, 5, 3-21.
- Ortiz, M. & Nazábal, J. (1988). Una nueva especie de anfipodo del género *Lembos* (Crustacea, Amphipoda), de aguas cubanas. *Revista de Investigaciones Marinas*, 9, 29-35.
- Ortiz, M. & Pérez, A. (1995). Una nueva especie de anfipodo cavernícola Hadzioideo (Amphipoda, Gammaridea) de Cuba. *Graellsia*, 51, 165-168.
- Ortiz, M. & Thomas, J. D. (2007). *Cerapus ortei* (Corophioidea: Corophiidae) un nuevo anfipodo de la costa caribeña de Costa Rica. *Avicennia*, 19, 17-24.
- Ortiz, M., Varela, C. & Lalana, R. (2011). Especie nueva de anfipodo del género *Photis* (Gammaridea: Photidae) del Archipiélago cubano. *Novitates Caribaeae*, 4, 10-16.
- Ortiz, M. & Vellido, T. (1985). Una nueva especie de anfipodo del género *Garosyrrhoë* (Synopiidae, Gammaridea) de aguas cubanas. *Revista de Investigaciones Marinas*, 6, 14-18.
- Ruffo, S. (1950). Studi sui crostacei anfipodi. XXII. Anfipodi del Venezuela raccolti dal dott. G. Marcuzzi. *Memorie del Museo Civico di Storia Naturale di Verona*, 2, 49-65.
- Ruffo, S. (1954). Studi sui crostacei anfipodi. XL. Nuovi anfipodi raccolti nel Venezuelano dal Prof. G. Marcuzzi. *Memorie del Museo Civico di Storia Naturale di Verona*, 4, 117-125.
- Ruffo, S., Krapp-Schickel, T. & Gable, M. F. (2000). The genus *Maera* (Crustacea: Amphipoda: Melitidae) from Bermuda. *Postilla*, 221, 1-35.
- Sánchez, L. (1985). *Algunos aspectos biológicos de Talorchestia margaritae Stephensen, 1948 (Amphipoda: Talitridae), de la Isla de Margarita, Venezuela*. (Tesis de licenciatura). Universidad de Oriente, Cumaná, Venezuela.
- Shih, C-t. (1969). The systematic and biology of the family Phronimidae (Crustacea: Amphipoda). *Dana Reports*, 74, 1-100.
- Shoemaker, C. R. (1926). Amphipods of the family Bateidae in the collection of the United States Museum. *Proceeding of the United States of Natural Museum*, 68(2626), 1-26.
- Shoemaker, C. R. (1933a). Two new genera and six new species of Amphipoda from Tortugas. Papers of the Tortugas Laboratory. *Carnegie Institution of Washington*, 28(435), 245-256.
- Shoemaker, C. R. (1933b). Amphipoda from Florida and the West Indies. *American Museum Novitates*, 598, 1-24.
- Shoemaker, C. R. (1934). Reports on the collections obtained by the first Johnson-Smithsonian deep-sea expedition to the Puerto Rican deep. Three new amphipods. *Smithsonian Miscellaneous Collections*, 9(12), 1-6.
- Shoemaker, C. R. (1935). The Amphipods of Porto Rico and the Virgin Islands. *New York Academy of Sciences Scientific Survey of Porto Rico and the Virgin Islands*, 15(2), 229-253.
- Shoemaker, C. R. (1945). The Amphipoda of the Bermuda Oceanographic Expeditions 1929-1932. *Zoologica*, 30, 185-266.
- Shoemaker, C. R. (1948). The Amphipoda of the Smithsonian-Roebbling Expedition to Cuba in 1937. *Smithsonian Miscellaneous Collections*, 110(3), 1-15.
- Spalding, M. D. (2004). *A guide to the coral reefs of the Caribbean*. Berkeley: University of California Press.
- Stebbing, T. R. R. (1888). Report on the Amphipoda collected by H.M.S. Challenger during the years 1873-76. *Zoology*, 29, 1-1737.
- Stebbing, T. R. R. (1897). Amphipoda from the Copenhagen Museum and other sources. *Transactions of Linnean Society of London, Serie Zoology*, 7, 25-45.
- Stebbing, T. R. R. (1903). Amphipoda from Costa Rica. *Proceedings of the United States Natural Museum*, 26, 925-931.
- Stebbing, T. R. R. (1906). Amphipoda I. Gammaridea. *Das Tierreich*, 21, 806.

- Stebbing, T. R. R. (1908). A new amphipod crustacean, *Orchestoidea biolleyi*, from Costa Rica. *Proceedings of the United States Natural Museum*, 34, 241-244.
- Stephensen, K. (1933). Amphipoda from the marine salines of Bonaire and Curacao. *Zoologische Ergebnisse einer Reise nach Bonaire, Curacao und Aruba im Jahre 1930. Zoologische Jahrbucher, Systematik*, 64, 437-446.
- Stephensen, K. (1948). Amphipods from Curacao, Bonaire, Aruba and Margarita. *Studies on Fauna Curaçao, Aruba, Bonaire and the Venezuelan Islands*, 11(3), 1-20.
- Stock, J. H. (1976). A new member of the crustacean suborder Ingolfiellidea from Bonaire. *Studies on Fauna Curaçao*, 50, 56-75.
- Stock, J. H. (1977a). The taxonomy and zoogeography of the Hadziid amphipoda, with emphasis on the West Indian taxa. *Studies on Fauna Curaçao*, 55, 1-130.
- Stock, J. H. (1977b). The zoogeography of the crustacean suborder Ingolfiellidea with descriptions of new West Indian taxa. *Studies on Fauna Curaçao*, 55, 131-146.
- Stock, J. H. (1978). *Bogidiella martini*, a new hypogean amphipod from the island of Saint-Martin (West Indies) and the zoogeography of the Bogidiellidae. *International Journal of Speleology*, 9(2), 103-113.
- Stock, J. H. (1979). New data on taxonomy and zoogeography of Ingolfiellid crustacea. *Bijdragen tot de Dierkunde*, 49(1), 81-96.
- Stock, J. H. (1980). A new cave amphipod (Crustacea) from Curacao: *Psammogammarus caesicolus* n. sp. *Bijdragen tot de Dierkunde*, 50, 375-386.
- Stock, J. H. (1981). The taxonomy and zoogeography of the family Bogidiellidae (Crustacea, Amphipoda), with emphasis on the west indian taxa. *Bijdragen tot de Dierkunde*, 51, 345-374.
- Stock, J. H. (1983). The stygobiont amphipoda of Jamaica. *Bijdragen tot de Dierkunde*, 53, 267-286.
- Stock, J. H. (1985). Bogidiellidae (Amphipoda) from Haiti and some general rules on the occurrence of Crustacea Malacostraca in inland groundwaters of the West Indies. *Stygologia*, 1, 208-223.
- Stoner, A. W. & Lewis, F. G. (1985). The influence of quantitative and qualitative aspects of habitat complexity in tropical sea-grass meadows. *Journal of Experimental Marine Biology and Ecology*, 94(1-3), 19-40.
- Sturaro, N. & Guerra-García, J. M. (2012). A new species of *Caprella* (Crustacea: Amphipoda) from the Mediterranean Sea. *Helgolaender Marine Research*, 66(1), 33-42. doi: 10.1007/s10152-011-0244-5.
- Takeuchi, I. (1999). Checklist and bibliography of the Caprellidea (Crustacea: Amphipoda) from Japanese waters. *Otsuchi Marine Science*, 24, 5-17.
- Thiel, M., Guerra-García, J. M., Lancellotti, D. A. & Vásquez, N. (2003). The distribution of littoral caprellids (Crustacea: Amphipoda: Caprellidea) along the Pacific coast of continental Chile. *Revista Chilena de Historia Natural*, 76(2), 203-218.
- Thomas, J. D. (1983). *Curidia debrogania*, a new genus and species of amphipod (Crustacea: Ochlesidae) from the Barrier Reef of Belize, Central America. *Proceedings of Biological Society of Washington*, 96(1), 127-133.
- Thomas, J. D. (1993). Biological monitoring and tropical biodiversity in marine environments: a critique with recommendations, and comments on the use of amphipods as bioindicators. *Journal of Natural History*, 27(4), 795-806.
- Thomas, J. D. (1997). Systematics, ecology and phylogeny of the Anamixidae (Crustacea: Amphipoda). *Records of Australian Museum*, 49, 35-98.
- Thomas, J. D. & Barnard, J. L. (1983). The Platyschnopidae of America (Crustacea: Amphipoda). *Smithsonian Contributions to Zoology*, 375, 1-33.
- Thomas, J. D. & Barnard, J. L. (1985). *Periculodes cerasinus*, n.sp., the first record of the genus from the Caribbean Sea (Amphipoda: Oedicerotidae). *Proceedings of Biological Society of Washington*, 98(1), 98-106.
- Thomas, J. D. & Barnard, J. L. (1986). New genera and species of the *Megaluropus* group (Amphipoda, Megaluropidae) from American Seas. *Bulletin of Marine Science*, 38(3), 442-476.
- Thomas, J. D. & Barnard, J. L. (1989). *Gammaropsis arawakia*, a new species of marine amphipoda (Crustacea) from Jamaica. *Proceedings of Biological Society of Washington*, 102(1), 89-94.
- Thomas, J. D. & Barnard, J. L. (1990). *Gitana dominica*, a new species from the Caribbean Sea (Amphipoda: Amphiloichidae). *Proceedings of Biological Society of Washington*, 103(3), 617-623.
- Thomas, J. D. & Barnard, J. L. (1991a). *Photis trapherus*, a new Elephantine species from the Caribbean Sea (Crustacea: Amphipoda). *Proceedings of Biological Society of Washington*, 104(1), 96-100.
- Thomas, J. D. & Barnard, J. L. (1991b). Two new species of *Netamelita* from the Caribbean Sea (Crustacea: Amphipoda). *Proceedings of Biological Society of Washington*, 104(3), 583-592.
- Thomas, J. D. & Klebba, K. N. (2006). Studies of commensal leucothoid amphipods: two new sponge-inhabiting species from south Florida and the western Caribbean. *Journal of Crustacean Biology*, 26(1), 13-22.
- Thomas, J. D. & Klebba, K. N. (2007). New species and host associations of commensal leucothoid

- amphipods from coral reefs in Florida and Belize (Crustacea: Amphipoda). *Zootaxa*, 1494, 1-44.
- Vader, W. (2005). *How many Amphipod Species?*. Poster presented at XII International Amphipod Colloquium. Cork, Ireland.
- Vainola, R., Witt, J. D. S., Grabowski, M., Bradbury, J. H., Jazdzewski, K. & Sket, B. (2008). Global diversity of amphipods (Amphipoda: Crustacea) in freshwater. *Hydrobiologia*, 595, 241-255. doi: 10.1007/s10750-007-9020-6.
- Van der Ham, J. L. & Vonk, R. (2003). A phylogenetic analysis of the *Eriopisa* complex (Crustacea: Amphipoda: Melitidae) and a new species from beach interstitial in Venezuela. *Journal of Natural History*, 37(7), 779-796.
- Varela, C., Ortiz, M. & Lalana, R. (2003). Crustáceos (Percarida y Decapoda), de la costa sur de la Península de Guanahabibes, Cuba. *Revista de Investigaciones Marinas*, 24, 73-76.
- Villarroel, E. J. & Graziani, C. A. (1997). Primer registro de *Caprella danilevskii* Czerniavskii (Amphipoda: Caprellidae) en Venezuela. *Saber, Universidad de Oriente*, 9, 102-103.
- Vinogradov, G. (1999). Amphipoda. In D. Boltovskoy (Ed.), *Zooplankton of the Southwestern Atlantic*. (1141-1240). Backhuys, Leiden.
- Vonk, R. (1988). *Psammomelita uncinata* n.g., n.sp. (Crustacea, Amphipoda, Melitidae) from infralittoral sand interstices on Curaçao. *Stygologia*, 4, 166-176.
- Vonk, R. (1989). *Nuuanu curvata* n.sp and *Melita leio-telson* n.sp. (Crustacea, Amphipoda) from beach interstitial on Curaçao. *Foundation for Scientific Research in Surinam and the Netherlands Antilles, Amsterdam*, 123, 185-198.
- Vonk, R. (1990). *Thalassostygius exiguus* n.g., n. sp., a new marine interstitial melitid (Crustacea, Amphipoda) from Curaçao and Klein Bonaire (Netherlands Antilles). *Stygologia*, 5, 43-48.
- Vonk, R. (1991). Two marine interstitial *Metaniphargus* species (Crustacea, Amphipoda) from Hawaii and the Cayman Islands. *Stygologia*, 6, 111-118.
- Vonk, R. & Schram, F. R. (2003). Ingolfiellidea (Crustacea, Malacostraca, Amphipoda): a phylogenetic and biogeographic analysis. *Contributions to Zoology*, 72(1), 39-72.
- Vonk, R. & Stock, J. H. (1987). *Psammogammarus longidactylus* n.sp., a new cave amphipod (Crustacea) and other stygobiont amphipods from Bonaire. *Stygologia*, 3, 241-251.
- Wakabara, Y., Tararam, A. S., Valério-Berardo, M. T., Duleba, W. & Pereira-Leite, F. P. (1991). Gammaridean and caprellidean fauna from Brazil. *Hydrobiologia*, 223, 69-77.
- WoRMS. (2012). Amphipoda. In J. Lowry (Ed.), *World Amphipoda database*. Recuperado de <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1135>
- Zeidler, W. & De Broyer, C. (2009). Volume 3: Catalogue of the Hyperiid Amphipoda (Crustacea) of the Southern Ocean with distribution and ecological data. *Bulletin de L'Institut Royal des Sciences Naturelles de Belgique*, 79(1), 1-96.



APPENDIX 1

Taxonomic classification of the amphipod fauna from Caribbean countries or subregion: PRC, Puerto Rico, HIS, Hispaniola, JAM, Jamaica, CUB, Cuba, CAY, Cayman Islands, MEX, Mexico, BEL, Belize, GUA, Guatemala, HON, Honduras, NIC, Nicaragua, CRC, Costa Rica, PAN, Panama, COL, Colombia, VEN, Venezuela, ABC, Netherlands Antilles, TYT, Trinidad & Tobago, LAN, Lesser Antilles

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY
Suborder Caprellidea Leach, 1814	<i>Aeginella spinosa</i> Boeck, 1861	X																
Family Caprellidae Leach, 1814	<i>Caprella andreae</i> Mayer, 1890	X	X															
	<i>Caprella damilevskii</i> Czerniavskii, 1868		X										X	X				
	<i>Caprella equilibra</i> Say, 1818		X										X	X				
	<i>Caprella penantis</i> Leach, 1814		X										X	X				
	<i>Caprella scaura</i> Temptelom, 1836		X													X		
	<i>Metaprotella hummelincki</i> McCain, 1968	X																
	<i>Paracaprella digitimanus</i> Quiete, 1971		X										X	X				
	<i>Paracaprella pusilla</i> Mayer, 1890	X	X										X	X				
	<i>Paracaprella tenuis</i> Mayer, 1903		X															
Family Caprellinoidea Laubitz, 1993	<i>Pseudaeiginella antigueae</i> Barnard, 1932																	
	<i>Pseudaeiginella biscaynensis</i> (McCain, 1968)	X	X															
	<i>Pseudaeiginella columbiana</i> Guerra-Garcia, Krapp-Schickel & Müller, 2006	X											X					
Family Parriambiidae Laubitz, 1993	<i>Cubadeutella cavernicola</i> Ortiz, Guerra-Garcia & Lalana, 2009	X																
	<i>Deutella californica</i> Mayer, 1890		X															
	<i>Deutella caribensis</i> Guerra-Garcia, Krapp-Schickel & Müller, 2006	X	X										X	X				
	<i>Deutella incerta</i> (Mayer, 1903)		X										X	X				
	<i>Deutella margaritae</i> Guerra-Garcia, 2003												X	X				
	<i>Deutella mayeri</i> Stebbing, 1895	X											X	X				
Family Phitsidae Vassilenko, 1968	<i>Hemiaegina minuta</i> Mayer, 1890	X	X										X	X				
	<i>Hemiprotos wigleyi</i> McCain, 1968	X	X										X	X				
	<i>Phitsica antillensis</i> Mayer, 1903																	
	<i>Phitsica marina</i> Slabber, 1769																	
Family Protellidae McCain, 1970	<i>Mayerella redunca</i> McCain, 1968	X	X											X				
	<i>Triella chibcha</i> Guerra-Garcia, Krapp-Schickel & Müller, 2006												X					

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY
Suborder Gammaridea Latreille, 1802																		
Family Ampeliscidae Costa, 1857	<i>Ampelisca abdita</i> Mills, 1964		X															
	<i>Ampelisca abyssicola</i> Stebbing, 1888			X														
	<i>Ampelisca agassizi</i> (Hudd, 1896)		X						X			X	X					
	<i>Ampelisca bicarinata</i> Croeke & Heard, 1983		X										X	X				
	<i>Ampelisca brevisimulata</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca burketi</i> J. L. Barnard & Thomas, 1989		X										X	X				
	<i>Ampelisca cristata cristata</i> Holmes, 1908		X										X	X				
	<i>Ampelisca cristata microdentata</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca cristoides</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca holmesi</i> Pearse, 1908		X										X	X				
	<i>Ampelisca lobata</i> Holmes, 1908		X										X	X				
	<i>Ampelisca mexicana</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca pacifica</i> Holmes, 1908		X										X	X				
	<i>Ampelisca parapacifica</i> Goetze & Heard, 1984		X										X	X				
	<i>Ampelisca parapanamensis</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca parva</i> Barnard & Agard, 1986		X										X	X				
	<i>Ampelisca pugetica</i> Stimpson, 1864		X										X	X				
	<i>Ampelisca romigi</i> J.L. Barnard, 1954		X										X	X				
	<i>Ampelisca schellenbergi</i> Shoemaker, 1933		X										X	X				
Family Ampeliscidae Costa, 1857	<i>Ampelisca spinipes</i> Boeck, 1861		X										X	X				
	<i>Ampelisca vadorum</i> Mills, 1963		X										X	X				
	<i>Ampelisca venetiensis</i> Shoemaker, 1916		X										X	X				
	<i>Ampelisca verrilli</i> Mills, 1967		X										X	X				
	<i>Byblis cubensis</i> (Ortiz & Gómez, 1979)		X										X	X				
	<i>Byblisoides blasensis</i> J.L. Barnard, 1964		X										X	X				
Family Amphilochoidae Boeck, 1871	<i>Amphilochous acaciicola</i> Martin, Ortiz & Aienza, 2001		X										X	X				
	<i>Apolochus casahoya</i> (McKinney, 1978)		X										X	X				
	<i>Apolochus delacaya</i> (McKinney, 1978)		X										X	X				
	<i>Apolochus neopolitanus</i> (Della Valle, 1893)		X										X	X				
	<i>Apolochus pillaii</i> (J. L. Barnard & Thomas, 1983)		X										X	X				

APPENDIX 1 (Continued)

Suborder / Family	Species	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY	
Family Amphilocheidae Boeck, 1871	<i>Gitana dominica</i> Thomas & Barnard, 1990	X											X					
	<i>Gitanopsis rempladoi</i> Ortiz & Lalana, 1995	X											X					
	<i>Houstonius laguna</i> (McKinney, 1978)	X	X									X	X					
	<i>Houstonius torugae</i> (Shoemaker, 1933)	X	X									X	X					
	<i>Ampithoe hirsutimanus</i> Ortiz & Lemaitre, 1997											X	X					
	<i>Ampithoe longimana</i> Smith, 1873											X	X					
	<i>Ampithoe marcuizi</i> Ruffo, 1954	X	X	X						X		X	X					
	<i>Ampithoe pollex</i> Kunkel, 1910	X				X							X					
	<i>Ampithoe ramondi</i> Audouin, 1826	X	X										X					
	<i>Ampithoe valida</i> Smith, 1873	X	X											X				
Family Aoridae Walker, 1908	<i>Cynadusa compta</i> (Smith, 1873)	X	X										X					
	<i>Cynadusa filosa</i> Savigny, 1816	X	X										X					
	<i>Pseudampithoides bacescui</i> Ortiz, 1976	X	X									X	X					
	<i>Pseudampithoides bacescui</i> Ortiz, 1976	X	X									X	X					
	<i>Pseudampithoides incurvata</i> (Just, 1977)									X								
	<i>Sumampithoe pelagica</i> (Milne-Edwards, 1830)	X	X										X					
	<i>Ampideutopus dolichocephalus</i> Myers, 1968												X					
	<i>Bemios barnarati</i> (Ortiz & Nazábal, 1988)	X												X				
	<i>Bemios brunneomaculatus</i> Myers, 1977	X												X				
	<i>Bemios dentischiium</i> (A. A. Myers, 1977)	X												X				
<i>Bemios foresti</i> (Mateus & Mateus, 1966)	X											X	X					
<i>Bemios intermedius</i> (Schellenberg, 1938)	X											X	X					
<i>Bemios kunkelae</i> (A. A. Myers, 1977)	X	X											X					
<i>Bemios longicornis</i> (A. A. Myers, 1978)	X	X	X									X	X					
<i>Bemios rolani</i> Ortiz & Lalana, 2002	X												X					
<i>Bemios sammartini</i> Ortiz, Lalana & López, 1992	X												X					
<i>Bemios scolosternum</i> Ortiz & Lemaitre, 1997	X												X					
<i>Bemios spinicarpus inermis</i> (A. A. Myers, 1979)	X	X											X					
<i>Bemios spinicarpus spinicarpus</i> (Pearse, 1912)	X	X	X										X					
<i>Bemios unicomis</i> (Brynum & Fox, 1977)	X	X											X					
<i>Globosolembos francanni</i> (Reid, 1951)	X												X					
<i>Globosolembos smithi</i> (Holmes, 1905)	X	X											X					

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY
Family Aoridae Walker, 1908	<i>Granditerella bonnieroides</i> Stephensen, 1947	X	X										X	X	X			
	<i>Lembos mayensis</i> Ortiz & Nazábal, 1984	X																
	<i>Lembos unifasciatus reductus</i> Myers, 1979			X									X					
	<i>Lembos unifasciatus unifasciatus</i> Myers, 1977		X	X										X				
	<i>Puramicrodeuopius myersi</i> (Brynum & Fox, 1977)												X	X				
	<i>Pedicrophium laminosum</i> (Pearse, 1912)		X											X				
	<i>Plestolombos ovalipes</i> (A. A. Myers, 1979)													X				
	<i>Plestolombos rectangularis</i> (A. A. Myers, 1977)		X											X				
	<i>Rudilemboides heardi</i> Ortiz & Lalana, 1996		X											X				
	<i>Rudilemboides naglei</i> Bousfield, 1973			X										X				
	<i>Argissa hamatipes</i> (Norman, 1869)			X														
	<i>Aristias bicornuta</i> Ortiz, Lalana & Varela, 2007		X															
	<i>Aristias captiva</i> Lowry & Stoddart, 1997																	
	<i>Boca campi</i> Lowry & Stoddart, 1997																	
<i>Miramarassa sanchezi</i> Ortiz, Lalana & Lio, 1999		X					X											
<i>Batea aldebaranae</i> Coleman, 2011																		
<i>Batea bousfieldi</i> (Ortiz, 1991)		X																
<i>Batea campi</i> (Ortiz, 1991)		X																
<i>Batea carinata</i> (Shoemaker, 1926)		X	X				X						X	X				
<i>Batea catharinensis</i> Müller, 1865		X	X										X	X				
<i>Batea cuspidata</i> (Shoemaker, 1926)		X	X										X	X				
<i>Batea schoffi</i> Ortiz & Lemaitre, 1997		X	X				X						X	X				
<i>Batea transversa</i> Shoemaker, 1926		X	X										X	X				
<i>Biancolina lowryi</i> Ortiz & Lalana, 1996		X																
<i>Actogidiella cultrifera</i> Stock, 1981																X		
<i>Amillogidiella martini</i> (Stock, 1978)																X		
<i>Bogidiella hamatula</i> Stock, 1985																		
<i>Bogidiella perla</i> Stock, 1981																		
<i>Bogidiella prionura</i> Stock, 1985																		
<i>Bogidiella virginialis</i> Stock, 1981																		
<i>Martigidiella crassipes</i> Stock, 1981																		

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Callipteridae Sars, 1893	<i>Stenopleura atlantica</i> Stebbing, 1888	X																
Family Cebocariidae Lowry & Stoddart, 2011	<i>Cebocaris grutesca</i> J.L. Barnard, 1964			X														
Family Cheluridae Allman, 1847	<i>Tropichelura gomezi</i> Ortiz, 1976	X	X	X		X								X			X	
Family Colomastigidae Stebbing, 1899	<i>Tropichelura insilae</i> (Calman, 1910)			X							X			X				
	<i>Colomastix bousfieldi</i> LeCroy, 1995	X	X											X				
	<i>Colomastix camura</i> LeCroy, 1995		X									X		X				
	<i>Colomastix falcirama</i> LeCroy, 1995		X															
	<i>Colomastix gibbosa</i> LeCroy, 1995		X															
	<i>Colomastix halichondriae</i> Bousfield, 1973	X	X								X			X				
	<i>Colomastix heardi</i> LeCroy, 1995	X												X				
	<i>Colomastix irciniae</i> LeCroy, 1995		X															
	<i>Colomastix janiceae</i> Heard & Perlmutter, 1977	X	X											X				
	<i>Colomastix tridentata</i> LeCroy, 1995	X	X															
	<i>Americorophium rioplatense</i> (Giambiagi, 1926)																	
	<i>Apocorophium acutum</i> (Chevreux, 1908)	X	X															
	<i>Apocorophium lacustre</i> (Vanhöffen, 1911)																	
<i>Apocorophium louisianum</i> (Shoemaker, 1934)			X															
<i>Apocorophium simile</i> (Shoemaker, 1934)		X																
<i>Caribboeetes barbadensis</i> Just, 1983																		
<i>Caribboeetes crassicornis</i> Just, 1984																		
<i>Caribboeetes intermedius</i> Just, 1984																		
<i>Caribboeetes justi</i> Ortiz & Lemaitre, 1997																		
<i>Caribboeetes pteryornis</i> Just, 1984																		
<i>Crassicorophium bonellii</i> (Milne Edwards, 1830)		X																
<i>Laticorophium baconi</i> (Shoemaker, 1934)			X															
<i>Leptochelirus rhizophorae</i> Ortiz & Lalana, 1980		X	X											X				
<i>Monocorophium acherusicum</i> (Costa, 1853)		X	X											X				
<i>Monocorophium insidiosum</i> (Crawford, 1937)		X	X										X	X				
<i>Cyphocaris anonyx</i> Boeck, 1871		X																
<i>Cyphocaris johnsoni</i> Shoemaker, 1934																		

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY
Family Cyproideidae Barnard, 1974	<i>Hoplopleoneoides obesa</i> Shoemaker, 1956		X			X					X							
	<i>Hoplopleoneoides shoemakeri</i> Ortiz, Lalana & Sanchez-Diaz, 2000	X																
Family Dexaminidae Leach, 1814	<i>Aberraplus aberrantis</i> (J.L. Barnard, 1962)		X	X														
	<i>Atylus urocarimatus</i> McKimney, 1980																	
	<i>Lepechinella raua</i> J.L. Barnard, 1973											X	X	X				
Family Endeavouridae Lowry & Stoddart, 1997	<i>Nototropis minikoi</i> (A.O. Walker, 1905)	X	X				X											
Family Eurytheneidae Stoddart & Lowry, 2004	<i>Ensayara jumane</i> Barnard & Thomas, 1990	X	X															
Family Eusiridae Stebbing, 1888	<i>Eusirogenes adad</i> J.L. Barnard, 1964											X						
	<i>Eusiroides monoculoides</i> (Haswell, 1880)													X				
	<i>Eusiroides yucatanensis</i> McKimney et al., 1980													X				
	<i>Nasagenia bacescui</i> Ortiz & Lalana, 1994	X											X					
	<i>Nasagenia comisariensis</i> Ortiz & Lemaitre, 1997												X	X				
	<i>Nasagenia yucatanensis</i> Ledoyer, 1986												X	X				
	<i>Pleustiroides alcoladoi</i> Ortiz, Lalana & Varela, 2007	X	X															
	<i>Pontogenia bartschi</i> Shoemaker, 1948	X																
	<i>Rhachotropis lobata</i> Shoemaker, 1934																	
	<i>Rhachotropis portoricana</i> J.L. Barnard, 1964																	
	<i>Rhachotropis wimaderi</i> Ortiz, Lalana & Varela, 2007	X																
	<i>Sennaia bidactyla</i> Bellan-Santini, 1997																	
	<i>Telhygenia longleyi</i> (Shoemaker, 1933)	X																
Family Gammaridae Latreille, 1802	<i>Gammarus mucronatus</i> Say, 1818		X															
	<i>Gammarus tigrinus</i> Sexton, 1939																	
Family Hadziidae Karaman, 1943	<i>Allovecchia gurneei</i> Holsinger & Peck, 1968			X														
	<i>Apoveckelia serrata</i> Stock, 1985																	
	<i>Bahadzia bazonici</i> Holsinger, 1992																	
	<i>Bahadzia jaraguensis</i> Jaume & Wagner, 1998		X															
	<i>Bahadzia latipalpus</i> Stock, 1985																	
	<i>Bahadzia setodactylus</i> Holsinger, 1992		X															
	<i>Bahadzia yagerrae</i> Ortiz & Perez, 1995	X																

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Hadziidae Karaman, 1943	<i>Crangoveckelia mixta</i> Stock, 1985				X													
	<i>Crangoveckelia spincicauda</i> Stock, 1985				X													
	<i>Mayaveckelia cenoticoala</i> Holsinger, 1977		X															
	<i>Mayaveckelia yucatanensis</i> Holsinger, 1977		X															
	<i>Metaniphargus anchihalinus</i> Stock, 1983					X												
	<i>Metaniphargus bousfieldi</i> Stock, 1977			X														
	<i>Metaniphargus chaetodactylus</i> Stock, 1985				X													
	<i>Metaniphargus crenatus</i> Stock, 1985				X													
	<i>Metaniphargus curasavicus curasavicus</i> Stephensen, 1933														X			
	<i>Metaniphargus curasavicus orientalis</i> Stock, 1977														X			
	<i>Metaniphargus haitianus</i> Stock, 1985				X													
	<i>Metaniphargus hyporheicus</i> Stock, 1983					X												
	<i>Metaniphargus jamaicae</i> (Holsinger, 1974)						X											
	<i>Metaniphargus juberthiei</i> Stock & Vermeulen, 1985						X									X		
	<i>Metaniphargus longipalpus</i> Stock, 1985				X													
	<i>Metaniphargus palpator</i> Stock, 1977					X												
	<i>Metaniphargus pedunculatus</i> Stock, 1985				X													
	<i>Metaniphargus plamicauda</i> Stock, 1985				X													
	<i>Metaniphargus sablonis</i> Vonk, 1991					X												
<i>Pintaveckelia grandis</i> Stock, 1985					X													
<i>Protohadzia schoenerae</i> (Fox, 1973)		X	X	X														
<i>Saliveckelia emarginata</i> Stock, 1977														X				
<i>Saliveckelia holsingeri</i> Stock, 1977														X				
<i>Tulaveckelia cernua</i> Holsinger, 1990			X															
<i>Haustorius mexicanus</i> Ortiz, Cházaro-Olvera & Winfield, 2001														X				
<i>Parhyalella nisibetae</i> Lazo-Wasem & Gable, 2001		X																
<i>Parhyalella whepleyi</i> (Shoemaker, 1933)		X	X										X				X	
<i>Apothyale media</i> (Dana, 1853)		X	X										X					
<i>Apothyale perieri</i> (Lucas, 1849)		X	X											X				
<i>Hyale pygmaea</i> Ruffo, 1950		X	X											X				
<i>Parargissa galathea americana</i> J.L. Barnard, 1961												X						



APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Hyalidae Bulycheva, 1957	<i>Parhyale fascigera</i> Stebbing, 1897	X	X	X									X	X	X	X		
	<i>Parhyale hawaiiensis</i> (Dana, 1853)		X	X									X	X	X	X		
	<i>Parhyale inyacka</i> K.H. Barnard, 1916	X														X		
	<i>Protohyale diploclacyla</i> (Stebbing, 1899)													X				
	<i>Protohyale macrodactyla</i> Stebbing, 1899	X	X											X	X	X		
	<i>Audulla chelifera</i> Chevreux, 1901	X	X										X					
	<i>Chevalia aviculae</i> Walker, 1904	X	X															
	<i>Chevalia carpenteri</i> Barnard & Thomas, 1987	X	X				X											
	<i>Chevalia mexicana</i> Pearse, 1912	X	X				X										X	
	<i>Gammaropsis aravakia</i> Thomas & Barnard, 1989						X											
	<i>Latigammaropsis atlantica</i> (Stebbing, 1888)	X	X											X	X	X		
	<i>Latigammaropsis togoensis</i> (Schellenberg, 1925)		X											X	X	X		
	<i>Microprotopus raneji</i> Wigley, 1966													X	X	X		
	<i>Microprotopus shoemakeri</i> Lowry, 1972													X	X	X		
<i>Photis dentata</i> Shoemaker, 1945																		
<i>Photis lecrolyae</i> Ortiz, Varela & Lalana, 2011	X																	
<i>Photis longicaudata</i> (Bate & Westwood, 1862)		X																
<i>Photis macromana</i> McKinney, Kalke & Holland, 1978			X															
<i>Photis melanica</i> McKinney, 1980			X															
<i>Photis pugnator</i> Shoemaker, 1945			X															
<i>Photis sarue</i> Souza-Filho & Serejo, 2010																		
<i>Photis trapherus</i> Thomas & J. L. Barnard, 1991																		
<i>Cerapus benthophilus</i> Thomas & Heard, 1979		X	X		X									X	X	X		
<i>Cerapus cidjoe</i> Lowry & Thomas, 1991			X											X	X			
<i>Cerapus ortzei</i> Ortiz & Thomas, 2007																		
<i>Cerapus thomasi</i> Ortiz & Lemaitre, 1997																		
<i>Corocubanus guitari</i> Ortiz & Nazabal, 1984		X											X					
<i>Erichthonius brasiliensis</i> (Dana, 1853)		X	X										X	X	X	X		
<i>Erichthonius rubricornis</i> (Stimpson, 1853)		X	X										X	X	X	X		
<i>Microjassa tetradonta</i> Conlan, 1995																		
<i>Neischyrocerus vidali</i> Ortiz & Lalana, 2002		X																

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY
Family Leucothoidae Dana, 1852	<i>Anamixis cavatara</i> Thomas, 1997	X	X								X			X				
	<i>Anamixis hanseni</i> Stebbing, 1897	X					X						X					
	<i>Anamixis vanga</i> Thomas, 1997		X								X			X				
	<i>Leucothoe ashleyae</i> Thomas & Klebba, 2006		X					X										
	<i>Leucothoe aytonia</i> Bellan-Santini, 1997		X															
	<i>Leucothoe barana</i> Thomas & Klebba, 2007						X					X						
	<i>Leucothoe flammosa</i> Thomas & Klebba, 2007						X					X						
	<i>Leucothoe garfuna</i> Thomas & Klebba, 2007						X					X						
	<i>Leucothoe kensleyi</i> Thomas & Klebba, 2006						X											
	<i>Leucothoe laurensi</i> Thomas & Ortiz, 1995		X									X			X			
	<i>Leucothoe panpulco</i> J.L. Barnard, 1961		X									X						
	<i>Leucothoe saron</i> Thomas & Klebba, 2007							X										
	<i>Leucothoe spinicarpa</i> (Abildgaard, 1789)							X										
<i>Leucothoe ubouhi</i> Thomas & Klebba, 2007							X											
<i>Leucothoe wurii</i> Thomas & Klebba, 2007							X											
<i>Neponamixis dianthus</i> Thomas, 1997							X											
<i>Liljeborgia bousfieldi</i> McKinney, 1979		X	X															
<i>Listriella andresi</i> Martin, Ortiz & Atienza, 2000														X				
<i>Listriella barnardi</i> Wigley, 1966							X											
<i>Listriella carinata</i> McKinney, 1979			X											X				
<i>Listriella difficilis</i> J.L. Barnard, 1959		X																
<i>Listriella kensleyi</i> Ortiz & Lalana, 1996		X																
<i>Listriella quiniana</i> McKinney, 1979			X															
<i>Bonassa bonairensis</i> (Stephensen, 1933)															X			
<i>Concarnes concarus</i> (Shoemaker, 1933)			X				X				X	X						
<i>Eclecticus eclecticus</i> Lowry & Stoddart, 1997												X						
<i>Lepiduristes lepidus</i> (J.L. Barnard, 1964)																		
<i>Lysianassa hummelincki</i> Stephensen, 1933																		
<i>Lysianopsis alba</i> Holmes, 1903		X	X											X				
<i>Orchomene madagascarensis</i> (Shoemaker, 1942)		X																
<i>Orchomene stocki</i> Bellan-Santini, 1990																		
<i>Orenoqueia serrata</i> Bellan-Santini, 1997																		

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY	
Family Lysianassidae Dana, 1849	<i>Otenvaldiera kymbalion</i> Jaume & Wagner, 1998			X															
	<i>Paracentromedon carabicus</i> J.L. Barnard, 1964										X								
	<i>Shoemakerella cubensis</i> (Stebbing, 1897)	X	X										X						
	<i>Socarnopsis catacumba</i> (Clark & Barnard, 1985)	X																	
	<i>Socarnopsis karamani</i> Ortiz, Garcia-Debras & Lalana, 2003	X																	
	<i>Tryphosella apalachicola</i> Lowry & Stoddart, 1997														X				
	<i>Tryphosella uristesi</i> Bellan-Santini, 1997																		
	<i>Gibberosus myersi</i> (McKinney, 1980)		X																
	<i>Resupinus spinicaudatus</i> Thomas & Barnard, 1986				X														
	<i>Anamaera hixonii</i> Thomas & J. L. Barnard, 1985																		
Family Megaluroptidae Thomas & Barnard, 1986	<i>Anelasmopus kraui</i> Oliveira, 1953																		
	<i>Ceradocus (Denticeradocus) rubromaculatus</i> (Stimpson, 1855)																		
	<i>Ceradocus (Denticeradocus) shearafi</i> Shoemaker, 1948	X	X		X						X		X	X					
	<i>Ceradocus shoemakeri</i> Fox, 1973	X	X										X	X					
	<i>Confodiopisa caesicola</i> (Stock, 1980)																		
	<i>Dulichitella appendiculata</i> (Say, 1818)	X																	
	<i>Dumosus atari</i> Thomas & J. L. Barnard, 1985		X								X								
	<i>Elasmopus balkomanus</i> Thomas & J. L. Barnard, 1988		X																
	<i>Elasmopus bampo</i> J.L. Barnard, 1979																		
	<i>Elasmopus brasiliensis</i> (Dana, 1853)																		
	<i>Elasmopus eileri</i> Ortiz, Lalana & Varela, 2004																		
	<i>Elasmopus lenaitrei</i> Ortiz & Lalana, 1994	X																	
	<i>Elasmopus levis</i> (S. I. Smith, 1873)		X																
	<i>Elasmopus pectenarius</i> (Bate, 1862)	X	X												X				
	<i>Elasmopus pocillinus</i> (Bate, 1862)	X	X												X				
<i>Elasmopus rapax</i> Costa, 1853	X	X												X					
<i>Elasmopus spinidactylus</i> Chevreaux, 1907	X																		
<i>Elasmopus thomasi</i> Ortiz & Lalana, 1994	X												X						
<i>Eriopisa incisa</i> McKinney, Kalke & Holland, 1978													X						
<i>Eriopisa mochiniae</i> Van der Ham & Yonk, 2003																			
<i>Hamimuera hamigera</i> (Haswell, 1879)	X																		

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY	
Family Melitidae Bousfield, 1973	<i>Jerbaria americana</i> Wätling, 1981		X																
	<i>Maera jerrica</i> Krapp-Schickel & Jarrett, 2000		X											X					
	<i>Maeracoota galani</i> Krapp-Schickel & Ruffo, 2001													X					
	<i>Maeropsis rathbunae</i> (Pearse, 1908)	X																	
	<i>Mallacoota carausui</i> Ortiz, 1976	X																	
	<i>Mélita guanaense</i> Ortiz, Garcia-Debras & Lalana, 2002	X																	
	<i>Mélita letotelson</i> Vonk, 1989	X	X													X			
	<i>Mélita longisetosa</i> Sheridan, 1980	X												X					
	<i>Mélita nitida</i> Smith, 1873	X	X											X					
	<i>Mélita palmata</i> (Montagu, 1804)													X					
	<i>Mélita persona</i> G. Karaman, 1987													X					
	<i>Mélita planatarga</i> Kunkel, 1910		X											X					
	<i>Meximaera diffidentia</i> J.L. Barnard, 1969													X					
	<i>Netamaera barnardi</i> McKinney, Kalke & Holland, 1978													X					
	<i>Netamelita tabaci</i> Thomas & Barnard, 1991							X											
	<i>Nuamu curvata</i> Vonk, 1989																		
	<i>Nuamu muelleri</i> Ortiz, 1976	X	X												X				
	Family Melitidae Bousfield, 1973	<i>Psammogammarus bluefieldensis</i> Ortiz, Lalana & Beltran, 1993									X								
		<i>Psammogammarus longidactylus</i> Vonk & Stock, 1987														X			
		<i>Psammogammarus scopolorum</i> Stock, 1983													X				
<i>Psammomelita uncinata</i> Vonk, 1988														X					
<i>Quadrmaera cristianae</i> Krapp-Schickel & Ruffo, 2000														X	X			X	
<i>Quadrmaera inaequipes</i> (A. Costa, 1851)		X												X	X				
<i>Quadrmaera miranda</i> (Ruffo, Krapp & Gable, 2000)			X											X	X				
<i>Quadrmaera pacifica</i> (Schellenberg, 1938)														X					
<i>Quadrmaera pietieri</i> Krapp-Schickel & Ruffo, 2000		X	X											X	X				X
<i>Quadrmaera prope pacifica</i> (Schellenberg, 1938)																			
<i>Quadrmaera quadrimana</i> (Dana, 1852)		X	X												X				X
<i>Quadrmaera sarae</i> (Krapp-Schickel & Ruffo, 2000)		X	X												X				
<i>Quadrmaera serrata</i> (Schellenberg, 1938)		X	X												X				
<i>Quadrivisio luzi</i> (Shoemaker, 1933)															X				

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Melitidae Bousfield, 1973	<i>Raffomauera williamsi</i> (Bynum & Fox, 1977)	X																
	<i>Spathiopus cojimarensis</i> Ortiz & Lalana, 2002	X													X			
	<i>Thalassostygus exiguus</i> Vonk, 1990																	
Family Melphippidae Stebbing, 1899	<i>Victoriopsis guanarocana</i> Ortiz & Lalana, 1989	X																
	<i>Hornellia (Metaceradocus) atlantica</i> Thomas & Barnard, 1986					X												
	<i>Hornellia (Metaceradocus) habanensis</i> Ortiz, Lalana & Varela, 2008	X																
Family Neomegamphopidae Myers, 1981	<i>Hornellia (Metaceradocus) tequesiae</i> Thomas & Barnard, 1986													X				
	<i>Neomegamphopus heardi</i> Barnard & Thomas, 1987													X				
	<i>Neomegamphopus hiatus</i> J. L. Barnard & Thomas, 1987		X											X				
Family Neomegamphopidae Myers, 1981	<i>Neomegamphopus kalanii</i> J. L. Barnard & Thomas, 1987										X			X				
	<i>Pseudomegamphopus barnardi</i> Myers, 1968													X				
	<i>Pseudomegamphopus excavatus</i> Myers, 1968													X				
Family Ochlesidae Stebbing, 1910	<i>Curidia monticae</i> Ortiz, Lalana & Varela, 2007	X					X											
	<i>Curidia debrogania</i> Thomas, 1983		X															
	<i>Aceroides (Patoides) symparis</i> (J.L. Barnard, 1964)											X						
Family Oedicerotidae Liljeberg, 1865	<i>Americichlidium americanum</i> (Bousfield, 1973)		X				X						X					
	<i>Hartmannodes nyei</i> (Shoemaker, 1933)						X											
	<i>Monoculodes cf. carinatus</i> (Bate, 1857)	X																
Family Philantidae Stebbing, 1899	<i>Pericotulodes cerasinus</i> Thomas & J. L. Barnard, 1985		X				X										X	
	<i>Westwoodilla longimana</i> Shoemaker, 1934																	
	<i>Pariphinotus seclusus</i> (Shoemaker, 1933)	X	X				X							X				
Family Pardaliscidae Boeck, 1871	<i>Pariphinotus seticoxa</i> (Ortiz, 1976)	X																
	<i>Parpano cebus</i> J.L. Barnard, 1964												X					
	<i>Parpano compositus</i> J.L. Barnard, 1964																	
Family Phoxocephalidae Sars, 1891	<i>Eobolgus spinosus</i> Holmes, 1905	X	X											X				
	<i>Harpinia excavata</i> Chevreux, 1888																	
	<i>Harpinia intermedia</i> Bellan-Santini, 1997											X						X
Family Phoxocephalidae Sars, 1891	<i>Heterophoxus oculatus</i> (Holmes, 1908)	X												X				
	<i>Metharpinia floridana</i> (Shoemaker, 1933)	X	X								X	X	X	X				
	<i>Metharpinia oripacifica</i> J.L. Barnard, 1980										X							
	<i>Microphoxus minimus</i> J.L. Barnard, 1960										X							

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	LT	CAY
Family Phoxocephalidae Sars, 1891	<i>Proharpinia barbada</i> Bellan-Santini, 1997														X			
Family Platyschnopidae Barnard & Drummond, 1979	<i>Rhepoxymius epistomus</i> (Shoemaker, 1938)	X	X				X	X		X	X		X	X				
	<i>Eudevenopus hondurancus</i> Thomas & J. L. Barnard, 1983			X						X				X				
	<i>Eudevenopus megracilis</i> (J.L. Barnard, 1964)			X						X	X			X				
	<i>Tiburonella morrocoyensis</i> Ortiz, Martin & Ateenza, 2000						X						X	X	X	X	X	
	<i>Tiburonella viscana</i> (J.L. Barnard, 1964)												X	X	X	X	X	
Family Podoceridae Leach, 1814	<i>Podocerus brasiliensis</i> (Dana, 1853)	X	X	X										X				
	<i>Podocerus cristatus</i> Thomson, 1879													X				
	<i>Podocerus fassipes</i> Serejo, 1995													X		X		
	<i>Podocerus jarecki</i> Baldinger & Gable, 2002													X				
	<i>Podocerus kleidus</i> Thomas & J. L. Barnard, 1992	X												X				
	<i>Podocerus lazovaseni</i> Baldinger & Gable, 1994													X				
Family Scopelocheiridae Lowry & Stoddart, 1997	<i>Aroni americana</i> Lowry & Stoddart, 1997		X											X				
Family Sebidae Walker, 1908	<i>Seba robusta</i> Ortiz & Lemaitre, 1997												X					
	<i>Seba tropica</i> McKinney, 1980		X				X							X				
Family Stegocephalidae Dana, 1853	<i>Stegocephaloides calypsonis</i> Verge, Vader & Galan, 2001													X				
Family Stenothoidae Boeck, 1871	<i>Parametopella texensis</i> McKinney, Kalke & Holland, 1978		X											X				
	<i>Stenothoe gallensis</i> Walker, 1904	X	X								X			X				
	<i>Stenothoe valida</i> Dana, 1852	X											X					
Family Synopiidae Dana, 1853	<i>Brizelia perica</i> J.L. Barnard, 1972											X						
	<i>Brizetopsis turba</i> J.L. Barnard, 1964											X						
	<i>Garosyrhoe bigarra</i> (J.L. Barnard, 1962)													X				
	<i>Garosyrhoe luquei</i> Ortiz & Veledo, 1988													X				
	<i>Ilerastroe torpens</i> (J.L. Barnard, 1962)	X												X				
	<i>Latacunga comanita</i> J.L. Barnard, 1972												X					
Family Synopiidae Dana, 1853	<i>Metaitron bellairsi</i> (Just, 1981)													X		X		
	<i>Metaitron tropakis</i> (J. L. Barnard, 1972)		X				X							X		X		
	<i>Synopia scheeleana</i> Bovallius, 1886													X		X		
	<i>Synopia ultramarina</i> Dana, 1853													X		X		
	<i>Syrhoe papyracea</i> Stebbing, 1888		X										X	X		X		
	<i>Tiron biocellata</i> J.L. Barnard, 1962	X												X		X		

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY	
Family Talitridae Rafinesque, 1815	<i>Chelorchestia costaricana</i> (Stebbing, 1906)			X						X					X				
	<i>Floresorchestia guadalupensis</i> Ciavatti, 1989												X		X		X		
	<i>Orchestia grillus</i> (Bosc, 1802)												X						
	<i>Platorchestia platensis</i> (Kroyer, 1845)		X	X												X			
	<i>Pseudorchestioidea biolleyi</i> (Stebbing, 1908)									X		X							
	<i>Talorchestia frizi</i> Stebbing, 1903	X												X					
	<i>Talorchestia marcuzzi</i> Ruffo, 1950													X					
	<i>Talorchestia margaritae</i> Stephensen, 1948													X					
	<i>Talorchestia sulenstoni</i> (Stebbing, 1899)														X		X		
	<i>Telorchestia antillensis</i> Bousfield, 1984		X	X											X	X			
Family Uristidae Hurlay, 1963	<i>Telorchestia karukarae</i> Ciavatti, 1989															X			
	<i>Ullorchestia uhleri</i> (Shoemaker, 1930)		X																
	<i>Stephonyx bicayensis</i> (Chevreux, 1908)		X	X												X			
	<i>Stephonyx carinatus</i> Bellan-Santini, 1997															X			
	<i>Stephonyx incertus</i> Bellan-Santini, 1997															X			
	<i>Vemana compressa</i> J.L. Barnard, 1964										X								
	<i>Vemana lizata</i> J.L. Barnard, 1964													X					
	Suborder Hyperidea Milne Edwards, 1830																		
	Family Brachyscelidae Stephensen, 1923	<i>Brachyscelus cruscilum</i> Bate, 1861	X	X			X												
		<i>Brachyscelus globiceps</i> (Claus, 1879)	X	X											X				
Family Brachyscelidae Stephensen, 1925	<i>Brachyscelus macrocephalus</i> Stephensen, 1925	X																	
	<i>Brachyscelus rapacioides</i> Stephensen, 1925		X																
	<i>Thamneus rostratus</i> Bovallius, 1887		X																
Family Cystisomatidae Willemoes-Suhm, 1875	<i>Cystisoma latipes</i> (Stephensen, 1918)		X																
	<i>Cystisoma longipes</i> (Bovallius, 1886)		X																
Family Dairellidae Bovallius, 1887	<i>Dairella californica</i> (Bovallius, 1885)		X																
	<i>Tryphana malmii</i> Boeck, 1871		X																
Family Lyceidae Claus, 1879	<i>Iulopsis loveni</i> Bovallius, 1887		X																
	<i>Iulopsis loveni</i> Bovallius, 1887		X																
Family Iulopidae Zaidler, 2004	<i>Lanceola sayana</i> Bovallius, 1885		X																

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY	
Family Lestrigomidae Zeidler, 2004	<i>Hyperietta luzoni</i> (Stebbing, 1888)		X											X					
	<i>Hyperietta parviceps</i> Bowman, 1973		X																
	<i>Hyperietta stebbingi</i> Bowman, 1973		X													X			
	<i>Hyperietta stephensi</i> Bowman, 1973		X																
	<i>Hyperietta vosselei</i> (Stebbing, 1904)		X																
	<i>Hyperioides longipes</i> Chevreux, 1900		X														X		
	<i>Hyperionyx macrodactylus</i> (Stephensen, 1924)		X																
	<i>Lestrigonus bengalensis</i> Giles, 1887		X				X										X		
	<i>Lestrigonus crucipes</i> (Bovallius, 1889)		X																
	<i>Lestrigonus latissimus</i> (Bovallius, 1889)		X																
	<i>Lestrigonus macrophthalmus</i> (Vosselei, 1901)		X												X				
	<i>Lestrigonus schizogeneios</i> (Stebbing, 1888)		X												X				
	<i>Phronimopsis spinifera</i> Claus, 1879		X																
	<i>Themistella fusca</i> (Dana, 1853)		X																
	Family Lycaeidae Claus, 1879		X																
	Family Lycaeidae Claus, 1879	<i>Lycaea bojenensis</i> Shoemaker, 1925		X															
<i>Lycaea bovallii</i> Chevreux, 1900			X																
<i>Lycaea bovallioides</i> Stephensen, 1925			X																
<i>Lycaea pachypoda</i> (Claus, 1879)			X																
<i>Lycaea pauli</i> Stebbing, 1888			X																
<i>Lycaea pullex</i> Marton, 1874			X																
<i>Lycaea vincentii</i> Stebbing, 1888			X																
<i>Simorhynchotus antennarius</i> (Claus, 1871)			X																
Family Lycaeopsidae Chevreux, 1913			X																
Family Lycaeopsidae Chevreux, 1913		<i>Lycaeopsis themistoides</i> Claus, 1879		X															
	<i>Lycaeopsis zamboangae</i> (Stebbing, 1888)		X																
Family Mimonectidae Bovallius, 1887		X																	
Family Oxycephalidae Dana, 1853	<i>Mimonectes gaussi</i> (Woltereck, 1904)		X																
	<i>Craniocephalus scleroticus</i> (Streets, 1878)		X																
	<i>Glossocephalus mlinebwartzi</i> Bovallius, 1887		X																
	<i>Leptocotis tenuirostris</i> (Claus, 1871)		X																
	<i>Oxycephalus clausi</i> Bovallius, 1887		X																
<i>Oxycephalus piscator</i> Milne Edwards, 1830		X																	
<i>Rhabdosoma minor</i> Fage, 1954		X																	



APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Oxycephalidae Dana, 1853	<i>Rhabdosoma whitei</i> Bate, 1862	X	X													X		
	<i>Streetsia challengeri</i> Stebbing, 1888		X															
	<i>Streetsia mindanaonis</i> (Stebbing, 1888)		X															
	<i>Streetsia porcella</i> (Claus, 1879)	X	X															
	<i>Streetsia steenstrupi</i> (Bovallius, 1887)		X															
Family Paraphronimidae Bovallius, 1887	<i>Paraphronima crassipes</i> Claus, 1879		X													X		
	<i>Paraphronima gracilis</i> Claus, 1879		X													X		
Family Parascelidae Bate, 1862	<i>Parascelus edwardsi</i> Claus, 1879		X													X		
	<i>Parascelus typhoides</i> Claus, 1879		X													X		
Family Parascelidae Bate, 1862	<i>Schizoscelus ornatus</i> Claus, 1879		X															
	<i>Thyropus sphaeroma</i> (Claus, 1879)		X											X				
Family Phronimidae Rafinesque, 1815	<i>Phronima colletti</i> Bovallius, 1887	X	X															
	<i>Phronima curvipes</i> Vosseler, 1901		X															
Family Phrosinidae Dana, 1853	<i>Phronima pacifica</i> Streets, 1887		X															
	<i>Phronima sedentaria</i> (Forsk., 1775)		X															
	<i>Phronima solitaria</i> Guérin-Ménéville, 1836		X															
	<i>Phronima stebbingi</i> Vosseler, 1901	X	X															
	<i>Phronimella elongata</i> (Claus, 1862)		X															
	<i>Anchylomera blossevilleti</i> Milne Edwards, 1830		X													X		
	<i>Phrosina semilunata</i> Risso, 1882		X													X		
	<i>Prinno abyssalis</i> Bowman, 1968		X															
	<i>Prinno brevidens</i> Bowman, 1978		X															
	<i>Prinno evansi</i> Shearer, 1986		X															
Family Platyscelidae Bate, 1862	<i>Prinno johnsoni</i> Bowman, 1978		X															
	<i>Prinno latreillei</i> Stebbing, 1888		X															
	<i>Prinno macropa</i> Guérin-Ménéville, 1836		X															
	<i>Amphithyrus bispinosus</i> Claus, 1879		X															
	<i>Amphithyrus glaber</i> Spandl, 1924		X															
	<i>Amphithyrus muratus</i> Volkov, 1982		X															
	<i>Amphithyrus sculpuratus</i> Claus, 1879		X															
<i>Hemityphis tenuimanus</i> Claus, 1879	X	X																

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYL	CAY
Family Platyscelidae Bate, 1862	<i>Paratyphus maculatus</i> Claus, 1879	X	X															
	<i>Paratyphus parvus</i> Claus, 1887		X															
	<i>Paratyphus promontori</i> Stebbing, 1888	X	X															
	<i>Paratyphus spinosus</i> Spandl, 1924		X															
Family Platyscelidae Bate, 1862	<i>Platyscelus armatus</i> (Claus, 1879)		X											X				
	<i>Platyscelus crustulatus</i> (Claus, 1879)	X	X											X				
	<i>Platyscelus ovooides</i> (Risso, 1816)	X	X											X				
	<i>Platyscelus serratulus</i> Stebbing, 1888	X	X											X				
	<i>Tetrahyrus forcipatus</i> Claus, 1879	X	X											X				
	<i>Eupronoe intermedia</i> Stebbing, 1888		X															
	<i>Eupronoe laticarpa</i> Stephensen, 1925		X															
Family Pronoidea Dama, 1853	<i>Eupronoe maculata</i> Claus, 1879	X	X											X				
	<i>Eupronoe minuta</i> Claus, 1879		X											X				
	<i>Paratyphus gracilis</i> Claus, 1879		X											X				
	<i>Paratyphus hoylei</i> Stebbing, 1888		X											X				
	<i>Parapronoe crustulum</i> Claus, 1879	X	X											X				
	<i>Parapronoe parva</i> Claus, 1879		X											X				
	<i>Pronoe capito</i> Guérin-Ménéville, 1836		X											X				
Family Scimidae Stebbing, 1888	<i>Acanthoscina acanthodes</i> (Stebbing, 1895)		X											X				
	<i>Scina borealis</i> (Sars, 1882)		X											X				
	<i>Scina crassicornis</i> (Fabricius, 1775)		X											X				
	<i>Scina curvialcyla</i> Chevreux, 1914		X											X				
	<i>Scina damasi</i> Priot, 1929		X											X				
	<i>Scina hurleyi australis</i> Zeidler, 1998		X											X				
	<i>Scina indica</i> M. Vinogradov, 1964		X											X				
	<i>Scina langhansii</i> Wäglér, 1926		X											X				
	<i>Scina marginata</i> (Bovallius, 1885)		X											X				
	<i>Scina nana</i> Wäglér, 1926		X											X				
	<i>Scina pacifica</i> (Bovallius, 1887)		X											X				
	<i>Scina similis</i> Stebbing, 1895		X											X				
	<i>Scina stenopus</i> Stebbing, 1895		X											X				

APPENDIX 1 (Continued)

Suborder / Family	Species	CUB	MEX	PRC	HIS	JAM	BEL	HON	GUA	NIC	CRC	PAN	COL	VEN	ABC	LAN	TYT	CAY	
Family Scimidae Stebbing, 1888	<i>Scina submarginata</i> Tattersall, 1906		X																
	<i>Scina tullbergi</i> (Bovallius, 1885)		X																
	<i>Scina vosseleri</i> Tattersall, 1906		X																
	<i>Scina wagneri</i> atlantis Thurston, 1976		X																
	<i>Vibilia armata</i> Bovallius, 1887	X	X																
Family Vibiliidae Dana, 1853	<i>Vibilia australis</i> Stebbing, 1888	X	X											X					
	<i>Vibilia chuni</i> Behning & Woltereck, 1912		X																
	<i>Vibilia gibbosa</i> Bovallius, 1887		X																
	<i>Vibilia jeungerardi</i> Lucas, 1845	X	X																
	<i>Vibilia propinqua</i> Stebbing, 1888	X	X																
	<i>Vibilia stebbingi</i> Behning & Woltereck, 1912	X	X																
	<i>Vibilia viatrix</i> Bovallius, 1887		X															X	
	Suborder Ingolfiellidea Hansen, 1903	<i>Ingolfiella fontinalis</i> Stock, 1977														X			
		<i>Ingolfiella fuscina</i> Dojiri & Sieg, 1987		X												X			
	Family Ingolfiellidae Hansen, 1903	<i>Ingolfiella grandispina</i> Stock, 1979													X				
<i>Ingolfiella margaritae</i> Stock, 1979																			
<i>Ingolfiella putealis</i> Stock, 1976															X				
<i>Ingolfiella quadridentata</i> Stock, 1979															X				
	<i>Ingolfiella tabularis</i> Stock, 1977														X				