

Abundance, diversity and community characterization of aquatic Coleoptera in a rice field of Northeastern Argentina

María Constanza Gómez Lutz¹, Arturo I. Kehr¹ & Liliana A. Fernández²

1. Centro de Ecología Aplicada del Litoral (CCT Nordeste - CONICET- UNNE), Ruta 5 km 2.5, (3400) Corrientes, Argentina; cotygoomezlutz@hotmail.com; arturokehr@yahoo.com.ar
2. División Entomológica, Museo de La Plata (CCT - CONICET), Paseo del Bosque s/n, 1900. La Plata, Buenos Aires (Argentina); liliafer@fcnym.unlp.edu.ar

Received 13-VIII-2014. Corrected 11-III-2015. Accepted 13-IV-2015.

Abstract: Rice fields occupy large areas in Northeastern Argentina, and in Corrientes this widespread activity has become a feature in the landscape, as it is one of the main producing provinces. The aquatic Coleoptera is part of the fauna inhabiting these artificial environments but little is known about this group in irrigated rice fields. The aims of this study were to determine the diversity and species richness of coleopterans in a typical rice field, and to characterize the community of water beetles through different abundance models. For this, samples were collected from an active rice field located in “El Sombrero” town, in Corrientes Province, between November 2011 and April 2012. An entomological net of 30 cm diameter was used, and species richness, diversity and equitability were calculated monthly; besides, the community composition was characterized by means of rank-abundance models. A total of 74 species of aquatic coleopterans were identified. January and February resulted the months with the highest diversity. The aquatic Coleoptera species found in most of the sampled months were adjusted to the logarithmic rank-abundance model. The data suggests that, if it is properly managed, rice cropping in Northeastern Argentina can support a diverse aquatic coleopteran fauna. *Rev. Biol. Trop.* 63 (3): 629-638. Epub 2015 September 01.

Key words: biodiversity, aquatic coleoptera, rice field, Argentina.

In the world, approximately 151 million hectares are cultivated with rice (FAO Stat, 2008). In Argentina, irrigated rice production is carried out mainly in Corrientes and Entre Ríos Provinces because of the agroecological characteristics (water, climate and soil) of this region, and nowadays, an approximate of 164 300 hectares are planted with this crop in both Provinces (Asociación Correntina de Plantadores de Arroz y Bolsa de Cereales de Entre Ríos, 2013). The irrigated rice fields behave as temporary wetlands, with alternating periods of flooding in summer and drought in winter (Fasola & Ruiz, 1996). These rice fields act as artificial aquatic ecosystems which connect and share water with natural wetlands (rivers and water courses, groundwater or dams filled with rainwater) (Blanco & Balze, 2011). These

man-made environments, are included in the definition of wetlands adopted by the Wetlands Convention (Secretaría de la Convención de Ramsar, 2010), and harbor a great variety of aquatic invertebrates (Fernando, 1993; Fernando, Furtado, & Lim, 1979), including insects (Bambaradeniya, 2000; Bambaradeniya et al., 2004). Most aquatic invertebrates, mainly microcrustaceans, insect larvae, gastropods, and oligochaetes, contribute to nutrient cycling in rice fields, and play important roles in the decomposition of the photosynthetic aquatic biomass that develops in rice field floodwater (Simpson & Roger, 1991).

In South America, studies about aquatic insects in rice fields are scarce. Some studies about the arthropod community have been carried out in Southern Brazil (Stenert, Bacca,



Maltchik, & Rocha, 2009; Fritz et al., 2011); besides, a paper on rice fields biodiversity was conducted in South-Central Corrientes Province (Blanco & Balze, 2011). Although beetles are important components of the aquatic fauna, little is known about their ecology in these artificial ecosystems in this study area. A pioneering study by Yano, Chu, and Satô (1983) documented 117 species of water beetles belonging to 14 families in rice fields worldwide. Most of the studies on the biodiversity of aquatic insects in rice fields have been conducted in Asian countries (e.g. Rozilah & Ali, 1998; Bambaradeniya & Amerasinghe, 2003; Bahaar & Bhat, 2011), probably because Asia stands out as the largest producer and consumer of rice in the world.

Abundance and species richness of aquatic beetles have been reported to be strictly related to the physical variables of the habitat such as vegetation and the chemistry of water (e.g. Nilsson & Söderberg, 1996). Most studies in this type of environment have dealt mainly with agronomic aspects and rice pests; however, studies on biodiversity in these agroecosystems fauna remain scarce. The aims of this study were to determine the diversity and species richness of coleopterans in a typical rice field, and to characterize the community of water beetles through different abundance models.

MATERIALS AND METHODS

Study site: This study was conducted in a rice field located 30 km Southwest of Corrientes city, Argentina, in the “El Sombro-ro” town (27°42'12" S - 58°46'2" W). The site has belonged to the National Institute of Agricultural Technology (INTA) since 1958. This field has an area of ~1 175 ha. Agricultural and livestock activities are carried out in the site.

Irrigated rice production: Rice sowing began in September and October. The field was then flooded with water drawn from the Paraná River, pumped from the main and secondary channels. The water is maintained on the plots by containment edges called “taipas” until April, and it is drained for later harvest.

In the initial phase of flooding, no vegetation was observed in the irrigation channels; later on, the predominant hydrophilic vegetation consisted of *Sagittaria montevidensis* Cham. and Schlecht, *Hydrocotyle ranunculoides* L. f., *Limnobium* sp. and *Ludwigia* sp.

Sampling methods: Samples were carried out monthly during a rice cultivation cycle (November 2011-April 2012). Aquatic insects were collected with an aquatic hand net (mesh size 300 µm, diameter 30 cm) both inside the rice field and in channels that drain the water from the river into the plantation, and connect to the different sowed areas. The sampling was started immediately after the field preparation work, which coincided with spring and sowing of the rice. The random sampling technique was used in different parts of the site, usually between 10 a.m. and 1 p.m. The samples were taken manually by one person who sampled for one hour, from the cultivated parcels and irrigation channels, in order to avoid underestimation of the biodiversity. During the sampling months, water depth ranged between 5 and 10 cm in the cultivated parcels, and between 10 and 50 cm in the irrigation channels. Table 1 shows data on monthly temperature, precipitation, water temperature, conductivity, dissolved oxygen and pH for this period.

The insect samples were fixed *in situ* in formaldehyde 5 %, transferred to the laboratory for identification, and then stored in 70 % ethanol. Taxa were classified according to Lawrence and Newton (1995). Hydrophiloidea was classified following Hansen (1999) and Short and Fikáček (2011). Hydrophilidae was classified following Short and Fikáček (2013), whereas Dytiscidae was classified following Nilsson (2001). Species were identified following Trémouilles and Bachmann (1981), Trémouilles (1984, 1989, 1995), Trémouilles, Oliva and Bachmann (1995), Trémouilles, Michat and Torres (2005), Oliva, Fernández and Bachmann (2002) and Vondel and Spangler (2008). The material collected was deposited in the collection of the Centro de Ecología Aplicada

TABLE 1

Temperature (°C): minimum-maximum (mean \pm standard deviation); mean precipitation (mm) \pm SD; water temperature (°C) \pm SD, conductivity (%) \pm SD, dissolved oxygen \pm SD and pH \pm SD during a rice cultivation cycle in a rice field of Corrientes province, Argentina

Month	Temperature	Precipitation	T°C - H ₂ O	Cond. (%)	Dis. O ₂	pH
November	13.2-36.4 (24.3 \pm 2.8)	18.7 \pm 16.8	27 \pm 2.7	49 \pm 13.2	71.4 \pm 30.5	7.94 \pm 0.7
December	12.2-40.7 (25.7 \pm 3.3)	10.6 \pm 15.3	28 \pm 3.3	40 \pm 10.0	21.5 \pm 9.6	8.03 \pm 0.9
January	16.2-38.9 (27.5 \pm 2.3)	13.1 \pm 14.6	30 \pm 2.7	48 \pm 13.4	0	6.35 \pm 0.4
February	15.4-38.7 (28.8 \pm 2.7)	7.2 \pm 7.6	28 \pm 3.0	90 \pm 25.0	68.3 \pm 28.0	7.01 \pm 0.6
March	8.6-39.3 (25.5 \pm 4.7)	14.8 \pm 18.5	25.2 \pm 2.0	60 \pm 17.0	0	6.19 \pm 0.4
April	6.6-33.8 (21.15 \pm 4.1)	15.8 \pm 23.3	18 \pm 1.0	40 \pm 11.1	29.2 \pm 13.1	7.69 \pm 0.6

del Litoral (CECOAL-(CONICET-UNNE)), Corrientes Province.

Only adult individuals were taken into consideration, for different analysis. Richness and diversity species were calculated on a monthly basis. The diversity was calculated by the Shannon diversity index (Shannon & Weaver, 1949). Rarefaction methods were first used to compare the average diversity and the average richness between the different months. Rarefaction uses a probability theory to derive expressions for the expectation and variance of diversity and richness of species for a sample of a given size. This method lowers the sample to a common abundance level and then compares species diversity and richness. Ecosim 7.0 software (Gotelli & Entsminger, 2004) was used for this calculation. The richness and diversity results were obtained for 21 classes of abundance (10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200 and 205 individuals). Subsequently, the diversity and species richness for each of the 21 classes of abundance was compared through the non-parametric Kruskal-Wallis test and then by the Dunn test with the Bonferroni correction. The statistical analyses were calculated with XIStat 7.5 software (Addinsoft, 2006).

The equitability for each month sampled was calculated by the formula of the Shannon equitability index (Magurran, 2004).

The community composition for each sampling date was characterized by means of rank-abundance models. Three of the most common

series (logarithmic, geometric or “broken stick”) presented a better adjustment to the data that was specified for each case. The different variables were compared by means of the Chi-square test. Every calculation was made with the Bio-Dap software (Thomas & Clay, 2000), which is based on the methods proposed by Magurran (1988).

RESULTS

A total of 74 species of Coleoptera, belonging to 28 genera and seven families, were identified. Dytiscidae was the richest genera family, while Hydrophilidae was the richest species family (Dytiscidae: 13 genera; 22 species, Hydrophilidae, 7 genera; 36 species). The following richest families were Noteridae (3 genera; 9 species), Dryopidae (2 genera; 3 species), Hydrochidae (1 genus; 2 species), Haliplidae (1 genus; 1 species) and Gyrinidae (1 genus; 1 species) (Table 2).

In order to compare richness and diversity of the aquatic Coleoptera community among the six months sampled, the parameters were rarefied to the smallest individual number (April, N = 205). Significant differences were observed in diversity and species abundance in the six months analyzed (Kruskal-Wallis Test; K = 66.94, d.f. = 5, P < 0.05, and K = 19.37, d.f. = 5, P < 0.05, respectively). According to the Dunn Test with Bonferroni correction (= 0.0033), the diversity in January and February was significantly higher than in the other months, where diversity showed similar lower

TABLE 2
 Aquatic Coleoptera species collected in a rice field located in Corrientes province (Argentina)
 from November 2011 to April 2012

	November	December	January	February	March	April
ADEPHAGA						
GYRINIDAE						
<i>Gyrinus ovatus</i>	6	2	12	0	0	0
HALIPLIDAE						
<i>Haliphus maculicollis</i>	0	0	0	1	3	0
DYTISCIDAE						
Copelatinae						
Copelatini						
<i>Copelatus alternatus</i>	0	1	0	1	2	0
<i>Copelatus longicornis</i>	1	0	0	0	11	0
Dytiscinae						
Aciliini						
<i>Thermonectus succinctus</i>	4	2	8	4	4	1
<i>Thermonectus nobilis</i>	0	0	1	0	0	0
Cybistrini						
<i>Megadytes laevigatus</i>	0	0	1	1	0	0
<i>Megadytes</i> sp.	0	0	1	0	0	0
Hydroporinae						
Bidessini						
<i>Anodocheilus maculatus</i>	0	0	4	0	0	0
<i>Hemibidesus conicus</i>	0	0	0	0	2	1
<i>Liodesus</i> sp.	0	4	1	6	1	0
Hydrovatini						
<i>Hydrovatus caraibus</i>	0	0	0	3	0	7
<i>Hydrovatus turbinatus</i>	0	0	0	0	0	1
Hyphydrini						
<i>Desmopachria concolor</i>	1	27	9	0	6	2
<i>Desmopachria</i> sp.	0	0	1	0	0	0
<i>Pachydrus globosus</i>	0	1	2	1	0	10
<i>Pachydrus obesus</i>	1	0	0	0	0	0
Methlini						
<i>Celina</i> sp. 2	0	1	0	0	0	0
Vatellini						
<i>Vatellus</i> sp.	0	11	3	3	0	0
<i>Derovatellus</i> sp.	0	1	0	1	6	1
Laccophilinae						
Laccophilini						
<i>Laccophilus</i> sp. 1	6	18	67	16	22	44
<i>Laccophilus</i> sp. 2	16	5	1	2	1	3
<i>Laccophilus</i> sp. 3	0	6	15	3	3	0
<i>Laccophilus</i> sp. 4	0	0	3	0	0	0
NOTERIDAE						
Noterinae						
Noterini						
<i>Hydrocanthus sharpi</i>	0	0	0	1	0	0

TABLE 2 (Continued)

	November	December	January	February	March	April
<i>Hydrocanthus debilis</i>	47	0	0	0	7	10
<i>Hydrocanthus paraguayensis</i>	1	0	7	0	2	4
<i>Suphis maculicollis</i>	2	0	0	0	0	0
<i>Suphis fluviatilis</i>	0	0	0	1	0	0
<i>Suphisellus nigrinus</i>	2	10	6	0	0	3
<i>Suphisellus flavopictus</i>	0	0	0	0	0	1
<i>Suphisellus</i> sp. 3	2	0	0	2	0	0
<i>Suphisellus</i> sp. 4B	18	25	16	7	0	1
POLYPHAGA						
DRYOPIDAE						
<i>Pelonomus</i> sp. 1	14	4	1	0	1	2
<i>Pelonomus</i> sp. 2	1	0	0	0	0	0
<i>Onopelmus</i> sp.	0	0	0	0	1	0
HYDROCHIDAE						
<i>Hydruchus</i> sp. 1	1	0	1	3	1	3
<i>Hydrochus</i> sp. 2	0	0	0	0	0	2
HYDROPHILIDAE						
Hydrophilinae						
Berosini						
<i>Derallus angustus</i>	38	8	3	1	1	1
<i>Derallus paranensis</i>	1	0	30	5	2	10
<i>Berosus rufulus</i>	0	0	0	0	3	9
<i>Berosus reticulatus</i>	0	0	0	138	251	1
<i>Berosus truncatipennis</i>	0	0	0	41	0	0
<i>Berosus stenocoptus</i>	0	0	0	3	0	0
<i>Berosus patruelis</i>	0	0	0	44	292	1
<i>Berosus minimus</i>	0	0	0	7	5	1
<i>Berosus unguidentatus</i>	0	0	0	1	0	0
<i>Berosus pedregalensis</i>	0	0	0	0	2	0
<i>Berosus</i> sp. 1	0	0	0	1	22	3
<i>Berosus</i> sp. 13	0	0	0	0	29	1
<i>Berosus</i> sp. 14	0	76	3	1	0	0
Laccobiini						
<i>Paracymus rufocinctus</i>	0	4	0	53	0	0
<i>Paracymus limbatus</i>	0	0	0	2	0	0
<i>Paracymus graniformis</i>	0	0	0	2	1	1
<i>Paracymus granulum</i>	0	0	0	0	0	2
<i>Paracymus</i> sp. 2	0	0	1	0	0	0
Hydrophilini						
<i>Tropisternus collaris</i>	3	0	5	9	0	1
<i>Tropisternus ovalis</i>	36	0	8	3	5	1
<i>Tropisternus lateralis</i>	8	16	5	38	11	0
<i>Tropisternus laevis</i>	10	3	17	112	6	72
<i>Tropisternus apicipalpis</i>	0	1	0	1	0	0
<i>Tropisternus longispina</i>	0	0	2	0	0	0
<i>Tropisternus burmeisteri</i>	0	0	3	4	9	0
<i>Tropisternus mergus</i>	0	0	0	13	0	0
<i>Tropisternus carinispina</i>	0	0	0	0	12	0

TABLE 2 (Continued)

	November	December	January	February	March	April
<i>Tropisternus latus</i>	0	0	0	0	40	0
Enochrinae						
<i>Enochrus vulgaris</i>	1	2	0	43	7	0
<i>Enochrus variegatus</i>	0	0	0	0	6	0
<i>Enochrus sublongus</i>	0	0	0	0	0	1
<i>Enochrus circumcinctus</i>	0	0	0	0	0	1
Acidocerinae						
<i>Helobata larvalis</i>	0	0	1	0	1	3
<i>Helochaeres femoratus</i>	0	0	0	1	2	0
<i>Helochaeres tectiformis</i>	0	0	1	0	0	0
<i>Helochaeres spatulatus</i>	1	0	0	0	0	0

values. The species richness during December and March was significantly higher than in April, whereas in the other months was similar (Table 3).

TABLE 3

Richness parameters summary (N): number of individuals; (S): species richness; (H): Shannon Index, and (J'): Equitability Index. Dunn test with Bonferroni correction for diversity results (calculated using the Shannon Index) and species richness in the rice field

Months	N	S	H'	J'
November	221	24 _{a,b}	2.43 _{b,c}	0.77
December	228	22 _a	2.37 _b	0.77
January	239	32 _{a,b}	2.68 _c	0.77
February	579	39 _{a,b}	2.53 _{b,c}	0.69
March	780	36 _a	2.00 _a	0.56
April	208	33 _b	2.42 _b	0.69

*Same letters indicate no significant differences; different letters indicate significant differences between groups.

The greatest number of species was recorded in February (S = 39); and the highest abundance was observed in March (N = 780) (Table 3). The greatest diversity was recorded in January (H' = 2.68) and the lowest in March (H' = 2.00). The Shannon equitability index indicated that evenness of most of the months sampled was high (J' < 0.69), except for March (J' = 0.56) (Table 3).

The distribution of species abundance in the different months was adjusted to a log-series model; nevertheless, the April distribution of species abundance did not fit any of the series models (Table 4). The test of goodness of fit (Chi-square test) indicated that no other model fitted well the data of species distribution in the others analyzed communities.

TABLE 4

Rank - abundance model results of aquatic Coleoptera in the different sampled months, where: (S) species richness; (d.f.) degrees of freedom; (Prob. X²) X² probability

Months	S	Geometric series			Log series			Broken-Stick		
		df	X ²	Prob X ²	df	X ²	Prob X ²	df	X ²	Prob X ²
Nov. 11	24	23	34.23	0.06190484	5	3.47	0.62793179	5	27.54	4.47641E-05
Dec. 11	22	21	50.35	0.00032604	6	2.45	0.87402096	6	57.28	1.60334E-10
Jun. 12	32	31	94.27	2.6449E-08	6	4.68	0.58546012	6	517.94	1.147E-108
Feb. 12	39	38	262.88	2.0439E-35	7	10.06	0.18519875	7	404.7	2.34188E-83
Mar. 12	36	35	950.42	7.416E-177	8	10.93	0.20569713	8	32 054.5	0
Apr. 12	33	32	233.37	1.8716E-32	6	16.54	0.01113112	6	5 435.45	0

DISCUSSION

Previous studies recognize the Coleoptera as the **most diverse insect order** living in rice fields in the province Corrientes (Blanco & Balze, 2011). In the present study, the aquatic coleopteran assemblage was characterized by the presence of Dytiscidae, Hydrophilidae, Noteridae, Dryopidae, Hydrochidae, Haliplidae and Gyrinidae, also reported in several rice field studies (e.g. Leitão, Pinto, Pereira, & Brito, 2007; Stenert et al., 2009; Bahaar & Bhat, 2011). However, the most representative families regarding their abundance were Hydrophilidae, Dytiscidae and Noteridae. Similar results were found in a study carried out in two permanent ponds in Corrientes province, where eight families were recorded, and where Hydrophilidae (48%), Dytiscidae (27%) and Noteridae (18%) showed the highest species percentage (Gómez Lutz, Fernández, & Kehr, 2012).

Hydrophilidae has also been reported as the family with significantly higher number of species among aquatic Coleoptera (Bahaar & Bhat, 2011; Gómez Lutz et al., 2012), and *Berosus* and *Tropisternus* have been the most important genera. Similar results have been obtained in permanent ponds from Corrientes Province, where both genera had more species number inside Hydrophilidae (Gomez Lutz et al., 2012). *Berosus reticulatus*, *Berosus patruellis* and *Tropisternus laevis* were the most abundant during February and March in this study. It is possible that rice fields in this area provide similar natural environmental conditions to the aquatic Coleoptera community.

Different studies on the fauna have also showed that these artificial environments may harbor a fauna similar to undisturbed wetlands (Bambaradeniya et al., 2004; Duré, Kehr, Schaefer, & Marangoni, 2008; Maltchik, Rolon, Stenert, Machado, & Rocha, 2011). These rice fields are a refuge to a great variety of organisms, all well adapted to this temporarily and highly manipulated ecosystem that is periodically disrupted by various agricultural practices (Fernando, 1995). The community

of water beetles was rich and abundant during the study, and similar to one reported for a national park (128 species included in 44 genera and seven families) and natural environments of this Province (127 species included in 40 genera and eight families) (Torres et al., 2012; Gómez Lutz et al., 2012). Most of the months sampled showed high diversity; the highest diversity index was observed in January. March was the month with the highest species richness but also the month with the lowest value of diversity. In contrast, January showed an intermediate number of species but the highest diversity index of all months. All these species were similarly represented. The equitability test was generally high, which showed a heterogeneous representation in the number of individuals within the community. However, it is important to note that this lack of homogeneity could be given by an explosion in the abundance of two *Berosus* species in that month. January and February are the most favorable months for studies related with the coleopteran communities in the rice field, since during these months the highest richness and diversity were registered.

The rank-abundance models aided the interpretation of the data. The most commonly used models are the geometric, the logarithmic and the “broken stick”; all of them were required to fit data from the different months of rice cultivation. The geometric series describes communities of highly uneven species-abundance distribution and low diversity characterized by a few dominant species; the “broken stick” model characterized by reflecting a more equitable condition among species and for having a more uniform distribution (Magurran, 1988); and the logarithmic series fitted better to communities characterized as being influenced by only one or a few determining factors affecting ecological interactions, with a high proportion of species found to be very uncommon (Magurran, 1988). The analyzed communities in most of the months (except April) fitted only the rank-abundance models of the logarithmic series. This is where the species

with intermediate abundance were those which dominated the different analyzed communities.

In a study of artificial environments in Argentina, Schnack, Francesco, Colado, Novoa, and Schnack (2000) suggested that these environments would contribute preserve the diversity. Our results on the diversity and richness of water beetles species in rice fields, showed that these can function as important systems that may contribute to biodiversity conservation, especially in those widespread areas where these environments have become part of the landscape.

The results suggest that the rice irrigation fields in Corrientes Province can host a wide richness of aquatic Coleoptera. This feature would be justified because the community structure was characterized by being conformed by both species, common and uncommon, (logarithmic model features) with similar abundances and slightly fluctuating. Therefore, these environments could be an alternative for biodiversity conservation in the region where the rice fields expand rapidly, due to the optimal production characteristics of soil and climate in the province. Understanding the factors regulating coleopteran abundance and diversity in agricultural areas will help to elucidate possible effects of human activities and the perturbations on natural populations.

RESUMEN

Abundancia, diversidad y caracterización de la comunidad de coleópteros acuáticos en una plantación de arroz al noreste de Argentina. Las plantaciones de arroz ocupan grandes áreas en el NE argentino. Corrientes es una de las principales provincias productoras de este cereal y al ser un cultivo extensivo ocupa grandes áreas, las cuales pasan a ser una parte característica del paisaje. Los coleópteros acuáticos forman parte de la fauna que habita en estos ambientes artificiales y poco se sabe de la fauna que habita en ellos. Los objetivos de este estudio fueron determinar la diversidad y la riqueza de especies de coleópteros en un campo típico de arroz, y caracterizar la comunidad de escarabajos de agua a través de los diferentes modelos de abundancia. Para ello, se realizaron muestreos con una red entomológica de 30 cm de diámetro en un campo de arroz ubicada en la localidad “El Sombrero”, en la provincia de Corrientes. Las muestras fueron recolectadas entre noviembre 2011 y abril 2012. La

riqueza de especies, diversidad y equitatividad para todos los meses muestreados fueron calculadas. La composición de la comunidad se caracterizó por medio de modelos de ranking-abundancia. Se identificaron 74 especies de coleópteros acuáticos. Enero y febrero fueron los meses con mayor diversidad. Las especies de coleópteros acuáticos que se registraron en la mayoría de los meses muestreados se ajustaron al modelo logarítmico. Los datos sugieren que, si se gestiona adecuadamente, los cultivos de arroz en el noreste de Argentina pueden albergar una fauna diversa de coleópteros acuáticos.

Palabras clave: biodiversidad, coleópteros acuáticos, campos de arroz, Argentina.

REFERENCES

- Addinsoft. (2006). XLSTAT (Versión 7.5 for Excel interface). U. K.: Addinsoft.
- Asociación Correntina de Plantadores de Arroz y Bolsa de Cereales de Entre Ríos. (2013). *Relevamiento Nacional-Campaña 2012-2013* Retrieved from <http://www.acpaarrozcorrientes.org.ar/>
- Bahaar, S. W. N., & Bhat, G. A. (2011). Aquatic Bio diversity in the paddy fields of Kashmir valley (J and K), India. *Asian Journal of Agricultural Research*, 5, 269-276.
- Bambaradeniya, C. N. B. (2000). *Ecology and biodiversity in an irrigated rice field ecosystem in Sri Lanka* (Ph.D. Thesis). University of Paradeniya, Sri Lanka.
- Bambaradeniya, C. N. B., & Edirisinghe, J. P. (2003). *Biodiversity associated with the rice field agro- ecosystem in Asian countries: a brief review*. Sri Lanka: Working paper 63. International Water Management Institute (IWMI).
- Bambaradeniya, C. N. B., Edirisinghe, J. P., de Silva, D. N., Gunatilleke, C. V. S., Ranawana, K. B., & Wijekoon, S. (2004). Biodiversity associated with an irrigated rice agro-ecosystem in Sri Lanka. *Biodiversity and Conservation*, 13, 1715-1753.
- Blanco, D. E. & Balze, V. M. (2011). *Conservación de los recursos acuáticos y la biodiversidad en arrozceras del noreste de Argentina*. Buenos Aires: Fundación para la conservación y el uso sustentable de los humedales.
- Duré, M. I., Kehr, A. I., Schaefer, E. F., & Marangoni, F. (2008). Diversity of amphibians in rice fields from Northeastern Argentina. *Interciencias*, 33, 523-527.
- FAO STAT. (2004). *FAO statistical database*. Retrieved from <http://apps.fao.org>.
- Fasola, M., & Ruiz, X. (1996). The value of rice field as substitutes for natural wetlands for waterbirds in the mediterranean region. *Colonial Waterbirds (Special Publication 1)*, 19, 123-128.

- Fernando, C. H. (1993). Rice field ecology and fish culture - an overview. *Hydrobiologia*, 259, 91-113.
- Fernando, C. H. (1995). Rice fields are aquatic, semi-aquatic, terrestrial and agricultural: A complex and questionable limnology. In K. H. Timotius, & F. Goltenboth (Eds.), *Tropical Limnology* (pp. 121-148). Indonesia: Satya Wacana Christian University, Salatiga.
- Fernando, C. H., Furtado, J. I., & Lim, R. P. (1979). Aquatic fauna of the world's rice fields. *Welleaceana Supplement Kuala Lumpur*, 2, 1-105.
- Fritz, L. L., Heinrichs, E. A., Machado, V., Andreis, T. F., Pandolfo, M., de Salles, S. M., de Oliveira, J. V., & Fiuza, L. M. (2011). Diversity and abundance of arthropods in subtropical rice growing areas in the Brazilian south. *Biodiversity Conservation*, 20, 2211-2224.
- Gómez Lutz, M. C., Fernández, L. A., & Kehr, A. I. (2012). Coleópteros acuáticos de lagunas situadas en el noroeste de la provincia de Corrientes, Argentina. *Revista de la Sociedad Entomológica Argentina*, 71, 73-85.
- Gotelli, N., & Entsminger, G. (2004). (Version 7.0). Software Ecosim. Null Models Software for Ecology. Acquired Intelligence Inc and Kesey-Bear. Retrieved from <http://together.net/~gentsmin/ecosim.htm>
- Hansen, M. (1999). Hydrophiloidea (s. str.) (Coleoptera). In M. Hansen (Ed.), *World Catalogue of Insects* (Vol. 2, pp. 1-416). Stenstrup, Denmark: Apollo Books.
- Lawrence, J. F., & Newton, Jr. A. F. (1995). Families and subfamilies of Coleoptera (with selected genera, notes, references and data on family-group names). In J. Pakaluk, & S. A. Slipinski (Eds.), *Biology, Phylogeny, and classification of Coleoptera: Papers Celebrating the 80th Birthday of Roy A. Crowson* (pp. 779-1092). Warszawa: Muzeum i Instytut Zoologii.
- Leitão, S., Pinto, P., Pereira, T., & Brito, M. F. (2007). Spatial and temporal variability of macroinvertebrate communities in two farmed Mediterranean rice fields. *Aquatic Ecology*, 41, 373-386.
- Magurran, A. E. (1988). *Ecological Diversity and its Measurement*. Princeton, N.J., U.S.A.: Princeton University Press.
- Magurran, A. E. (2004). *Measuring Biological Diversity*. Oxford, U.K.: Blackwell Publishing.
- Maltchik, L., Rolon, A. S., Stenert, C., Machado, I. F., & Rocha, O. (2011). Can rice field channels contribute to biodiversity conservation in Southern Brazilian wetlands? *Revista de Biología Tropical*, 59, 1895-1914.
- Nilsson, A. N. (2001). Dytiscidae (Coleoptera). In A. N. Nilsson (Ed.), *World Catalogue of Insects* (Vol. 3, pp. 1-395). Stenstrup, Denmark: Apollo Books.
- Nilsson, A. N., & Söderberg, H. (1996). Abundance and species richness patterns of diving beetle (Coleoptera: Dytiscidae) from exposed and protected sites in 98 northern Swedish lakes. *Hydrobiologia*, 321, 83-88.
- Oliva, A., Fernández, L. A., & Bachmann, A. O. (2002). Sinopsis de los Hydrophiloidea acuáticos de la Argentina (Insecta, Coleoptera). *Monografías del Museo Argentino de Ciencias Naturales*, 2, 1-67.
- Rozilah, I., & Ali, A. B. (1998). Aquatic insect populations in the Muda rice agroecosystem. In B. N. Nashriya, N. K. Ho, B. S. Ismail, B. A. Ahyaudin, & K. Y. Lum (Eds.), *Rice Agroecosystem of the Muda Irrigation Scheme, Malaysia* (pp. 97-110). Malaysia: Malaysian Institute of Nuclear Technology Research and Muda Agricultural Development Authority.
- Secretaría de la convención de RAMSAR. (2010). *Inventario de humedales: Marco de Ramsar para el inventario y la descripción de las características ecológicas de los humedales*. Manuales Ramsar para el uso racional de los humedales (4ª ed., Vol. 15). Retrieved from http://www.ramsar.org/pdf/lib/lib_rtr01_s.pdf
- Shannon, C. E., & Weaver, W. (1949). *The Mathematical Theory of Communications*. Urbana: University of Illinois Press.
- Schnack, J. A., Francesco, F. O., Colado, U. R., Novoa, M. L., & Schnack, E. J. (2000). Humedales antrópicos: su contribución para la conservación de la biodiversidad en los dominios subtropical y pampásico de la Argentina. *Ecología Austral*, 10, 63-80.
- Short, A. E. Z., & Fikáček, M. (2011). World catalogue of the Hydrophiloidea (Coleoptera): additions and corrections II (2006-2010). *Acta Entomologica Musei Nationalis Pragae*, 51, 83-122.
- Short, A., & Fikáček, M. (2013). Molecular phylogeny, evolution and classification of the Hydrophilidae (Coleoptera). *Systematic Entomology*, 38, 723-752.
- Simpson, I. C., & Roger, P. A. (1991). The impact of pesticides on non target aquatic invertebrates in wetland ricefields: A Review. In P. A. Pingali, & P. A. Roger (Eds.), *Impact of pesticides on farmer health and the rice environment* (pp. 249-270). Philippines: Kluwer Academic Publishers.
- Stenert, C., Bacca, R. C., Maltchik, L., & Rocha, O. (2009). Can hydrologic management practices of rice fields contribute to macroinvertebrate conservation in Southern Brazil wetlands? *Hydrobiologia*, 635, 339-350.
- Thomas, G., & Clay, D. (2000). *BIODAP-ecological diversity and its measurement. Resource Conservation*. New Brunswick, Canada: Fundy National Park. Retrieved from <http://nhsbig.inhs.uiuc.edu/populations/bio-dap.zip>
- Torres, P. L. M., Michat, M. C., Libonatti, M. L., Fernández, L. A., Oliva, A., & Bachmann, A. O. (2012). Aquatic Coleoptera from Mburucuyá National Park



- (Corrientes Province, Argentina). *Revista de la Sociedad Entomológica Argentina*, 71(1-2), 57-71.
- Trémouilles, E. R. (1984). El género *Rhantus* Dejean en la Argentina (Coleoptera, Dytiscidae). *Physis*, 42, 9-24.
- Trémouilles, E. R. (1989). Contribución para el conocimiento del género *Thermonectus* Dejean en la Argentina y áreas limítrofes. *Revista de la Sociedad Entomológica Argentina*, 46, 95-115.
- Trémouilles, E. R. (1995). Insecta, Coleoptera, Dytiscidae: Methlinae - Hydroporinae. In *Fauna de agua dulce de la República Argentina*. FECIC, Buenos Aires, 37, 1-82.
- Trémouilles, E. R., & Bachmann, A. O. (1981). La tribu Cybisterini en la Argentina (Coleoptera, Dytiscidae). *Revista de la Sociedad Entomológica Argentina*, 39, 101-125.
- Trémouilles, E. R., Oliva, A., & Bachmann, A. O. (1995). Insecta Coleoptera. In E. C. Lopretto, & G. Tell (Eds.), *Ecosistemas de aguas continentales, Metodologías para su estudio* (pp. 1133-1197). La Plata: Ediciones Sur.
- Trémouilles, E. R., Michat, M. C., & Torres, P. L. M. (2005). A synopsis of the South American *Hydrovatus* (Coleoptera: Dytiscidae: Hydroporinae), with notes on habitat and distribution, and a key to species. *Revista de la Sociedad Entomológica Argentina*, 64, 61-69.
- Vondel Van, B. J., & Spangler, P. J. (2008). Revision of the Haliplidae of the Neotropical Region including Mexico (Coleoptera: Haliplidae). *Koleopterologische Rundschau*, 78, 69-194.
- Yano, K., Chu, Y. I., & Satô, M. (1983). Faunal and biological studies on the insects of paddy fields in Asia. XI. Records on Aquatic Coleoptera from paddy water in the world. *Chinese Journal of Entomology*, 3, 15-31.