

Does damming of the Colorado River affect the nursery area of blue shrimp *Litopenaeus stylirostris* (Decapoda: Penaeidae) in the Upper Gulf of California?

Eugenio Alberto Aragón-Noriega^{1,2} and Luis Eduardo Calderón-Aguilera¹

1 Centro de Investigación Científica y Educación Superior de Ensenada Km 107 Carr. Tijuana Ensenada, Ensenada, Baja California, 22800 México.

2 Centro de Investigaciones Biológicas del Noroeste. Apdo. Postal 349 Guaymas, Sonora, 85400 México.
Mailing Address: P. O. Box 43-4844 San Diego, CA. 92143-4844, USA. Phone 01 52 (6) 174-5050.
Fax 01 52 (6) 175 0545. E-mail: earagon@cicese.mx aaragon@cibnor.mx

Received 4-VIII-1999. Corrected 24-V-2000. Accepted 16-VI-2000.

Abstract: After damming the Colorado River the freshwater flow was reduced to 1 % of its virgin flow to the Upper Gulf of California (UGC). The ecological effects need to be properly documented. The UGC is the nursery area for *Litopenaeus stylirostris*, the most profitable fishery in the zone. In order to know the relative abundance of *L. stylirostris* postlarval stage we conducted a sampled survey every 14 days in 1993, 1994 and 1997, plus an intensive sampling during a complete tide cycle in July 1995 and 1996. We did 10 min trawls each hour during the flood tide. Relative abundance of postlarvae was higher ($p < 0.05$) in those years when freshwater flow reached the UGC.

Key words: Upper Gulf of California, Colorado River, shrimp postlarvae, *Litopenaeus stylirostris*, abundance.

The impact of man-made land forms such a dam can drastically change the condition of any ecosystem. Large reservoirs serve as sediment traps for much of the eroded material, disrupting the sediment budgets of the drainage systems below the dams (Dolan *et al.* 1974).

Before the construction of the Hoover Dam, the Colorado River discharged about 16 600 hm³ of water in the Gulf of California. Today less than 1 % of its virgin flow ever reaches the mouth of the Colorado River (Turner and Karpiscak 1980). Flow has been regulated since storage began February 1, 1935. Mexico built Morelos Dam at the end of the Colorado River, in 1950. It is 1.1 mi south of the northerly international boundary with Mexico and 21.9 mi north of the southerly international boundary (Fig. 1).

It is commonly acknowledged that damming the Colorado River was detrimental to the environment of the Upper Gulf of California (UGC). This is a nursery area for many commercially important species. However, those ecological effects have not been properly documented. Alvarez-Borrego *et al.* (1975) and Lavín *et al.* (1998) studied the hydrography of the UGC. Those authors reported that, after null input of freshwater, the UGC can be regarded as an inverse estuary, and salinity in these conditions are higher in the North (39 ‰) than in the South (35.5 ‰). However, in a later study it was found that when up stream dams overflow because of abnormal snow melts in the Basin of the Colorado River and/or because heavy rain during El Niño events, a considerable amount of freshwater discharges




Fig. 1. Colorado River Basin, with dams and aqueducts in the Upper Gulf of California. Sampling locations are also shown (After Lavín and Sánchez 1999).

in UGC. The hydrography of the area shifts severely to estuary conditions; salinity then is lower (32 ‰) in the North than in the South (35.4 ‰) (Lavín and Sánchez 1999).

The shrimp fishery is the most profitable fishery in the UGC, supported by two species, brown shrimp *Farfantepenaeus cali-*

forniensis (Holmes 1900) and blue shrimp *Litopenaeus stylirostris* (Stimpson 1871). The latter represent over 80 % of total shrimp catch (Mathews 1981). The blue shrimp peak spawning occurs in June and July, and postlarvae use the Colorado River delta as a nursery area.

It is widely accepted that penaeid shrimp postlarval stage is critical in their life span, because it is affected by higher natural mortality. Consequently, catches depend on the strength of the year class. We, therefore, need to acquire knowledge on the dynamics of the nursery area to understand the dynamics of postlarval development. In order to assess the abundance of blue shrimp postlarvae in the nursery area of the UGC, we conducted a five year sampling survey during the spawning period of shrimp.

MATERIALS AND METHODS

Shrimp postlarvae are found in the shallow zone, close to the shore at depth of less than 2.0 m, so they have to be collected manually. We used a custom designed plankton net (505 *m) attached to a stainless steel support, trawled by two persons. The net had a calibrated flowmeter in its mouth to estimate water volume filtered. We did 10 min trawls each hour during the flood tide (tides in the Gulf of California are semi diurnal, so we sampled day and night, fig 2). Surveys were conducted every 14 days in 1993, 1994 and 1997, plus an intensive sampling during a complete tide cycle from July 12 -27, 1995 and from June 30 - July 16, 1996 in a location on the peninsula near San Felipe, Baja California (Fig. 1). Species identification was done after Mair (1979) and Calderón-Perez *et al.* (1989), checked with lab reared postlarvae from known progenitors.

Relative abundance of postlarvae (Pl/m³) was used to compare differences between years. Data were monthly averages of the spawning peak months, June and July, of each year from 1993 through 1997. A Tukey test or honestly significant difference procedure were performance after a significant ANOVA shows differences in abundance between years.

Our sampling period included two years (1993, 1997) with river discharge and three years without discharge, a more normal condition. In order to correlate postlarvae relative abundance against Colorado River water flow,




Fig 2. Tide height in mm at location during the sampling period.

we compared the monthly average of Colorado River flow, from January to June with postlarval abundance for 1993 – 1997.

The Colorado River water flow data at the Southern International Border, were obtained from the United States Bureau of Reclamation, Yuma, Arizona.

RESULTS

Relative abundance of postlarvae was higher (Tukey $p < 0.05$) in 1993 than in the other years. Abundances in 1994-1996 were the lowest and formed a different group (Table 1). Lower salinity was on those years where abundance was higher: 1993 and 1997. Temperature was not significantly different for the same months in all five years (ANOVA $p = 0.708$).

The Colorado River discharged an unusually high amount of freshwater into the UGC on the years 1993 and 1997. During 1994 to 1996 no discharge was observed. The relative abundance of postlarvae had the same pattern of the river flow. The correlation coefficient was high and significant ($r = 0.8815$; $p < 0.05$)

DISCUSSION

Undoubtedly the river is responsible for the salinity condition in the UGC. Carvajal *et al.* (1997) developed a numerical model to predict salinity as function of the river flow and to assess the environmental impact of the reduction of river discharge on the area.

TABLE 1

Colorado River flow amount crossing Mexico-US border (monthly average from January throughout June) and Litopenaeus stylirostris postlarvae relative abundance (PI/m³) (average of June and July) from 1993 throughout 1997.

Year	River flow (X 10 ⁶ m ³)	Postlarvae relative abundance (PI/m ³)	Standard error
	Average	Average	
1993	312.01	43.6 a	13.6
1994	67.28	11.63 b	1.35
1995	76.25	11.20 b	2.25
1996	71.42	16.01 b	3.37
1997	115.65	33.32 c	8.06

Values with different letters are significantly different at $p < 0.05$

During 1993 and 1997 the amount of freshwater discharged by the river diluted the salinity of the UGC. Low salinity is reported as preferred for blue shrimp on Sinaloan coastal lagoons (Mair 1980). Studies have shown the effect of salinity on growth rate of penaeid postlarvae (Rothlisberg 1998).

On other penaeid shrimps low salinity signals postlarvae to find the way to a nursery area (Hughes 1969, Mathews *et al.* 1991). In the UGC, an inverse estuary, there is no low salinity as a signal, yet postlarvae nurse in that area. Mair *et al.* (1982) conducted a laboratory experiment which demonstrated that shrimp selected the estuary water over marine water even with the same salinity. They mentioned that the "taste" of water may be important, because terrigenous material is a factor that

postlarvae select for within nursery areas where enough food is available. Whether postlarvae are adapted to low salinity or follow the water taste, our data show that river inflow prompts shrimp postlarvae to use the improved UGC as a nursery area.

Another important factor is that expanded habitat is available when the Colorado River flow increases, because freshwater floods areas that are dry on those years when river discharge is null or too low.

On years 1994 through 1996 the salinity in UGC was higher than marine water. Presence of postlarvae was still observed during this period, but at lower concentration than in those years when the Colorado river discharged water. Even though our data are just for five years and the correlation is weak, there is a

significant positive effect of the river water over nursery area. Freshwater discharged by the Colorado river lowers salinity, and possibly expands shrimp postlarvae habitat, which contain more food and sheltered areas.

ACKNOWLEDGMENTS

The authors thank Miguel Cisneros, Artemisa Castro and anonymous reviewers for their comments and suggestions. National Council of Science and Technology of Mexico (REF. CONACYT 3876-T) supported the present study, and granted a fellowship to the first author (CONACYT 92829).

RESUMEN

El represamiento del Río Colorado ha ocasionado que el flujo de agua dulce sobre el Alto Golfo de California (AGC) se haya reducido hasta el 1 % del flujo original. Se ha documentado el efecto de la reducción de agua dulce sobre las condiciones hidrográficas del AGC, pero las repercusiones ecológicas no se han descrito apropiadamente. El AGC ha sido área de crianza para especies comerciales como el camarón *Litopenaeus stylirostris*. Se hicieron recolectas de postlarvas de *L. stylirostris* en el AGC durante cinco años consecutivos. Los muestreos fueron catorcenalmente en los años de 1993, 1994 y 1997 y se realizó una recolecta diaria durante 15 días consecutivos en los años 1995 y 1996. Para ello se arrastró una red de plancton de 505 μ durante 10 min cada hora durante el flujo de marea. La abundancia relativa de las postlarvas de camarón en esta zona varía considerablemente en años cuando el flujo de agua dulce incrementa. La abundancia es mayor hasta en un 200 % ($p < 0.05$) cuando existe descarga de agua dulce al AGC.

REFERENCES

- Alvarez-Borrego, S., B.P. Flores Báez & L. Galindo Bect. 1975. Hidrología del Alto Golfo de California II. Condiciones durante invierno, primavera y verano. Cien. Mar. 2: 21-36.
- Calderón-Pérez, J.A., R.E. Macías & R.S. Rendón. 1989. Clave de identificación para los estadios de postlarva y primeros juveniles de camarón del género *Penaeus* (Crustacea: Decapoda) del Golfo de California. México. Cien. Mar. 15 : 57-70.
- Carvajal, N., A. Souza & R. Durazo. 1997. Anumerical study of the ex ROFI of Colorado River. J. of Mar. Syst. 12: 17-33.
- Dolan, R., A. Howard & A. Galleon. 1974. Man's Impact on the Colorado River in the Grand Canyon. Amer. Sci.62: 390-401.
- Hughes, D.A. 1969. Responses to salinity change as a tidal transport mechanism of pink shrimp, *Penaeus duorarum*. Biol. Bull. 136: 43-53.
- Lavín, M. F. & S. Sánchez. 1999. On how the Colorado River affected the Hydrography of the Upper Gulf of California. Cont. Shelf Res.19: 1545-1560.
- Lavín, M. F., V. M. Godínez & L. Alvarez. 1998. Inverse-Estuarine features of the upper Gulf of California. Estuarine, Coastal and Shelf Sci. 47: 769-795.
- Mair, J. McD. 1979. The identification of postlarvae of four species of *Penaeus* (CRUSTACEA: DECAPODA) from the Pacific Coast of México. J. of Zool. 188: 347-351.
- Mair, J.McD., J.L. Watkins & D.I. Williamson. 1982. Factors affecting the migration of postlarval Penaeid shrimp into a Mexican Lagoon Syst.. Oceanol. Acta, No. S.P. Proc. Int. Symp.on Coastal Systems. SCOR/IABO/UNESCO; Bordeaux, France. 1981. 339-345.
- Mair, J. McD. 1980. Salinity and water type preferences of four species of postlarval (*Penaeus*) from west México. J. Exp. Mar. Biol. Ecol. 45: 69-82.
- Mathews, C.P. 1981. A review of the North American penaeid fisheries, with particular reference to Mexico. Kuwait Bull.of Mar. Sci. 2: 325-409.
- Matthews, T.R., W.W. Schoroeder & D.E. Stearns. 1991. Endogenous rhythm, light and salinity effects on postlarval brown shrimp *Penaeus aztecus* Ives recruitment to estuaries. J. Exp. Mar. Biol. Ecol. 154: 117-189.
- Rothlisberg, P. C. 1998. Aspects of penaeid biology and ecology of relevance to aquaculture: a review. Aquaculture 164: 49-65.
- Turner, R. M. & M.M. Kapiscak. 1980. Recent vegetation changes along the Colorado River between Glen Canyon Dam and Lake Mead Arizona. U. S. Geological Survey Professional paper. Raston, Virginia 125 p.

