

## Antimicrobial activity of *Cymbopogon citratus* (Poaceae) on *Streptococcus mutans* biofilm and its cytotoxic effects

Mailen Ortega-Cuadros<sup>1, 2\*</sup>, Adriana Patricia Tofiño-Rivera<sup>3, 4</sup>, Luciano José Merini<sup>5</sup>,  
María Cecilia Martínez-Pabón<sup>6</sup>

1. PhDe Doctorado en Biotecnología, Universidad de Antioquia. mailen.ortega@udea.edu.co
2. Grupo de investigación Biodiversidad, Corporación para Investigaciones Biológicas-CIB, Cra 72A #78b-141, Medellín, Antioquia, Colombia. mailen.ortegac@gmail.com
3. Corporación Colombiana de Investigación Agropecuaria - AGROSAVIA. Centro de Investigación Motilonia – kilómetro 5 vía a Becerril, Agustín Codazzi, Cesar, Colombia; atofino@agrosavia.co
4. Universidad de Santander, Facultad de ciencias de la salud. Grupo de investigación CIENCIAUDES, Valledupar, Colombia
5. EEA-Anguil, Instituto Nacional de Tecnología Agropecuaria – INTA-CONICET, La Pampa, Argentina. Ruta Nacional 5, Km 580, La Pampa, Argentina; merini.luciano@inta.gov.ar
6. Facultad de Odontología, Universidad de Antioquia, Medellín, Colombia; mcecilia.martinez@udea.edu.co

\* Correspondencia.

Received 28-IV-2018. Corrected 21-VII-2018. Accepted 18-IX-2018.

**Abstract:** Dental caries is a pathology of multifactorial origin and currently natural products are an efficient alternative treatment. The work sought to evaluate the antimicrobial activity of the three substances (*Cymbopogon citratus* essential oil and the citral and myrcene components) against *Streptococcus mutans* ATCC UA159, as well as their cytotoxicity on keratinocytes and human fibroblasts. Through Scanning electron microscopy, the typical morphology of *S. mutans* was observed and, a characteristic formation of polysaccharides, with more structured arrangement at 48 h. The bacterial viability effect against *S. mutans* on biofilms was evaluated through exposure to the three substances by using the MBEC technique-high-throughput at concentrations of 1, 0.1, and 0.01 µg/mL and chlorhexidine as positive control. The cytotoxicity of the compounds was evaluated on keratinocytes and fibroblasts through the MTT reduction technique, using 0.5 mM H<sub>2</sub>O<sub>2</sub> as cell-death control (negative control) and ethanol 1 % as vehicle control (positive control). The three substances evaluated had effects on the bacterial viability of *S. mutans* with mortality between 74 and 96 %, without significant difference among them ( $P > 0.393$ ); additionally, no cytotoxicity was evident on keratinocytes and fibroblasts in a 24 h treatment. The substances evaluated showed significant antimicrobial effects; hence, these should be studied further as potential co-adjuvants to prevent dental caries that cause minor adverse effects.

**Key words:** biofilms; terpenes; dental caries; phytotherapy; *Cymbopogon citratus*; essential oils.

Ortega-Cuadros, M., Tofiño-Rivera, A. P., Merini, L. J., & Martínez-Pabón, M. C. (2018). Antimicrobial activity of *Cymbopogon citratus* (Poaceae) on *Streptococcus mutans* biofilm and its cytotoxic effects. *Revista de Biología Tropical*, 66(4), 1519-1529.

The oral cavity is a dynamic environment that hosts a vast variety of microorganisms, whose lifestyle is characterized by the formation of biofilms or dental plaque (Kolenbrand, 2000). The biofilm develops naturally over the dental surface from extracellular polymers

and microbial consortia (Costerton et al., 1994), which offer protection and resistance for microbial species against environmental variations and antimicrobial agents (Marsh, 2005). Dental plaque prevents oral cavity colonization by exogenous species, thus, maintaining

a beneficial balance with the host. However, when this homeostasis is lost, it alters the metabolism of microbial communities, transforming the biofilm composition to a pathogenic state that permits diseases emergence such as dental caries (Marsh, 2005).

From the Public Health perspective, this disease represents a global problem (OMS, 2012) with incidence between 33.27 and 98 % in the general population of Latin America and Colombia (Bönecker & Cleaton-Jones, 2003; Minsalud, 2014). Treatments to control dental plaque and prevent dental caries include brushing (Ghassemi et al., 2013), dental floss (Lõe, 2000), mouthwash, chlorhexidine, fluoride (Rodrigues, Lussi, Seemann, & Neuhaus, 2011; James et al., 2017), phenols and derivatives of quaternary ammonium (Twetman, 2004). Although there is enough evidence on the preventive effect of several of these substances (Bascones & Morantes, 2006; Kalesinskas, Kačergius, Ambrozaitis, Pečiulienė, & Ericson, 2014), it has been found that they can produce adverse effects when used in concentrations above therapeutic dosages or over a long period of time. These products can induce dental fluorosis (fluorides) (Jha, Mishra, Sharma, & Damodaran, 2011), dysgeusia and dental pigmentation (chlorhexidine) (Varoni, Tarce, Lodi, & Carrassi, 2012), burning sensation, and erosive effects on the enamel (quaternary ammonium compounds and phenols) (Bascones & Morantes, 2006; de Araújo, Gonçalves, Martins, Lima, & Araújo, 2015). In addition, there are groups with special needs regarding oral hygiene, in whom a chemical option for prevention and therapy may have higher utility. Some examples are children with autistic disorder (Lowe & Lindemann, 1985), individuals with Sjögren's syndrome (Mavragani & Moutsopoulos, 2010), and patients with head and neck cancer which, as consequence of treatments associated to its pathology (Pfister et al., 2015), are more susceptible to caries (Jensen et al., 2010). Renewed interest is noted on the use of oral hygiene products formulated from natural compounds, among which

essential oils are highlighted (Lang & Buchbauer, 2012).

As natural products, essential oils have a great chemical diversity and a huge potential in antimicrobial therapies (Lang & Buchbauer, 2012). The species *Cymbopogon citratus* (DC) Stapf is a perennial aromatic plant belonging to the Poaceae (Gramineae) family, cultivated in tropical and sub-tropical regions around the world (Gupta & Jain, 1978). It is commonly known in several countries as “hierba luisa”, “citronella”, “lemongrass”, “lemon tea”, “holy cane” (Negrelle & Gomes, 2007), and “limoncillo” in Colombia (Tofiño-Rivera et al., 2016).

From the bio-prospection perspective, *C. citratus* essential oils and their citral and myrcene compounds have been reported as substances with antimicrobial properties on *S. mutans* planktonic cultures (Bernardes et al., 2010; Vargas et al., 2010) and *S. mutans* biofilms (Tofiño-Rivera et al., 2016; Almeida, Akisue, Cardoso, Junqueira, & Jorge, 2013; Oliveira et al., 2017). However, regarding its adverse effects on oral mucosa, several cell lines present very variable results (Koba et al., 2009; Correa-Royero, Tangarife, Durán, Stashenko, & Mesa-Arango, 2010; Vargas et al., 2010; Kpoviessi et al., 2014; Tofiño-Rivera et al., 2016).

The aim of this study was to assess the antimicrobial activity of *C. citratus* essential oil and its main components, citral and myrcene, on the bacterial viability of *S. mutans* biofilms and their cytotoxicity on HaCaT cells and human fibroblasts.

## MATERIALS AND METHODS

**Plant material:** Plant material used was classified in the “José Cuatrecasas” Herbarium at Universidad Nacional in Palmira with Code 4952 20/10/2014. *C. citratus* essential oil was obtained by the Association of Kankuamos Indigenous (ASOPROKAN), by steam distillation of plant leaves. These leaves were harvested during the rainy season, October-November. Phytochemical analysis for the oil were carried out at the Chemistry of Natural Products

research group, Universidad de Córdoba in Montería, Colombia. Gas chromatography was used with mass selective detector (GC-MS) (Tofiño-Rivera et al., 2016).

**Biofilm cultivation:** To obtain biofilms, *S. mutans* ATCC UA159 was cultured by using the MBEC™ 19131 commercial system (Innovotech, 2015), according to the protocol described by Tofiño-Rivera et al. (2016) with some modifications. First, 4-6 colonies were taken from a Brain Heart agar plate (BHI) (BHI; Merck KGaA, Darmstadt, Germany), transferred to 5 mL of BHI broth and incubated until obtaining an inoculum corresponding to the 1.0 McFarland (8 h ~). One mL was diluted in BHI broth to 1:30; inoculum control was verified through serial dilutions from 10<sup>-1</sup> to 10<sup>-8</sup> (UFC/mL) and with it to be establish the initial count. MBEC™ system was inoculated with the 1:30 dilution and cultured on a benchtop shaking agitator at 110 bpm and 37 °C for 48 h in controlled atmosphere with 5 % CO<sub>2</sub>, replacing the culture medium with fresh medium at 24 h.

**Biofilm growth control:** After incubation, biofilms formed on the pegs surface of the MBEC™ device lid were washed with saline solution at 0.9 % for 2 min to remove any poorly adhered bacterial cells. To verify the biofilm growth, seven pegs were removed from the lid of the MBEC™ device (positions A12 - G12) and processed as follow:

**Viable cell count (CFU/mL):** Three pegs (B12, C12 and E12) were immersed in a microtitre dish containing 200 µL of neutralizing solution and placed in a sonicator (MID-MARK Soniclean model M 150) at 42 kHz for 30 min. Then, serial dilutions were made from 10<sup>-1</sup> to 10<sup>-8</sup> for their subsequent seed in BHI agar plates, through the spot technique and CFU/mL count after 48 h of incubation.

**Crystal violet staining:** Two pegs of the same series (A12 y G12) were dipped in 1 mL of crystal violet 0.5 % and stained during

30 min. Then, pegs were soaked in sterile distilled water and dried at room temperature; thereafter, they were observed in a stereoscope at 6.5 magnification (Zeiss Stemi 2000-C) to verify any biofilm formation.

**Scanning electron microscopy (SEM):** to observe the typical morphology of biofilms and formation of polysaccharides by *S. mutans*, SEM was used. Pegs D12 and F12 were fixed overnight in glutaraldehyde 2.5 % at room temperature for 24 hours. Thereafter, they were washed three times with saline phosphate buffer (SPB) for 5 min each, and then washed with distilled water and dehydrated during 15 min in increasing concentrations of ethanol at 50 %, 75 %, and 95 %, followed by a final dehydration with ethanol at 100 % for 15 min with three repetitions. Finally, the biofilms were dried to a critical point, coated with gold, and analyzed using a scanning electron microscope (SEM) (JEOL JSM 6490 LV).

**Bacterial viability of *S. mutans* biofilms:** Susceptibility tests were conducted on a microtiter plate (challenge plate) (Nunclon Delta Surface; Nunc, Roskilde, Denmark) containing different concentrations of tested substances and their respective controls. For *C. citratus* essential oils and their main components citral (cis+trans 95 % by Alfa Aesar, Karlsruhe, Germany) and myrcene (technical grade, Sigma-Aldrich Chemical, St. Louis, MO, USA), concentrations of 1, 0.1, and 0.01 µg/mL dilution in ethanol at 1 % and chlorhexidine at 0.5 % was used as positive control. For growth controls, ethanol at 1 %, and growth in neutralizing solution were placed in column 1 and the sterility control in column 12. The lid with pegs was dipped in the challenge plate and incubated for 5 min at 37 °C with CO<sub>2</sub> and shaking agitation at 110 bpm. After this time, biofilms were transferred to another microtiter plate (recovery plate), which contained 180 µL of neutralizing solution in each well, (Innovotech, 2015). It was left to rest during 30 min and then brought to sonication for 30 min at 42 kHz to detach biofilms. Serial dilutions

from  $10^0$  to  $10^{-7}$  were prepared with the sonication products, which were seeded in BHI agar plates using the spot seeding technique. After 48 h of incubation, the count of CFU / mL was performed.

To calculate the percentage of dead bacterial cells the following formula was applied:

$$[1 - (\text{Mean CFU treated} / \text{Mean CFU initial bacterial number})] \times 100 \%$$

The reduction logarithm was calculated through the formula:

$$\text{Log}_{10} \text{ Reduction} = \text{Mean Log}_{10} \text{ Untreated Control Peg} - \text{Mean Log}_{10} \text{ Treated Pegs}$$

All tests were performed with seven replicates and three repetitions.

#### **Eukaryotic cells viability assessment through bromide reduction of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazole (MTT):**

Assessment of eukaryotic cells viability was performed according to a method developed by Mosmann (1983) as modified by Denizot and Lang (1986) and indications of the ISO 10993-5-2009 standard (ISO, 2009).

The HaCaT cells and human fibroblasts (skin from healthy patient) were kindly donated by the Tissue Engineering and Cellular Therapies group (GITTC) from University of Antioquia. The HaCaT cells were cultured in DMEM (Dulbecco's Modified Eagle Medium, Sigma-Aldrich Chemical, St Louis, MO, USA) supplemented with 5 % of fetal bovine serum (FBS, Gibco, Grand Island, NY, USA) and 100 µg/mL penicillin and streptomycin (Gibco, Grand Island, NY, USA); and incubated at 37 °C in moist atmosphere and CO<sub>2</sub> at 5 %. Primary cultures of fibroblasts were maintained in identical conditions except for the SBF content that was 10 %.

The initial inoculum was  $1 \times 10^4$  and  $5 \times 10^4$  cells/well for HaCaT cells and fibroblasts respectively. After allowing cell adhesion, treatments were conducted with solutions containing 1, 0.1, and 0.01 µg/mL of *C. citratus* essential oil and its components citral and

myrcene, using H<sub>2</sub>O<sub>2</sub> 0.5 mM as cellular death control and ethanol 1 % as vehicle control. After 24 h exposure, 10 µL of MTT 5 mg/mL were added (Sigma-Aldrich Chemical, St. Louis, MO, USA), followed by 4 h incubation in darkness. Finally, to dissolve the formazan crystals, 100 µL of isopropanol-HCl 0.04 M (Merck KGaA, Darmstadt, Germany) were added and absorbance was measured at 571 nm in an ELISA reader (Multiskan FC - Thermo Scientific). Viability was expressed as percentage respect to death control. All tests were performed by triplicate.


Data was analyzed using SPSS® statistical package version 22. Statistical assumptions were assessed by applying normality and homoscedasticity tests. Intra-group comparison was achieved through one-way ANOVA, Kruskal Wallis tests and multiple comparisons of means against a control (*post-hoc*) via Dunnet's test. Differences were considered statistically significant if  $P < 0.05$ .

## RESULTS

**Phytochemical analysis:** The principal components of *C. citratus* essential oil were citral (58 %) and myrcene (15.6 %). The complete chemical profile of essential oil used was previously published (Toffiño-Rivera et al., 2016).

**Effect on *S. mutans* bacterial viability on biofilms:** Crystal violet staining was used as a visual control of biofilm growth, identifying a cover on the peg as of 24 h. Through SEM, the typical morphology of *S. mutans* was observed and, a characteristic formation of biofilm polysaccharides, with more structured arrangement at 48 h (Fig. 1).

The *C. citratus* essential oil and the citral and myrcene components have the highest mortality percentages at a concentration of 1 µg/mL (93, 96 y 91 % respectively), with a decrease in the inhibitory effect at the lowest concentrations (Table 1). According to the statistical analysis, no statistically significant difference were found among the substances (essential oil, citral and myrcene; Kruskal



**Fig. 1.** SEM of *S. mutans* biofilm on the surface of MBEC pegs after 48 hs of incubation. a) x1,000. Biofilm structure attached to the peg surface; b) x6,000. Typical morphology of *S. mutans*; c) x15,000. The size and typical morphology of *S. mutans* cells; d) x30,000. The presence of the extracellular matrix structuring the biofilm is observed.

**TABLE 1**  
Antimicrobial activity of *C. citratus* essential oil and its main components (citral and myrcene) on the bacterial viability of *S. mutans* biofilms

Substances assessed	Concentration	Viable cell count (Log <sub>10</sub> CFU/ml)	% Mortality	Reduction Log
<i>C. citratus</i> essential oil	1 μg/mL	5.71	93	1.19
	0.1 μg/mL	6.01	87	0.90
	0.01 μg/mL	6.31	74	0.59
Citral	1 μg/mL	5.50	96	1.41
	0.1 μg/mL	5.89	90	1.01
	0.01 μg/mL	5.94	89	0.96
Myrcene	1 μg/mL	5.85	91	1.06
	0.1 μg/mL	5.95	89	0.96
	0.01 μg/mL	5.96	88	0.94
Chlorhexidine	0.5 %	--	100	--
Neutralizing solution control	–	7.44	0	--
Vehicle control (Ethanol)	1 %	7.60	0	--

Wallis tests = 0.393) in addition, these also did not present significant difference with chlorhexidine (Kruskal Wallis tests = 0.06).

**Cellular viability using MTT test:** The MTT test on HaCaT cells and fibroblasts showed that none of the concentrations evaluated for *C. citratus* essential oil or its components, citral and myrcene had toxicity on the cells exposed during 24 h; with highly significant differences in relation to the “death control” H<sub>2</sub>O<sub>2</sub> 0.5 mM (Anova test < 0.05). The treatments do not have significant differences with the positive vehicle control (one-way ANOVA and multiple comparisons of means against a control via Dunnet’s test P > 0.05) (Fig. 2).

## DISCUSSION


Aromatic plants and their derivatives, and especially essential oils, are of biotechnological interest due to their vast chemical diversity and biological properties (Lang & Buchbauer, 2012). The chemical composition of *C. citratus* essential oil from plants cultivated in the Kankuamo indigenous reservation in Atanquez, Cesar Department, Colombia, presents citral (58 %) and myrcene (15.6 %) as main components, classifying this plant within the citral chemotype (Tofiño-Rivera et al., 2016). Similar results were obtained by other groups using a similar method of extraction. They identified citral (46.97 %) and myrcene (27.83 %) or citral (75 %) and β-Pinene (10.1 %) as main components of the *C. citratus* oil (Gbenou et al., 2013; Kpoviessi et al., 2014). Differences can be attributed to the collected leaf maturity state and the distillation time applied (Durán, Monsalve, Martínez, & Stashenko, 2007).

The bacterial viability effect on biofilms, assessed through the MBEC technique, demonstrated that *C. citratus* essential oil, citral and myrcene at all the tested concentrations strongly reduce the viability of *S. mutans* ATCC UA159 on biofilms cultured during 48 h (mortality between 74 % and 96 %) without significant differences against chlorhexidine

(P = 0.06). These finds are interpreted favorably, given that chlorhexidine is one of the most studied and effective anti-plaque treatments (Gunsolley, 2010). Besides, these results are in accordance with previous reports (Tofiño-Rivera et al., 2016) for *C. citratus* essential oils at concentrations 1, 0.1, 0.01 µg/100 mL, which found inhibition on biofilms (mortality between 93.1 and 95.4 %) at 24 h. Almeida et al. (2013) also identified the inhibitory effect of *C. citratus* essential oil at a concentration of 0.5 % on *S. mutans* biofilms ATCC 35688 cultured in acrylic discs within a 24 well microtiter plate for 120 h (Almeida et al., 2013)

The results obtained in this research demonstrate that citral and myrcene also present activity on the bacterial viability of *S. mutans* in biofilms either pure or mixed with other components of essential oil. In previous reports, this is identified and attributed to the antimicrobial effect of phytochemicals, which interact with bacterial cellular structures affecting their function (Saad, Muller, & Lobstein, 2013). Some phytochemicals action mechanisms on microbial cells consist in affecting the integrity of the cellular membranes whether by altering fatty acids composition (Di Pasqua, Hoskins, Betts, & Mauriello, 2006), or surface physicochemical properties (i.e.: hydrophobicity, electric conductivity, filtration of soluble proteins and reducing sugars) (Lopez-Romero, González-Ríos, Borges, & Simões, 2015; Li & Yu, 2015), or modifying the activity of calcium and potassium ion channels (Saad et al., 2013). These can also cause irreversible damage to cytoplasmic membranes, leading to microorganism’s death by unbalance in the ion homeostasis. Conservation of ionic balance is an integral part of maintaining the cell’s energy status, as well as transporting solutes, regulating metabolism, controlling turgor pressure and motility (Li & Yu, 2015). Membrane level damage is a mechanism identified by Saad et al. (2013) for *C. citratus* essential oil on *Escherichia coli*.

Oil safety was verified, in terms of cytotoxic effect on eukaryotic cells. Tissues fulfill biological functions that maintain mucous



**Fig. 2.** Cytotoxicity on HaCaT and fibroblasts cells of the essential oil of *C. citratus* and its components citral and myrcene. Cytotoxic effect of the three concentrations evaluated from each treatment in relation to the cell death control (H<sub>2</sub>O<sub>2</sub>). a) *C. citratus* essential oil on HaCaT cells. b) Citral component on HaCaT cells. c) Myrcene component on HaCaT cells. d) *C. citratus* essential oil on fibroblasts. e) Citral component on fibroblasts. f) Myrcene component on fibroblasts. Data expressed as mean ± DS, n = 3. \*\*\* = P < 0. 05.

membranes integrity and they should not be affected by exogenous substances (Newman, Takei, Klokkevold, & Carranza, 2006). The results show that the *C. citratus* essential oil and its components citral and myrcene, do not evidence cytotoxicity on the HaCaT cells or on human fibroblasts (Fig. 2) in a 24 h treatment. Similar results were reported by other authors, using HaCaT cells and concentrations of the essential oil between 25 and 100  $\mu\text{L}/\text{mL}$  (Koba et al., 2009). Neither Correa-Royero et al. (2010) exposing Vero cells to concentrations < 200  $\mu\text{g}/\text{mL}$  and Tofiño-Rivera et al. (2016) exposing CHO cells at concentrations of 1, 0.1, and 0.01  $\mu\text{g}/100 \text{ mL}$  identified essential oil cytotoxicity. Vargas et al. (2010) evidenced slight cytotoxicity on odontoblastoid cells at a concentration of 1 000  $\mu\text{g}/\text{mL}$  during 120 min, however, this may be attributed to the use of higher concentration, than prior studies (Vargas et al., 2010). Kpoviessi et al. (2014), identified toxicity on CHO cells and non-carcinogenic human fibroblasts with  $\text{IC}_{50} = 10.63$  and  $39.77 \mu\text{g}/\text{mL}$ , respectively (Kpoviessi et al., 2014), possibly due to the different essential oil chemical composition (citral and  $\beta$ -Pinene), usually obtained when plants are cultured at high temperatures. Plant material from which essential oil was extracted was collected during the rainy season, evidencing that the change on phytochemistry, associated to the effect of environmental factors and physiological status of the biomass collected, can influence upon its cytotoxic effect (Durán et al., 2007; Gbenou et al., 2013).

It must be considered that the cytotoxicity results reported in this research were obtained by exposing cells to 24 h treatments in order to identify residual effects of the substances evaluated, related to sub-lethal or lethal effects on the tissues that prolonged exposure may cause. Recommended time for mouthwash use is about 2 min (Duckworth, 2013), so the effectiveness on the control of *S. mutans* biofilms and harmlessness of the *C. citratus* essential oil and its main components with eukaryotic cells highlight the safety of the substances and backs the formulation of natural product.

These formulations would contribute to the selective control of microbial populations from dental plaque and pursue the maintenance of normal oral microbiota that performs beneficial functions for the host (Marsh, Head, & Devine, 2015).

We foresee the use of experimental models that permit a greater approach to the possible *in vivo* therapeutic effect of formulations designed for the conditions of the oral cavity. *C. citratus* is included in the list of medicinal plants recognized for therapeutic purposes and medicinal use, hence its formulation could be openly sold (Invima, 2016), according to the colombian law (MPS, 2004). Access to *C. citratus* and its benefits has been available for the Colombian population base, associated to its analgesic and anti-inflammatory effect in traditional medicine. For that reason, it has important ethnobotanic tradition in the different indigenous communities (Trujillo & Correa-Múnera, 2010; Ortega, 2015). In addition, regarding to its biocide effect, steam stripped essential oils derived from this plant have greater stability than its extracts (Bandoni, Retta, Di Leo Lira, & Baren, 2009). Both advantages can be used as alternative to boost the economy agro-industrial of the local communities where biomass is produced. It is important to remark the background of the Kankuamo Indigenous Association of Agroecological Producers (ASOPROKAN), located in Atanquez, Colombia in organic production of *C. citratus* essential oil. (Ulloa, 2011; Vélez, Rátiva, & Varela, 2012). Considering the obtained results and the biotechnologically promising performance showed by *C. citratus* essential oils, we recommend to move toward the formulation of products for oral health based on these bio-inputs, by prioritizing technical support to the communities that produce these bio-products and ensuring its availability for low-income communities.

#### ACKNOWLEDGMENTS

The authors thank Sandra Milena Morales Uchima (microbiologist) and the work team of the Oral Histopathology and Microbiology



Laboratory of the Faculty of Dentistry, Universidad de Antioquia, for her support during the experimental process. Gratitude is also expressed to Jorge Humberto Botero Garcés, professor in the Department of Microbiology and Parasitology, Faculty of Medicine, University of Antioquia, for his support with the statistical analysis of the data; to the mentorship committee, the Kankuamo association - ASOPROKAN, to Mario Zapata, director of Corpoica C.I Motilonia, Cesar. This project was funded by COLCIENCIAS project: "Restoration of soils degraded by mining by using rhizoremediation strategies based on the use of native aromatic species that promote development of regional micro-economies", Contract 1852 COLCIENCIAS-CORPOICA call 563 2014, chapter Argentina; CORPOICA project internal agenda: "Aromatic Medicinal and Condiment Plants; a Productive Alternative for Colombia" MADR-CORPOICA agreement TV15 resolution 000023 of 2015, FONTAGRO project 16598: "Development of regional micro-economies in the production of essential oils harvested in mining soils", 2016 announcement and the Faculty of Dentistry of the University of Antioquia. This work was reviewed and approved by the ethics committee of the Faculty of Dentistry of the University of Antioquia, as stated in Concept No 02-2017 of Act No. 01 of 2017.

## RESUMEN

**Actividad antimicrobiana de *Cymbopogon citratus* (Poaceae) sobre biopelícula de *Streptococcus mutans* y sus efectos citotóxicos.** La caries dental, es una patología de origen multifactorial y actualmente los productos naturales son un tratamiento alternativo eficiente. El trabajo buscó evaluar la actividad antimicrobiana del aceite esencial de tres sustancias (*Cymbopogon citratus* y los componentes citral y mircenol) contra *Streptococcus mutans* ATCC UA159, así como su citotoxicidad sobre queratinocitos y fibroblastos humanos. A través de Microscopía Electrónica de Barrido, se observó la morfología típica de *S. mutans*, y una formación característica de polisacáridos, con una disposición más estructurada a las 48 h. El efecto de viabilidad bacteriana contra *S. mutans* en biofilms se evaluó a través de la exposición a las tres sustancias mediante el uso de la técnica de alto rendimiento MBEC, a concentraciones de 1, 0.1 y 0.01 µg/ml y clorhexidina

como control positivo. La citotoxicidad de los compuestos se evaluó sobre queratinocitos y fibroblastos, mediante la técnica de reducción de MTT, utilizando 0.5 mM de H<sub>2</sub>O<sub>2</sub> como control de muerte celular (control negativo) y etanol 1 % como control del vehículo (control positivo). Las tres sustancias evaluadas, tuvieron efectos sobre la viabilidad bacteriana de *S. mutans* con una mortalidad entre 74 y 96 %, sin diferencia significativa entre ellos (P > 0.393); adicionalmente, no fue evidente la citotoxicidad sobre queratinocitos y fibroblastos en un tratamiento de 24 h. Las sustancias evaluadas mostraron efectos antimicrobianos significativos; por lo tanto, estos deben estudiarse más a fondo como posibles coadyuvantes para prevenir la caries dental que causa efectos adversos menores.

**Palabras clave:** biofilms; terpenos; caries dental; fitoterapia; *Cymbopogon citratus*; aceites essential.

## REFERENCES

- Almeida, R. B., Akisue, G., Cardoso, L. M. L., Junqueira, J. C., & Jorge, A. O. C. (2013). Antimicrobial activity of the essential oil of *Cymbopogon citratus* (DC) Stapf. on *Staphylococcus* spp., *Streptococcus mutans* and *Candida* spp. *Revista Brasileira de Plantas Mediciniais*, 15, 474-482.
- Bandoni, A. L., Retta, D., Di Leo Lira, P. M., & Baren, C. M. (2009). ¿Son realmente útiles los aceites esenciales? *Boletín Latinoamericano y del Caribe de Plantas Medicinales y Aromaticas*, 8, 317-322.
- Bascones, A. & Morantes, S. (2006). Antisépticos orales. Revisión de la literatura y perspectiva actual. *Avances en Periodoncia*, 18, 31-59.
- Bernardes, W. A., Lucarini, R., Tozatti, M. G., Flauzino, L. G., Souza, M. G., Turatti, I. C., ... Cunha, W. R. (2010). Antibacterial activity of the essential oil from *Rosmarinus officinalis* and its major components against oral pathogens. *Zeitschrift für Naturforschung*, C, 65, 588-93.
- Bönecker, M., & Cleaton-Jones, P. (2003). Trends in dental caries in Latin American and Caribbean 5-6- and 11-13-year-old children: a systematic review. *Community Dentistry and Oral Epidemiology*, 31, 152-157.
- Correa-Royero, J., Tangarife, V., Durán, C., Stashenko, E., & Mesa-Arango, A. (2010). In vitro antifungal activity and cytotoxic effect of essential oils and extracts of medicinal and aromatic plants against *Candida krusei* and *Aspergillus fumigatus*. *Revista Brasileira de Farmacognosia*, 20, 734-741.
- Costerton, J. W., Lewandowski, Z., DeBeer, D., Caldwell, D., Korber, D., & James, G. (1994). Biofilms, the customized microniche. *Journal of Bacteriology*, 176, 2137-2142.

- de Araújo, D. B., Gonçalves, B. E. M., Martins, G. B., Lima, M. J. P., & Araújo, M. T. B. (2015). Saúde bucal: a importância dos enxaguatórios com antisépticos. *Revista de Ciências Médicas e Biológicas*, *14*, 88-93.
- Denizot, F., & Lang, R. (1986). Rapid colorimetric assay for cell growth and survival. *Journal of Immunological Methods*, *89*, 271-277.
- Di Pasqua, R., Hoskins, N., Betts, G., & Mauriello, G. (2006). Changes in membrane fatty acids composition of microbial cells induced by addiction of thymol, carvacrol, limonene, cinnamaldehyde, and eugenol in the growing media. *Journal of Agricultural and Food Chemistry*, *54*, 2745-2749.
- Durán, D. C., Monsalve, L. A., Martínez, J. R., & Stashenko, E. E. (2007). Estudio comparativo de la composición química de aceites esenciales de *Lippia alba* provenientes de diferentes regiones de Colombia, y efecto del tiempo de destilación sobre la composición del aceite. *Scientia Et Technica 2007*, *XIII* (33), 435-438.
- Duckworth, R. M. (2013). Pharmacokinetics in the Oral Cavity: Fluoride and Other Active Ingredients. *Monographs in Oral Science*, *23*, 125-139.
- Gbenou, J. D., Ahounou, J. F., Akakpo, H. B., Laleye, A., Yayi, E., Gbaguidi, F., ... Kotchoni, S. O. (2013). Phytochemical composition of *Cymbopogon citratus* and *Eucalyptus citriodora* essential oils and their anti-inflammatory and analgesic properties on Wistar rats. *Molecular Biology Reports*, *40*, 1127-1134.
- Ghassemi, A., Vorwerk, L., Hooper, W., Patel, V., Sharma, N., & Qaqish, J. (2013). Comparative plaque removal efficacy of a new children's powered toothbrush and a manual toothbrush. *The Journal of Clinical Dentistry*, *24*, 1-4.
- Gunsolley, J. C. (2010). Clinical efficacy of antimicrobial mouthrinses. *Journal of Dentistry*, *38*, S6-10.
- Gupta, B., & Jain, N. (1978). Cultivation and utilization of Genus *Cymbopogon* in Indian. *Indian Perfumer*, *22*, 55-68.
- Innovotech. (2015). MBEC™ Assay For High-Throughput Antimicrobial Susceptibility Testing of Biofilms. Recuperado de <http://bit.ly/2ykWf6Q>
- International Organization for Standardization. (2009). ISO 10993-5 (3<sup>d</sup> ed.). 2009-06-01. Recuperado de [www.iso.org](http://www.iso.org)
- Invima. (2016). Listado de Plantas Medicinales Aceptadas con Fines Terapéuticos 21-12-2016. Recuperado de <http://bit.ly/2f6N0Xi>
- James, P., Worthington, H. V., Parnell, C., Harding, M., Lamont, T., Cheung, A., ... Riley, P. (2017). Chlorhexidine mouthrinse as an adjunctive treatment for gingival health (Review). *Cochrane Database of Systematic Reviews*, *3*, CD008676.
- Jensen, S. B., Pedersen, A. M. L., Vissink, A., Andersen, E., Brown, C. G., Davies, A. N., ... Brennan, M. T. (2010). A systematic review of salivary gland hypofunction and xerostomia induced by cancer therapies: Prevalence, severity and impact on quality of life. *Support Care Cancer*, *18*, 1039-1060.
- Jha, S. K., Mishra, V. K., Sharma, D. K., & Damodaran, T. (2011). Fluoride in the environment and its metabolism in humans. *Reviews of Environmental Contamination and Toxicology*, *211*, 121-142.
- Kalesinskas, P., Kačergius, T., Ambrozaitis, A., Pečiulienė, V., & Ericson, D. (2014). Reducing dental plaque formation and caries development. A review of current methods and implications for novel pharmaceuticals. *Stomatologija, Baltic Dental and Maxillofacial Journal*, *16*, 44-52.
- Koba, K., Sanda, K., Guyon, C., Raynaud, C., Chaumont, J. P., & Nicod, L. (2009). In vitro cytotoxic activity of *Cymbopogon citratus* L. and *Cymbopogon nardus* L. essential oils from Togo. *Bangladesh Journal of Pharmacology*, *4*, 29-34.
- Kolenbrander, P. E. (2000). Oral Microbial Communities: Biofilms, Interactions, and Genetic Systems. *Annual Review of Microbiology*, *54*, 413-437.
- Kpoviessi, S., Bero, J., Agbani, P., Gbaguidi, F., Kpadonou-Kpoviessi, B., Sinsin, B., ... Quetin-Leclercq, J. (2014). Chemical composition, cytotoxicity and in vitro antitypanosomal and antiplasmodial activity of the essential oils of four *Cymbopogon* species from Benin. *Journal of Ethnopharmacology*, *151*, 652-659.
- Lang, G., & Buchbauer, G. (2012). A review on recent research results (2008-2010) on essential oils as antimicrobials and antifungals. A review. *Flavour and Fragrance Journal*, *27*, 13-39.
- Li, C. M., & Yu, J. P. (2015). Chemical Composition, Antimicrobial Activity and Mechanism of Action of Essential Oil from the Leaves of *Macleaya Cordata* (Willd.) R. Br. *Journal of Food Safety*, *35*, 227-236.
- Löe, H. (2000). Oral hygiene in the prevention of caries and periodontal disease. *International Dental Journal*, *50*, 129-139.
- Lopez-Romero, J. C., González-Ríos, H., Borges, A., & Simões, M. (2015). Antibacterial Effects and Mode of Action of Selected Essential Oils Components against *Escherichia coli* and *Staphylococcus aureus*. *Evidence-Based Complementary and Alternative Medicine*, *2015*, 1-9.
- Lowe, O., & Lindemann, R. (1985). Assessment of the autistic patient's dental needs and ability to undergo dental examination. *Journal of Dentistry for Children*, *52*, 29-35.

- Marsh, P. D. (2005). Dental plaque: Biological significance of a biofilm and community life-style. *Journal of Clinical Periodontology*, 32, (suppl. 6), 7-15.
- Marsh, P. D., Head, D. A., & Devine, D. A. (2015). Ecological approaches to oral biofilms: Control without killing. *Caries Research*, 49, 46-54.
- Mavragani, C. P., & Moutsopoulos, H. M. (2010). The geo-epidemiology of Sjögren's syndrome. *Autoimmunity Reviews*, 9, A305-A310.
- Ministerio de la protección social (MPS). (2004). Decreto número 2266 de 2004. Recuperado de <http://bit.ly/2wCvFMj>
- Ministerio Nacional de Salud y Protección Social (Minsalud). (2014). IV estudio nacional de salud bucal (ENSAB). Bogota. Recuperado de <http://bit.ly/2wrBxgf>
- Mosmann, T. (1983). Rapid colorimetric assay for cellular growth and survival: Application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, 65, 55-63.
- Negrelle, R. R. B., & Gomes, E. C. (2007). *Cymbopogon citratus* (DC.) Stapf: chemical composition and biological activities. *Revista Brasileira de Plantas Mediciniais*, 9, 80-92.
- Newman, M., Takei, H., Klokkevold, P., & Carranza, F. (2006). *Periodontología Clínica* (10ª ed). México: McGraw-Hill.
- Oliveira, M. A. C., Borges, A. C., Brighenti, F. L., Salvador, M. J., Gontijo, A. V. L., & Koga-Ito, C. Y. (2017). *Cymbopogon citratus* essential oil: effect on polymicrobial caries-related biofilm with low cytotoxicity. *Brazilian Oral Research*, 31, 1-12.
- Organización mundial de la salud (OMS). (2012). Salud bucodental: Nota informativa N° 318. Recuperado de <http://bit.ly/2y9XSLD>
- Ortega, E. (2015). Traditional uses of plants in the Colombian Orinoquia. *Revista Universitaria en Ciencias de la Salud*, 21, 16-28.
- Pfister, D. G., Spencer, S., Brizel, D. M., Burtness, B., Busse, P. M., Caudell, J. J., ... Hughes, M. (2015). Head and Neck Cancers, Version 1.2015. *Journal of the National Comprehensive Cancer Network*, 13, 847-55.
- Rodrigues, J. A., Lussi, A., Seemann, R., & Neuhaus, K. W. (2011). Prevention of crown and root caries in adults. *Periodontology 2000*, 55, 231-249.
- Saad, N. Y., Muller, C. D., & Lobstein, A. (2013). Major bioactivities and mechanism of action of essential oils and their components. *Flavour and Fragrance Journal*, 28, 269-279.
- Toffiño-Rivera, A., Ortega-Cuadros, M., Galvis-Pareja, D., Jimenez-Rios, H., Merini, L. J., & Martínez-Pabón, M. C. (2016). Effect of *Lippia alba* and *Cymbopogon citratus* essential oils on biofilms of *Streptococcus mutans* and cytotoxicity in CHO cells. *Journal of Ethnopharmacology*, 194, 749-754.
- Trujillo, W., & Correa-Múnera, M. (2010). Plants used by a Coreguaje indigenous community in the Colombian Amazon. *Caldasia*, 32, 1-20.
- Twetman, S. (2004). Antimicrobials in future caries control? A review with special reference to chlorhexidine treatment. *Caries Research*, 38, 223-229.
- Ulloa, A. (2011). The Politics of Autonomy of Indigenous Peoples of the Sierra Nevada de Santa Marta, Colombia: A Process of Relational Indigenous Autonomy. *Latin American and Caribbean Ethnic Studies*, 6, 79-107.
- Vargas, F. S., Oliveira, C. F., Giro, E. M. A., Sacramento, L. V. S., Spolidorio, D. M. P., & Costa, C. A. S. (2010). Efeito antimicrobiano e citotóxico do óleo essencial de *Cymbopogon citratus* sobre células odontoblastóides. *Revista Odontológica do Brasil Central*, 19, 101-107.
- Varoni, E., Tarce, M., Lodi, G., & Carrassi, A. (2012). Chlorhexidine (CHX) in dentistry: state of the art. *Minerva Stomatologica*, 61, 399-419.
- Vélez, I., Rátiva, S., & Varela, D. (2012). Cartografía social como metodología participativa y colaborativa de investigación en el territorio afrodescendiente de la cuenca alta del río Cauca. *Cuadernos de Geografía*, 21, 59-73.