



<https://doi.org/10.15517/rev.biol.trop.v70i1.46904>

Altitudinal distribution of the functional feeding groups of aquatic macroinvertebrates using an ecological network in Andean streams

Juan Pablo Serna^{1*}; <https://orcid.org/0000-0002-2805-8218>

David Fernandez¹; <https://orcid.org/0000-0003-4898-4144>

Fabio Velez¹; <https://orcid.org/0000-0001-6348-6405>

Julián Ruiz¹; <http://orcid.org/0000-0001-5863-9761>

Broder Breckling²

Néstor Aguirre¹; <https://orcid.org/0000-0002-0847-7335>

1. Facultad de Ingeniería, Universidad de Antioquia, Medellín, Colombia; jpablo.serna@udea.edu.co (Correspondence*), david.fernandez@udea.edu.co, fabio.velez@udea.edu.co, julian.ruiz@udea.edu.co, nestor.aguirre@udea.edu.co
2. University of Bremen, General and Theoretical Ecology, broder@uni-bremen.de

Received 07-V-2021. Corrected 18-XI-2021. Accepted 31-I-2022.

ABSTRACT

Introduction: Analysis of functional feeding groups (FFG) in aquatic macroinvertebrates is important in understanding the structure, function, and dynamics of ecological processes in ecosystems. Modularity refers to the degree of compartmentalization of food webs and varies between -1 and 1. A network with a modularity value close to 1 is resilient to disturbances and can be interpreted as an indicating factor for the stability of communities.

Objective: In this study, we analyzed the trophic structure of benthic macroinvertebrates in La Nitrera stream, the San Juan River, and the Cauca River in the Colombian Andes.

Methods: The study was supported by ecological networking techniques using *Gephi* software. We studied nine sites in dry, rainy, and transition seasons in 2017 and 2018, monitoring changes in the altitude gradient. At each of the sites, the organisms were captured and determined, and physicochemical and hydraulic information was obtained.

Results: The variance component analysis allowed to explain the variability of the data by relating the following environmental variables: FFG, diversity, richness, modularity, season, and time. Simple multifactorial ANOVA indicated that significant changes in FFG were associated with altitude, and modularity to time. The allocation of the FFG was done by stomach analysis and secondary information.

Conclusion: The transition season had the highest modularity, possibly due to the recolonization of some biotopes caused by the decrease in the velocity of water currents. La Nitrera and San Juan presented higher values than the Cauca, which may indicate that the altitudinal change and velocity of water currents affects the compartmentalization of the network.

Key words: modularity; altitudinal gradient; stomach analysis; network analysis; invertebrate trophic structure.

Network analysis is an important element in understanding the structure, function, and dynamics of ecological systems. These systems are complex because they involve relationships

and interactions that are not easily quantifiable. However, progress has been made in understanding some interactions whose identity may change in space and time, such as

food webs, patch dynamics, and population fluctuations (Anand et al., 2010). Although networks have provided fundamental representations of ecological complexity, they need to be extended to systematically and simultaneously capture these multifaceted interactions (Pilosof et al., 2017).

The importance of aquatic macroinvertebrates as indicators of environmental conditions lies in the benthic life form of these organisms, their longer life cycles compared other aquatic organisms like bacteria and algae, and their specialized adaptation to specific environmental characteristics (Castillo-Figueroa et al., 2018; Roldán-Pérez, 2016). These aspects become a response to the assessment of the space-time conditions of the environment in which they are found (National Academy of Sciences, 1988; Rosenberg & Resh, 1993). The energy and nutrient transfer in aquatic food webs depends mainly on the diversity, quality, and quantity of resources available in the system, as well as on the species composition, the number of individuals and their trophic relationships (Cummins et al., 2005).

Food webs are important in the ecological processes that occur in river ecosystems. As such, information on food web relations serve as a relevant part of the scientific basis for the planning and management of river ecosystems (Tamaris-Turizo et al., 2018). In the case of aquatic macroinvertebrate communities, these trophic relationships relate to life cycles, habitat choice, behavior, predation, and other factors (Chará-Serna et al., 2010).

The complexity involved in analyzing these interactions, especially the species classification and taxonomy in tropical rivers in Colombia, has prompted studies in fields such as functional ecology (Schmera et al., 2013; Schmera et al. 2017), whereby functional traits are assigned that allow the measurement of morphological, physiological or phenological characteristics at the individual level (Violle et al., 2007). In this respect, some authors have made advances in the analysis, study and functional classification of the aquatic insects feeding groups in the Neotropics (Chará-Serna

et al., 2010; Ramírez & Gutiérrez-Fonseca, 2014; Tamaris-Turizo et al., 2018; Tomanova et al., 2006). This has improved the understanding of the predator-prey relationship function among organisms, and thereby the determination of the balance of the communities that inhabit a biotope.

In Colombia, the altitudinal variation influences the physico-biotic structure of the rivers because at short distances in the Colombian Andean it is possible to have changes in the biotopes and species composition. These changes are also accompanied by rheological variability due to the annual cycles of rainfall and dry seasons (Latrubesse et al., 2005). For example, in the rainy season the rivers are abundant and turbid, while in the dry season the rivers significantly reduce their flow and increase their transparency of water (Gutiérrez, 2018). Thus, spatial variability – meters elevation - and variability in time, i.e., annual rainfall cycle, influence the structure and functioning of aquatic macroinvertebrate.

According to Ramírez and Gutiérrez-Fonseca (2014), aquatic macroinvertebrates can be classified into six major categories. Scrapers (Sc) generally live attached to rocks and other substrates and feed on periphyton (algae, bacteria, and fungi). Piercers (Pc) feed mainly on vascular plants by cutting the tissue and consuming its fluid. Shredders (Sh) chew up plant and wood debris, breaking down large particles into smaller ones that can be assimilated by other organisms, and generally facilitating the organic matter decomposition process. Collectors (GC) can screen small particles, although, having smaller oral devices, they also feed on material that can be re-suspended in the water column. Filter feeders (Ft) remove particles from the water column; some may consume bacteria and suspended organic particles by filtering. Finally, predators (Pr) are those that can consume other organisms, some of which have mouth structures like jaws.

In ecology, the idea of modularity is widely used (Olesen et al., 2007; Ruiz-Toro et al., 2020). One application of it is to measure the degree of compartmentalization of food



webs, which can be used to indicate the stability relations of communities. A network with high modularity would be considered as more resilient to external disturbances or changes because influences to a limited number of the parts would less intensely propagate throughout the network (Gauzens et al., 2015; Stouffer & Bascompte, 2011). The idea that a modular organization would be beneficial for the local stability of ecological communities and their ability to recover from small perturbations (Grilli et al., 2016).

The question that this paper seeks to answer is whether the altitudinal gradient and temporal variation influence the distribution and resistance of aquatic macroinvertebrate communities in a river basin. Therefore, this research aims to analyze the distribution of aquatic macroinvertebrates according to the altitudinal gradient in two rivers and a stream in the same basin in Antioquia, Colombia. This is done using ecological networks analysis and modularity as a response variable, to determine whether there are differences in the resistance characteristics of macroinvertebrate communities according to the typology of each river or stream.

MATERIALS AND METHODS

Study area and site characteristics: The study area is located in the department of Antioquia in Colombia (Fig. 1) and includes three different catchment areas: a micro-basin where the La Nitrera stream is located; an intermediate basin whose main drainage is the San Juan, between the municipality of Jardín and its mouth on the Cauca River in the place known as Peñalisa; and a section of the Cauca River forming a macro-basin, between the village of Bolombolo and the municipality of Caucasia in the North of the department.

Three sampling stations were located at different altitudinal gradients at each study site. At each of the sites, the physicochemical, hydraulic, and biological variables were measured. La Nitrera stream is in the Southwest of the department of Antioquia, in the

municipality of Concordia. Its source is 2 190 elevation meters, and it is one of the main tributaries of the reservoir located in La Nitrera protected area that supplies water to the municipality of Concordia. This natural park is classified as low mountain rainforest with a highly rugged topography (Morales-Quintero et al., 2019; Ruiz-Toro et al., 2020).

The San Juan River basin is in the Southwest region of the Department of Antioquia, with an area of approximately 1 400 km², and elevations between 534 and 3 920 meters elevation San Juan River, within the classification of types of streams, can be classified as a river or mountain stream. The Cauca River originates in the Colombian Massif between the Western and Central Andes Mountains and has a drainage area of 59 000 km², which represents 5 % of the national territory (Puertas et al., 2011).

The fluvial system of the Cauca River runs 1 350 km from its source in the Colombian massif (Cauca Department) to its mouth in the “Brazo de Loba” (Bolívar Department). The basin is polluted by mining, agro-industrial waste and wastewater treatment plants and hydroelectric activities, resulting in large amounts of sediments with heavy metals, xenobiotics, and cyanides along its path (López, 2013; Torres & Pinilla, 2011). Table 1 presents the geographical information on latitude, longitude, and altitude.

Field sampling collection: The information for site description was gathered at each of the nine stations (Table 1). Hydraulic and physicochemical data, as well as information on aquatic macroinvertebrates, were sampled. Each of these variables was repeatedly recorded. The fieldwork was carried out over dry (February, July), rainy (April, September) and transition (August, September) seasons.

Environmental and biological variables: The parameters measured in the field were: dissolved oxygen, electrical conductivity, pH, turbidity, and water temperature. WTW 3330 multiparametric cells were used for this purpose. The aquatic macroinvertebrates were




Fig. 1. Sampling site locations with the three streams: La Nitrera stream (NS1, NS2, NS3), San Juan River (SJS1, SJS2, SJS3) and Cauca River (CS1, CS2, CS3).

TABLE 1
Geographic information

Site	Longitude (North)	Latitude (West)	Altitude (m)
NS1	6°2.129'	75°56.026'	2 220
NS2	6°2.139'	75°55.669'	2 183
NS3	6°2.119'	75°55.951'	2 096
SJS1	5°35' 14"	75°48' 44"	2 134
SJS2	5°35' 58"	75°49' 03"	1 892
SJS3	5°55' 56"	75°51' 34"	556
CS1	5°58' 31"	75°50' 14"	555
CS2	6°2.13'	75°56.026'	453
CS3	6°2.14'	75°55.669'	50

collected with a Surber net of 0.09 m² area placed against the current. Large material such as stones was extracted and the material in the quadrant was removed by hand, ensuring that the sediment and organisms remained in

the net. This process was carried out for one minute in one of the margins of each stream. Also, a semiquantitative sampling using a triangular net was carried out in a 10 m section for five minutes and the sampling effort



in each station was the same. The extracted material was separated in situ in plastic trays and later stored in plastic bottles with 70 % alcohol duly labeled with date, place, type of substrate, replica, responsible, to later be taken to the Laboratory of Sanitary Hydrobiology of the University of Antioquia.

All these measurements were taken in duplicate. Different substrates were chosen according to the station: leaf and rock for La Nitrera and rock for San Juan and Cauca. The extracted material was separated in situ in plastic trays and later stored in plastic bottles with 70 % alcohol, labeled with date, place, type of substrate, replicate, and person responsible. These were then transported to the sanitary hydrobiology laboratory of the University of Antioquia. In the laboratory, all samples of the organisms were determined to the lowest possible taxonomic level using taxonomic keys (Álvarez & Daza, 2005; Boltovskoy et al., 1995; Cummins et al., 2005; Domínguez & Fernández, 2009; Govedich & Moser, 2015; Lasso et al., 2018; Petersen, 2002; Ramírez & Gutiérrez-Fonseca, 2014; Roldán, 1988, 2003; Thorp & Covich, 2009). A BST-606 stereomicroscope was used for this purpose.

Stomach analysis of functional feeding groups: The protocol proposed by Muñoz et al. (2009) in the trophic relationships chapter for the river ecosystem was used to analyze the stomach contents of aquatic macroinvertebrates (Elosegui & Sabater, 2009). The digestive analysis was performed for the most representative morphotypes (Table 2). From these three specimens were selected. This was to identify

the food preference of each organism reported in the literature.

For this analysis, the digestive material closest to the mouth was extracted with the help of dissection forceps and deposited on a slide before a drop of glycerol (85 %) was added. The sample was then homogenized with the assistance of a coverslip, and the observation was made under the inverted microscope, making an inventory of the material present in each plate (Muñoz et al., 2009). The observations were made with an inverted *Boeco* BIB100 microscope using the 40X (0.05 mm) objective, except for the *Phylloicus*, where the 10X objective was used.

Biomass determination: The biomass was determined in the sanitary hydrobiology laboratory by relative dry weight following the protocol of Puerta et al. (2009) his purpose, semi-quantitative sampling of La Nitrera stream was performed, integrating the biotopes (rock and leaf litter), which were then arranged in porcelain crucibles. Initially, the sample was filtered for 30 minutes, removing as much alcohol as possible. The organisms were then separated according to their functional feeding group and placed in an MF-2001 electric muffle furnace at 105 °C for two hours. The fixed solids determination (inorganic matter) was then performed, first calcining the sample at 550 °C for half an hour, then weighing the sample on a Shimadzu TX323L analytical balance with a sensitivity of 0.001 mg. This procedure was performed by replication. In this way, the volatile solids were determined via the

TABLE 2
Diversity indices in the three streams

Index	Nitrera			San Juan			Cauca		
	Mean (\bar{x})	Standard deviation (σ)	Variance coefficient (VC)	Mean (\bar{x})	Standard deviation (σ)	Variance coefficient (VC)	Mean (\bar{x})	Standard deviation (σ)	Variance coefficient (VC)
Richness	9.541	2.604	27.29	6.65	2.158	32.46	4.5	3.326	73.92
Diversity	1.812	0.283	5.66	1.329	0.387	29.18	0.979	0.738	75.41
Evenness	0.820	0.080	9.816	0.723	0.122	16.87	0.773	0.211	27.34
Dominance	0.198	0.080	40.47	0.354	0.164	46.25	0.407	0.348	85.41

difference in weight, indicating the available biomass that can be used as energy for higher-order organisms (Ruiz-Toro et al., 2020).

Modularity Analysis using ecological network: The modularity value was obtained through the software *Gephi* 0.9.2. This metric indicates the feeding relationships of the aquatic macroinvertebrates present in three rivers under altitudinal gradient distributions. This modularity value allows the determination of the clustering intensity degree of a network. This measures the quality of the resulting partitions and can have a scalar value between -1 and 1, with positive values indicating the possible presence of community structure. If the value is negative, it indicates a bad grouping (Blondel et al., 2008).

Gephi 0.9.2 software was used as an easily accessible tool for the representation and analysis of ecological networks (Bastian et al., 2009; Martin, 2015). This software was applied to the data of the functional feeding groups of the semi-quantitative sampling. The networks were made up of two components: the list of nodes that made up the network (functional feeding groups and morphospecies abundances) and a list of interactions between the nodes (biomass).

Analysis and processing of data: Descriptive analyses were performed using inferential statistics relating to the physico-chemical, hydro morphological, and biological variables for each environment, considering averages, maxima, and minima, to visualize trends and extremes in each of the variables. The modularity associated with trophic interactions was used as a response variable. In this way, through multifactorial ANOVA of two factors, it was possible to determine whether there were significant differences associated with time or site of sampling. Finally, the diversity indices were determined through the *Biodiversity Pro* software.

RESULTS

Abundance and ecological structure of aquatic macroinvertebrates: A total of 7 702 benthic macroinvertebrates were collected and identified, comprising 150 taxa. *Smicridea*. (N = 1 096) was the most abundant taxon in all streams, followed by *Thraulodes* sp. (N = 971) and *Anacroneuria*. (N = 576). In La Nitrrera, the most abundant taxa were *Smicridea* and *Phylloicus* (N = 935 and 431). In the San Juan River, *Thraulodes* (N = 546) *Traverella* (N = 325) had the highest abundances in the Cauca River. Table 2 presents in terms of richness, La Nitrrera had the highest mean value per station (\bar{x} = 9.5) followed by San Juan (\bar{x} = 6.6) and Cauca (\bar{x} = 4.5).

Analysis of stomach content and FFG assignment: Stomach contents of the most representative morphotypes were analyzed and grouped according to their feeding groups' preferences. The material found was classified according to the nutrition types according to Cheshire et al. (2005). Classifications were as follows: FPOM (fine particulate organic matter, < 50 μ m); CPOM (coarse particulate organic matter, 50 μ m - 1 mm); OR (organisms' remains); PA (periphytic algae); and PT (plant tissue, > 1 mm).

Table 3 presents comparative information with the FFG allocation found through the diet presented in the stomach analysis of some representative taxa from tropical rivers in this study, and that reported by previous authors.

12 taxa were analyzed. 5 taxa had FPOM in their stomachs; 7 taxa had CPOM; 8 had plant tissue; 4 had traces of organisms, and only one had periphytic algae. This made it possible to assign a feeding function to each of them. The findings for nearly 92 % of the groups are consistent with reports by authors such as (Ramírez & Gutiérrez-Fonseca, 2014).

Table 4 presents the summary of the multifactorial ANOVA relating the abundances of



TABLE 3
Diet and Feeding Functional Group of some taxa identified in this research and reported in the literature

Taxa	Food diet	FFG in the current study	FFG reported in the literature
<i>Traverella</i>	FPOM, OR	CG, Pr	CG
<i>Baetodes</i>	FPOM, PT	Ft, Sc	Sc
<i>Prebaetodes</i>	FPOM	CG	CG
<i>Lepthophyes</i>	CPOM, PT	CG, Sh	CG
<i>Hetaerina</i>	FPOM, CPOM, OR, PA, PT	Pr	Pr
<i>Smicridea</i>	PT	Ft	Ft
<i>Phylloicus</i>	CPOM, PT	Sh	Sh
<i>Anacroneria</i>	CPOM, OR, PT	Pr	Pr
Chironomidae	CPOM	CG, Sh	CG
<i>Simulium</i>	FPOM, PT	Ft	Ft
<i>Physa</i>	CPOM, PT	Sc	Sc
<i>Gammarus</i>	CPOM, OR	Pr	Sh

CPOM: Coarse particulate organic matter, FPOM: Fine particulate organic matter OR: Organism remains, PA: Periphytic algae PT: Plant tissue, CG: Collector gatherers, Pr: Predator, Ft: Filter feeder, Sc: Scrapers, Sh: Shredders.

TABLE 4
Multifactorial variance analysis for the functional feeding groups

MAIN EFFECTS	Ft		CG		Pr		Sc		Sh	
	F	P Value	F	P Value	F	P Value	F	P Value	F	P Value
A: Season	0.49	0.6186	0.83	0.4444	0.48	0.6243	1.94	0.1583	0.76	0.4733
B: Altitude	4.70	0.0005	3.39	0.0051	5.14	0.0002	5.45	0.0001	4.75	0.0005
AxB	0.74	0.7394	0.24	0.9983	0.37	0.9809	1.88	0.0561	0.98	0.5002

the functional groups to the hydrological season and altitude factors, as well as the interaction between these factors.

These values show that statistically significant differences occur between FFG and Altitude. This last factor is directly related to the sampling site. The season does not appear to have a significant effect on the abundance of the functional feeding groups in the three streams.

Fig. 2 illustrates the spatial changes along the different basins studied. Pie charts, in different shades of gray, show the composition in percentages of the abundance of the functional feeding groups in the semi-quantitative sampling for each of the monitored stations. The numerical value in each segment of the pie chart corresponds to richness which, in the case of Nitrera, is higher than for the other two rivers. However, no differences are observed

between the sampling stations in La Nitrera. Meanwhile, for the San Juan River, there is a variation of the FFG in each period. Filter feeders were abundant at the time, while collectors were abundant in the middle and lower parts.

The diagrams presented in Fig. 3 shows the configuration of the ecological network through the relationships of the FFG with the modularity value for each of the stations in September 2018.

In the case of La Nitrera, the number of nodes is higher due to the presence of a greater number of morphotypes compared to the other rivers. The Shredder have then greatest contribution of biomass in La Nitrera (Fig. 3A, Fig. 3B, Fig. 3C), while in the San Juan River it is from collectors (Fig. 3D, Fig. 3E, Fig. 3F). In the Cauca River, the contribution of biomass varies between stations. At the station (Fig. 3G), the main contribution is from scrapers,




Fig. 2. Abundance and Richness distribution of the FFG following the altitudinal gradient in dry, transition and rainy seasons in 2017 and 2018.




Fig. 3. Trophic relationship diagrams for September 2018 at the different stations for the three riverine scenarios using ecological networks. **A.** Nitrera S1 September 2018 **B.** Nitrera S2 September 2018 **C.** Nitrera S3 September 2018 **D.** San Juan S1 September 2018 **E.** San Juan S2 September 2018 **F.** San Juan S3 September 2018 **G.** Cauca S1 September 2018 **H.** Cauca S2 September 2018 **I.** Cauca S3 September 2018. Each color indicates the number of modules

especially the Thiaridae family; at (Fig. 3H) it is from filter feeders although other groups contribute, while at CS3 (Fig. 3I) filter feeders were found as the only biomass contributors.

The analysis of variance components (AVC) using modularity as a dependent variable showed that the altitude is the factor that contributes most to the variance, with 65.49 %, followed by the season, with 27.57 %. Hence, a multifactorial ANOVA was carried out to establish whether there are statistical

differences for each principal or interaction effect in each watercourse studied. Table 5 presents multifactorial ANOVA indicating significant statistics differences between season and La Nitrera modularity.

Fig. 4 shows the variation of the modularity values in the different hydrological periods. Fig. 4A and Fig. 4B corresponding to July, Fig. 4C and Fig. 4D corresponding to August. Finally, Fig. 4E and Fig. 4F corresponding to September in 2017 and 2018 respectively. The

TABLE 5
Multifactorial variance analysis for Modularity

MAIN EFFECTS	La Nitrera		San Juan		Cauca	
	F	Value P	F	Value P	F	Value P
A: Season	3.15	0.0720	4.14	0.0490	2.64	0.1124
B: Altitude	3.24	0.0678	6.18	0.0179	0.01	0.9912
AxB	0.52	0.7207	3.37	0.0542	0.08	0.9856




Fig. 4. Modularity value hydrologic periods: dry (orange), transition (grey), rainy (blue) in the three streams in 2017 and 2018. A. July de 2017 B. July de 2018 C. August de 2017 D. August de 2018 E. September de 2017 F. September de 2018.

bars indicate the modularity value for each station. The shades of grey change according to the season. Spatial and temporal changes can be seen. The range of modularity in each figure varies from 0 to 1. If the value is closer to 1 then the degree of compartmentalization of the food web is greater, by contrast negative values indicate there are not a community structure base in the net. Nevertheless, all stations present a positive value. If the polygon remains spatially stable then it will close completely, which means that the modularity is homogeneous throughout the changes in elevation.

The most compartmentalized months are July 2018 (Fig. 4B) and September 2018 (Fig. 4F). It is important to bear in mind that, by August 2018 (Fig. 4D), the sample of organisms could not be taken because of a flood in the river at the SJS1 station. August 2018

for SJS1, CS3 stations and July, August, and September of 2017 for CS3 station were not possible to calculate modularity value due to the low or no number of organisms.

DISCUSSION

To the functional feeding groups analysis, the stomach determination show that fine organic matter was predominant, followed by organic matter in most of the digestive systems. There were some remains of organisms, generally mouthparts and antennae of *Diptera* sp. Additionally, there were plant structures (leaf litter, parts of internodes, seeds, shells, fibers, etc.). These results are in line with the statements of several authors that highlight the importance of FPOM as a dominant



food for most aquatic invertebrates (Ferreira et al., 2015; Moore et al., 2004; Tamaris-Turizo et al., 2018).

In the lower altitude areas, the richness of species is significantly reduced due to the reduction in slopes and the topology of the rivers, which results in a greater amount of deposition, which does not promote the availability of substrates. In general terms, La Nitrera presents a great abundance of FFG, mainly in the rainy season, due to the high heterogeneity of microhabitats (Tamaris-Turizo et al., 2018). Ft and CG are the most abundant in this stream in 2017, and Pr in 2018. The results indicate that, for the San Juan River, there is no significant variation between the FFGs during the hydrological seasons. The predominant FFGs are CG in the rainy season and Pr in the dry season. A high variation of FFGs based on the gradient was expected. However, lower richness was observed in the Cauca River, which is mainly due to aspects reported by other authors such as the geomorphology of the river, the short length of the main channel and the steep slopes of the river (Tamaris-Turizo et al., 2018; Webster & Meyer, 1997). Another point to consider is the impact of human activities as the gradient decreases, which have a significant effect on the composition of aquatic organisms. Baumgartner and Robinson (2017) mention due to natural gradients (stream order, substratum, altitude, etc.) and the dendritic structure of river networks cumulative environmental factors affecting biotic assemblages are expected in human dominated sites.

The changes in the distribution of aquatic macroinvertebrates according to their FFG showed a statistically significant variation due to altitude, which is associated with the sampling altitude gradient. If the gradient is lowered, the abundance of these FFGs also decreases. This variation occurs due to factors such as biotopes colonization availability and increases in water turbidity and water temperature (Dallas, 2007). The utilization of the model of ecological networks using the modular structure showed that the stations that presented greater FFG, that is to say, a greater diversity

of biotopes related to the number of ecological functions, may be able to buffer the propagation of disturbances, determining the resistance of these networks (Gauzens et al., 2015).

In the study, a variety of community responses were observed, depending on the type of stream and the intensity of hydrological alteration (Serna López et al., 2020). During the dry and rainy periods, the food webs had a smaller number of organisms, which can be seen in the modularity and compartmentalization of the webs. These aspects show how tributaries can influence longitudinal recovery trends (Mellado-Díaz et al., 2019). If a network has a high modularity degree, then this is reflected in the compartmentalization degree of the food webs. High modularity could indicate the network is resilient to external disturbances or changes.

As presented in the transition period, the potential recovery of aquatic communities along longitudinal gradients in the basin is maintained. The bars indicated that the transition period presented the highest modularity, which is due to the recolonization of some chorotypes resulting from the decrease in the bed speed. La Nitrera stream and the San Juan River presented higher values than the Cauca River, which may indicate that the altitudinal change affects the ecological network, presenting a greater degree of compartmentalization due to the number of nodes. Finally, this study represents a contribution to the analysis of aquatic macroinvertebrate communities through functional feeding groups, using the modularity of ecological networks along an altitude gradient in rivers in Colombia. Although further studies are required, it can be shown that the habitat conditions and changes in the configuration of the watercourses alter the stability of these relationships and reduce the species richness in low areas < 50 m.

Ethical statement: the authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal

procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

ACKNOWLEDGMENTS

This article was written within the framework of the doctoral thesis and with the support of *Minciencias*, convention 727 of 2015, and in cooperation with the research groups GeoLimna and Gepar of the faculty of engineering of the university of Antioquia. Some of the results obtained are thanks to work carried during the internship at the university of Vechta, Germany. This paper has not conflict of interests.

RESUMEN

Distribución altitudinal de los grupos funcionales de alimentación de macroinvertebrados acuáticos utilizando una red ecológica en arroyos andinos

Introducción: El análisis de grupos funcionales de alimentación (GFA) en macroinvertebrados acuáticos es importante para comprender la estructura, función y dinámica de los ecosistemas de procesos ecológicos. La modularidad se refiere al grado de compartimentación de las redes alimentarias y varía entre -1 y 1. Una red con un valor de modularidad cercano a 1 es resistente a las alteraciones y puede interpretarse como un factor indicativo para la estabilidad de las comunidades.

Objetivo: En este estudio se analizó la estructura trófica de los macroinvertebrados bentónicos, un elemento importante en la calidad ambiental, en el arroyo La Nitrera, el río San Juan y el río Cauca.

Métodos: El estudio contó con el apoyo de técnicas de redes ecológicas utilizando el software Gephi. En 2017 y 2018, estudiamos nueve sitios en estaciones secas, lluviosas y de transición, monitoreando cambios en el gradiente de altitud. En cada uno de los sitios se capturaron y determinaron los organismos y se recogió información fisicoquímica e hidráulica.

Resultados: El análisis de componentes de varianza permitió explicar la variabilidad de los datos relacionando las siguientes variables ambientales: GFA, diversidad, riqueza, modularidad, estación y tiempo. La ANOVA simple multifactorial indicó que existen cambios significativos en los GFA en relación con la altitud, y la modularidad con el tiempo. La asignación de los GFA se realizó mediante análisis estomacal e información secundaria.

Conclusión: La temporada de transición tuvo la mayor modularidad, posiblemente debido a la recolonización de

algunos biotopos provocada por la disminución de la velocidad del cauce. La Nitrera y San Juan presentaron valores superiores a los del Cauca, lo que puede indicar que el cambio altitudinal y la velocidad de las corrientes de agua influyen en la compartimentación de la red.

Palabras clave: modularidad; gradiente altitudinal; análisis estomacal; análisis de redes; estructura trófica de invertebrados.

REFERENCES

- Álvarez, L. F., & Daza, E. M. (2005). *Metodología para la utilización de los macroinvertebrados acuáticos como indicadores de la calidad del agua*. <http://repository.humboldt.org.co/handle/20.500.11761/31357>
- Anand, M., González, A., Guichard, F., Kolasa, J., & Parrott, L. (2010). Ecological systems as complex systems: Challenges for an emerging science. *Diversity*, 2(3), 395–410. <https://doi.org/10.3390/d2030395>
- Bastian, M., Heymann, S., & Jacomy, M. (2009). Gephi: An Open Source Software for Exploring and Manipulating Networks. *Third International AAAI Conference on Weblogs and Social Media*, 361–362. <https://doi.org/10.1136/qshc.2004.010033>
- Baumgartner, S. D., & Robinson, C. T. (2017). Changes in macroinvertebrate trophic structure along a land-use gradient within a lowland stream network. *Aquatic Sciences*, 79(2). <https://doi.org/10.1007/s00027-016-0506-z>
- Blondel, V. D., Guillaume, J. L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*. <https://doi.org/10.1088/1742-5468/2008/10/P10008>
- Boltovskoy, D., Izaguirre, I., & Correa, N. (1995). Feeding selectivity of *Corbicula fluminea* (Bivalvia) on natural phytoplankton. *Hydrobiologia*, 312(3), 171–182. <https://doi.org/10.1007/BF00015510>
- Castillo-Figueroa, D., Garzón-Salamanca, L. L., & Albaracín-Caro, J. F. (2018). Aquatic macroinvertebrates as water quality bioindicators in Colombia: A systematic review. *Neotropical Biology and Conservation*, 13(3), 235–248. <https://doi.org/10.4013/nbc.2018.133.06>
- Chará-Serna, A. M., Chará, J. D., Zúñiga, M. del C., Pedraza, G. X., & Giraldo, L. P. (2010). Trophic classification of aquatic insects in eight sheltered streams of the Colombian coffee ecoregion. *Universitas Scientiarum*, 15(1), 27–36. <https://doi.org/10.11144/javeriana.SC15-1.tcoa>
- Cheshire, K., Boyero, L., & Pearson, R. G. (2005). Food webs in tropical Australian streams: Shredders are not



- scarce. *Freshwater Biology*, 50(5), 748–769. <https://doi.org/10.1111/j.1365-2427.2005.01355.x>
- Cummins, K., Merritt, R., & Andrade, P. (2005). The use of invertebrate functional groups to characterize ecosystem attributes in selected streams and rivers in south Brazil. *Studies on Neotropical Fauna and Environment*, 40(1), 69–89. <https://doi.org/10.1080/01650520400025720>
- Dallas, H. F. (2007). The influence of biotope availability on macroinvertebrate assemblages in South African rivers: Implications for aquatic bioassessment. *Freshwater Biology*, 52(2). <https://doi.org/10.1111/j.1365-2427.2006.01684.x>
- Domínguez, E., & Fernández, H. (2009). *Macroinvertebrados bentónicos sudamericanos*. Fundación Miguel Lillo.
- Elosegui, E., & Sabater, S. (2009). *Conceptos y técnicas en ecología fluvial*. Fundación BBVA.
- Ferreira, W. R., Ligeiro, R., Macedo, D. R., Hughes, R. M., Kaufmann, P. R., Oliveira, L. G., & Callisto, M. (2015). Is the diet of a typical shredder related to the physical habitat of headwater streams in the Brazilian Cerrado? *Annales de Limnologie-International Journal of Limnology*, 51(2), 115–127. <https://doi.org/10.1051/limn/2015004>
- Gauzens, B., Thébault, E., Lacroix, G., & Legendre, S. (2015). Trophic groups and modules: Two levels of group detection in food webs. *Journal of the Royal Society Interface*, 12(106), 20141176. <https://doi.org/10.1098/rsif.2014.1176>
- Govedich, F. R., & Moser, W. E. (2015). Clitellata: Hirudinida and Acanthobdellida. *Thorp and Covich's Freshwater Invertebrates*, 565–588. <https://doi.org/10.1016/B978-0-12-385026-3.00023-1>
- Grilli, J., Rogers, T., & Allesina, S. (2016). Modularity and stability in ecological communities. *Nature Communications*, 7(1), 1–10. <https://doi.org/10.1038/ncomms12031>
- Gutiérrez, J. P. (2018). *Riverbank filtration in highly turbid rivers*. TU Delft University. <https://doi.org/10.4233/uuid:a6fe2afd-1f34-4b03-9687-1a5b627b64c2>
- Lasso, C. A., Morales-Betancourt, M. A., Vera-Ardila, M. L., & Linares, E. L. (2018). *Moluscos dulceacuícolas de Colombia*. Instituto Humboldt. <https://doi.org/10.21068/a2018n03>
- Latruesse, E. M., Stevaux, J. C., & Sinha, R. (2005). Tropical rivers. *Geomorphology*, 70(3–4), 187–206. <https://doi.org/10.1016/j.geomorph.2005.02.005>
- López, P. A. (2013). Propuesta de prevención y manejo de la contaminación por mercurio en la región de La Mojana. *Journal of Chemical Information and Modeling*, 53, 1689–1699. <https://doi.org/10.1017/CBO9781107415324.004>
- Martin, G. (2015). Gephi -Introduction to network analysis and visualization. *Digital Humanities*. <http://www.martingrandjean.ch/gephi-introduction/>
- Mellado-Díaz, A., Sánchez-González, J. R., Guareschi, S., Magdaleno, F., & Toro Velasco, M. (2019). Exploring longitudinal trends and recovery gradients in macroinvertebrate communities and biomonitoring tools along regulated rivers. *Science of the Total Environment*, 695, 133774. <https://doi.org/10.1016/j.scitotenv.2019.133774>
- Morales-Quintero, D. E., Velez-Macias, F. de J., & Guerrero-Hoyos, B. (2019). Caracterización hidrogeológica y geomorfológica de la cuenca y embalse del parque natural la Nitrera. *Revista de La Facultad de Ciencias*, 8(1), 73–99. <https://doi.org/10.15446/rev.fac.cienc.v8n1.71399>
- Muñoz, I., Romani, A. M., Rodrigues-Capítulo, A., Gonzales Esteban, J., & García-Berthou, E. (2009). Relaciones tróficas en el ecosistema fluvial. En A Elosegui (Ed.), *Conceptos y Técnicas En Ecología Fluvial* (pp. 1–22). Fundación BBVA.
- National Academy of Sciences. (1988). In E. O. Wilson (Ed.), *Biodiversity*. The National Academies Press. <https://doi.org/https://doi.org/10.17226/989>
- Olesen, J. M., Bascompte, J., Dupont, Y. L., & Jordano, P. (2007). The modularity of pollination networks. *Proceedings of the National Academy of Sciences of the United States of America*, 104(50), 19891–19896. <https://doi.org/10.1073/pnas.0706375104>
- Petersen, H. (2002). General aspects of collembolan ecology at the turn of the millennium. *Pedobiologia*, 46(3–4), 246–260.
- Pilosof, S., Porter, M. A., Pascual, M., & Kéfi, S. (2017). The multilayer nature of ecological networks. *Nature Ecology and Evolution*, 1(4), 1–9. <https://doi.org/10.1038/s41559-017-0101>
- Puerta, B., Molina, F., & Rodriguez, D. (2009). *Manual de laboratorio de procesos biológicos*. Reimpresos-Universidad de Antioquia.
- Ramírez, A., & Gutiérrez-Fonseca, P. E. (2014). Functional feeding groups of aquatic insect families in Latin America: A critical analysis and review of existing literature. *Revista de Biología Tropical*, 62, 155–167. <https://doi.org/10.15517/rbt.v62i0.15785>
- Roldán, G. (1988). *Guía para el estudio de los macroinvertebrados acuáticos del departamento de Antioquia*. Universidad de Antioquia.
- Roldán, G. (2003). *Bioindicación de la calidad del agua en Colombia (Uso del método BMWP/Col)*. Universidad de Antioquia.

- Roldán-Pérez, G. (2016). Los macroinvertebrados como bioindicadores de la calidad del agua: cuatro décadas de desarrollo en Colombia y Latinoamérica. *Revista de La Academia Colombiana de Ciencias Exactas, Físicas y Naturales*, 40(155), 254. <https://doi.org/10.18257/raccefyfyn.335>
- Rosenberg, D., & Resh, V. (1993). *Freshwater biomonitoring and benthic macroinvertebrates*. Chapman & Hall.
- Ruiz-Toro, J., Aguirre-Ramírez, N., Serna-López, J., Hernández-Atilano, E., & Vélez-Macias, F. (2020). Energía calórica, biomasa y estructura de los macroinvertebrados acuáticos en la reserva La Niterra, Concordia, Antioquia, Colombia. *Acta Biológica Colombiana*, 25(1), 8. <http://dx.doi.org/10.15446/abc.v25n1.76435>
- Schmera, D., Eros, T., & Heino, J. (2013). Habitat filtering determines spatial variation of macroinvertebrate community traits in northern headwater streams. *Community Ecology*, 14(1), 77–88. <https://doi.org/10.1556/ComEc.14.2013.1.9>
- Schmera, D., Heino, J., Podani, J., Erős, T., & Dolédec, S. (2017). Functional diversity: a review of methodology and current knowledge in freshwater macroinvertebrate research. *Hydrobiologia*, 787(1), 27–44. <https://doi.org/10.1007/s10750-016-2974-5>
- Serna López, J. P., Fernández Mc Cann, D. S., Vélez Macías, F. de J., & Aguirre Ramírez, N. J. (2020). An image processing method for recognition of four aquatic macroinvertebrates genera in freshwater environments in the Andean region of Colombia. *Environmental Monitoring and Assessment*, 192(10), 1–11. <https://doi.org/10.1007/s10661-020-08545-2>
- Stouffer, D. B., & Bascompte, J. (2011). Compartmentalization increases food-web persistence. *Proceedings of the National Academy of Sciences of the United States of America*, 108(9), 3648–3652. <https://doi.org/10.1073/pnas.1014353108>
- Tamaris-Turizo, C. E., Pinilla, G. A., & Muñoz, I. (2018). Trophic network of aquatic macroinvertebrates along an altitudinal gradient in a Neotropical mountain river. *Revista Brasileira de Entomologia*, 62(3), 180–187. <https://doi.org/10.1016/j.rbe.2018.07.003>
- Thorp, J. H., & Covich, A. P. (2009). *Ecology and Classification of North American Freshwater Invertebrates* (3rd Ed.). <https://doi.org/https://doi.org/10.1016/C2009-0-02669-5>
- Tomanova, S., Goitia, E., & Helešic, J. (2006). Trophic levels and functional feeding groups of macroinvertebrates in neotropical streams. *Hydrobiologia*, 556(1), 251–264. <https://doi.org/10.1007/s10750-005-1255-5>
- Torres, J. F., & Pinilla, G. A. (2011). Revisión de las características limnológicas de los sistemas acuáticos de la región de la Mojana. *Facultad de Ingeniería*, 70.
- Violle, C., Navas, M. L., Vile, D., Kazakou, E., Fortunel, C., Hummel, I., & Garnier, E. (2007). Let the concept of trait be functional! *Oikos*, 116(5), 882–892. <https://doi.org/10.1111/j.0030-1299.2007.15559.x>
- Webster, J. R., & Meyer, J. L. (1997). Organic matter budgets for streams: A synthesis. *Journal of the North American Benthological Society*, 16(1), 141–161. <https://doi.org/10.2307/1468247>