

Free-living *Acanthamoeba* and *Naegleria* spp. amebae in water sources of León, Nicaragua

Byron Leiva^{1,2,3}, Emma Clasdotter², Ewert Linder^{2,3} & Jadwiga Winięcka-Krusnell³

1. Departamento de Microbiología y Parasitología, Universidad Nacional Autónoma de Nicaragua, León, Nicaragua. Tel. 505-311 2947. Fax: 505-311 2947; byronleiva2001@yahoo.com.mx; byron.leiva@smi.ki.se
2. Microbiology and Tumor Biology Centre, Karolinska Institute, SE-171 77, Stockholm, Sweden; ewert.linder@smi.ki.se
3. Department of Parasitology, Mycology and Environmental Microbiology, Swedish Institute for Infectious Disease Control, SE-171 82 Solna, Sweden. Tel. +46-8-4572521. Fax +46-8-310525; jadwiga.winięcka-krusnell@smi.ki.se

Received 16-I-2007. Corrected 20-IX-2007. Accepted 04-X-2007.

Abstract: free-living amebae (FLA) are known to occur worldwide in water-related biotopes, but only limited information is available on these organisms in developing countries and so far no information on their presence is available from Nicaragua. The aims of this study were to evaluate the prevalence of potentially pathogenic *Acanthamoeba* spp. and *Naegleria* spp. in different water sources to which the population of León municipality is exposed. Since pathogenic amebae are thermotolerant, we were especially interested in the occurrence of FLA in geothermal areas. Water samples were collected from León area in Nicaragua: 88 samples were from rivers and springs, 111 from wells, 74 from water taps and 21 from water tanks in urban and suburban León and from three nearby geothermal areas of San Jacinto, Posoltega and Tipitapa. Amebae were identified using morphological and physiological criteria, immunohistochemical staining procedures and molecular methods. Indirect immunofluorescent test was performed on cysts and trophozoites fixed on microscopical slides and incubated for 30 min at room temperature in separate experiments with the following antibodies: rabbit-anti *N. fowleri*/*N. lovanesis* (Nf-Pab), mouse monoclonal antibody anti *N. fowleri* (Nf-5D12u), rabbit antibodies against *Acanthamoeba* spp. And fluorescent *in situ* hybridization (FISH) was performed using 18S rRNA-targeted fluorescent oligonucleotide probes. Probes: GSP for the detection of *Acanthamoeba* and NAEG1088 for the detection of *Naegleria*. Free-living amebae were recovered from approximately 43 % of the samples. *Acanthamoeba* spp was found in 21 % of samples from León municipality and in 2 % of samples from geothermal areas. Amoeboflagellates were found in 10 % of samples from León and in 19 % in geothermal areas. Fifty three percent of tested wells in the geothermal area contained thermotolerant amoeboflagellates. *Naegleria* spp. was identified in 24 out of 39 (61.5 %) of isolated amoeboflagellates. Twelve of them were assigned to *N. lovanesis* while none of the isolates could be identified as *N. fowleri*. However, the common presence of thermotolerant *Naegleria* in water, specially *N. lovanesis*, which is an indicator species for *N. fowleri*, suggests that also this pathogenic amoeba may pose a risk to public health in the area. On the other side, direct pathogenicity, free-living amebae are receiving increasing attention as reservoirs and potential vehicles for the transmission of bacteria in the environment. Thus the information provided in this study may serve as base-line for future studies on the role of free-living amebae e.g. in waterborne-disease outbreaks in the region. Among such potentially important enteropathogens are *Vibrio cholerae*, *E. coli* 0157, and *Helicobacter pylori*. Rev. Biol. Trop. 56 (2): 439-446. Epub 2008 June 30.

Key words: free-living amebae, amebic encephalitis, amebic keratitis; *Acanthamoeba*, *Naegleria*, water sanitation.

Free-living amebae (FLA) are a large group of organisms with worldwide distribution. Humans are naturally exposed to these amebae since they are common in the

environment, especially in all water-related biotopes. Among FLA there are facultative parasites of man and animals belonging to genera *Acanthamoeba*, *Naegleria* and *Balamuthia*

causing severe infections in the central nervous system. *Acanthamoeba* spp. are also capable of infecting the skin, the lungs, the eyes or causing disseminated infections in susceptible hosts, immunodeficiency apparently being an increasingly important risk factor (Visvesvara 1993, De Jonckheere 1980, De Jonckheere and Brown 1997, Senzsi *et al.* 1998, Maraciano-Cabral *et al.* 2000).

Notably *Naegleria* spp. grow preferentially in water temperatures of 25 to 44 °C. This parasite is potentially pathogenic in man causing fatal primary amoebic meningoencephalitis (PAM). Typically children and young adults are the victims and most cases have been described from the United States, Australia and the Czech republic. In Nicaragua, geothermal areas are frequently used for recreation, and thermal water may be the only water source of local households. However, there is no information on the presence of potentially pathogenic free-living amoebae in the region. Also it is not surprising that cases of PAM have not been described in Nicaragua, considering the difficulties in establishing the diagnosis by cytological analysis (Benson *et al.* 1985) and the restricted availability of NMR (Falcone *et al.* 1994). Importantly diagnostic difficulties occur even under favorable health care conditions (Hecht *et al.* 1972, Bakardjiev *et al.* 2002, Gyori 2003, Stephany *et al.* 2004).

The aims of this study were to evaluate the prevalence of FLA in different sources of water in the urban area of León municipality and in the geothermal area, and search for thermotolerant, potentially pathogenic *Acanthamoeba* spp. and *Naegleria* spp. in the isolated material. Collected data will serve as a base-line information for future studies on association between FLA and clinical conditions caused by these organisms. Furthermore such information may be important considering outbreaks of diarrhea caused by enteropathogenic bacteria in water.

MATERIALS AND METHODS

Sample collection and cultivation of amoebae: water samples were collected from León

area in Nicaragua: 88 samples were from rivers and springs, 111 from wells, 74 from water taps and 21 from water tanks in urban and suburban León and from 3 nearby geothermal areas of San Jacinto, Posoltega and Tipitapa.

Water samples were collected into sterile 50 ml tubes and transported to the laboratory at Microbiology Department, University of León where they were studied using established methods for identification essentially as described (Schuster 2002). Amoebae were isolated from specimens filtered through 0.45 µm filters attached to sterile syringes. Filters were placed on non-nutrient agar plates covered with viable *Escherichia coli* K12 strain. Part of the samples were centrifuged for 10 min at 4 000 rpm, supernatant was removed and the sediment was re-suspended with remaining water by tapping the tube. One or two drops were placed on agar-*E. coli* plates. Agar plates were incubated at 40 °C for the isolation of thermotolerant strains. Growing amoebae were repeatedly sub-cultured and cloned by limiting dilution.

Reference strains: strains of *Acanthamoeba castellanii*, *A. rhyssodes*, *Hartmannella vermiformis*, *Naegleria gruberi* and *N. lovansensis*. The ocular *Acanthamoeba* strains were isolated at SMI and typed at PHL, Bath, England. *Hartmannella* was obtained from Dr. E. Hadas, University of Poznan, Poland and *Naegleria gruberi* from ATCC Atlanta. Amoebae were cultivated on non-nutrient agar plates covered with *E. coli* and used as control material for identification tests.

Identification of isolates: initial identification of amoebae was based on microscopical observation of living and fixed cysts and trophozoites. Measurements were performed with an ocular micrometer at 400 x and 1 000 x magnification. Data from 50 individuals of each strain were collected and characterization was made using morphological criteria. Thermotolerance of amoebae was assessed by testing for ability to grow on agar plates incubated at 40, 42, 45 and 47 °C for five to seven days. The enflagellation test (Behets *et*

al. 2003) was performed on 24-wells plates. Trophozoites collected from agar plates were added to wells containing two ml of sterile, distilled water and incubated at 37 °C. The plates were inspected after 30, 60, 90 and 120 min for the presence of free-swimming, flagellated forms. Indirect immunofluorescent test was performed on cysts and trophozoites fixed on microscopical slides and incubated for 30 min at room temperature in separate experiments with the following antibodies: rabbit-anti *N. fowleri*/*N. lovaniensis* (Nf-Pab), mouse monoclonal antibody anti *N. fowleri* (Nf-5D12u) (both from Indicia Biotechnology, Oullins, France), rabbit antibodies against *Acanthamoeba* spp. (were from Dr J-Å Liljeqvist, University of Gothenburg, Sweden). After washing with PBS slides were incubated again for 30 min at room temperature with the appropriate FITC conjugated sheep anti-rabbit anti-immunoglobulin antibodies (Sigma F 1010, Saint Louis, Missouri). After subsequent washing with PBS, slides were mounted in non-fading medium (Vectashield, Vector laboratories, Inc. Burlingame, CA.) and observed in the microscope. The presence of cellulose in cyst wall of *Acanthamoeba* was demonstrated using the cellulose-binding domain of cellulase essentially as described (Linder *et al.* 2002). Fluorescent *in situ* hybridization (FISH) was performed using 18S rRNA- targeted fluores-

cent oligonucleotide probes according to the protocols described before (Grimm *et al.* 2001, Stothard *et al.* 1999). Probes: GSP for the detection of *Acanthamoeba* and NAEG1088 for the detection of *Naegleria* were synthesized by Scandinavian Gene Synthesis AB, 73122 Köping, Sweden.

RESULTS

One hundred twenty five out of the 294 samples (42.5%) collected from different water sources yielded growth of FLA. Amebae could be recovered from 75 out of 178 (42%) water samples collected in the León municipality and an almost identical proportion was recovered from samples from the geothermal areas (50 out of 116.43%) (Table 1 and 2). However, a detailed analysis showed some differences; the lowest prevalence of amebae, 17 out of 74 (23%) was found in tap water in León municipality and the highest prevalence in wells from geothermal areas, 25 out of 36 (69%), (Table 1 and 2). There were significantly higher levels of amoeboflagellates from wells in geothermal areas as compared to wells in the León municipality, 53% as opposed to 15%.

The majority of samples contained multiple species of amebae, however, only *Acanthamoeba* and amebae developing flagellated forms, were considered in this study.

TABLE 1
Samples from León municipality in Nicaragua

Sources	N°	<i>Acanthamoeba</i> spp. (%)	Amoeboflagellates (%)	¹ Other (%)	Total pos	
					N°	(%)
Water tank	21	(29)	(5)	(10)	9	43
Tap water	74	(19)	(4)	(0)	17	23
Wells	75	(21)	(15)	(21)	43	57
Stream	8	(13)	(25)	(38)	6	75
Total	178	(21)	(10)	(12)	75	42

1 Other = *Hartmanella* sp., Free living amoebae not identified.
pos = positive

TABLE 2
Samples from geothermal areas in Nicaragua

Sources	N°	<i>Acanthamoeba</i> spp.	Amoebaflagellates	¹ Other	Total pos	
		%	(%)	(%)	N°	(%)
Wells	36	(0)	(53)	(17)	25	69
Stream	80	(5)	(8)	(25)	25	31
Total	116	(2)	(19)	(22)	50	43

1 Other = *Hartmanella* sp., Free living amoebae not identified
 pos. = positive

TABLE 3
Growth of thermotolerant free-living amoebae at different temperatures

	<i>Acanthamoeba</i>			Amoebaflagellates		
	40°C	42°C	≥45°C	40°C	42°C	≥45°C
León Municipality	32	5	0	9	8	0
Geothermal areas	2	0	0	6	5	11
Total	34	5	0	15	13	11




Fig. 1. A Identification of cyst of *Acanthamoeba* spp. stained with monoclonal anti-cellulase antibodies. (B) Identification of *Acanthamoeba* spp. by fluorescence *in situ* hybridization (FISH) with 18S rRNA gene specific probe.

TABLE 4
Identification of Naegleria isolates by different methods

	Enflagellation	FISH	≥42 °C	Nf-Pab	Nf-5D12u
León Municipality	17	8	8	3	0
Geothermal areas	22	16	16	9	0
Total	39	24	24	12	0

FISH = fluorescent in situ hybridization

IF = immunofluorescence microscopy

Nf-Pab = anti *N. fowleri* /*N. lovaniensis* polyclonal antibodies

Nf-5D12u = anti *N. fowleri* mouse monoclonal antibodies




Fig. 2. A. Identification of *Naegleria* spp. by FISH with 18S rRNA *Naegleria* probe. (B) Identification of *Naegleria* with polyclonal rabbit-anti *N. fowleri*/*N. lovaniensis* (NF-Pab) by indirect immunofluorescence.

The average water temperature of samples collected in the area of León was 27.5 °C while the temperature of water collected in San Jacinto, Posoltega and Tipitapa was 38.5 °C ranging from 34-50 °C. Thermotolerant amoebae growing at ≥40 °C were identified among both *Acanthamoeba* and amoeboflagellate isolates. Recovery of amoebae was similar in the 3 geothermal areas. Half of the amoeboflagellates, 11 out of 22 isolated from the geothermal areas, showed growth at temperatures ≥45 °C (Table 3).

Thirty-nine isolates (31.2% of positive samples) were classified as *Acanthamoeba*

spp. by applying morphological criteria. Their identity was further confirmed by reactivity of trophozoites with anti-*Acanthamoeba* antibodies, the presence of cellulose in their cysts and by the FISH technique. All 39 isolates were positive by all methods used for identification.

Twenty-four flagellated isolates (19.2% of positive samples) were classified as *Naegleria* spp. based on positive FISH with *Naegleria* specific probe. Twelve isolates (50%) were tentatively assigned to *N. lovaniensis* based on the reactivity with Nf-Pab antibody (Table 4, Fig. 2), as all 12 failed to react with *N. fowleri* specific monoclonal Nf-5D12u antibodies.

DISCUSSION

The results in this study show that potentially pathogenic FLA are common in the León area and widely distributed even in drinking water. Morphological criteria and cultivation at different temperatures form the basis for species identification. It is known that the pathogenicity of FLA correlates with their capacity to tolerate temperatures of 37 °C or more (De Jonckheere 1980) and that non-virulent and non-pathogenic strains are unable to grow at temperatures above 37 °C (Visvesvara 1993).

All tests for *Acanthamoeba* were consistently positive. This is true also for the cellulase binding method described by us (Linder *et al.* 2002). The enflagellation test is used to identify genus *Naegleria*. However, when the detection of *Naegleria* spp. is based on the enflagellation test, false negatives are likely to occur, leading to significant underestimations (Behets *et al.* 2003). Specific PCR probes for *Naegleria* are valuable tools for the identification of environmental protozoa (Grimm *et al.* 2001). Apparently a higher sensitivity is achieved using such methods. Thus it was possible to detect *N. fowleri* in Yellowstone and Grand Teton National parks by PCR, which was not the case in a previous study using conventional cultivation and microscopy methods (Sheehan *et al.* 2003).

In our study FLA were found in 43% of water samples studied. Such a high prevalence is not unique. In a recent study from Bulgaria, the prevalence was 61% in natural fresh waters and in river samples as high as 94% (Tsvetkova *et al.* 2004). In Germany *Acanthamoeba* was found in 79% of isolates of river water (Hoffmann and Michel 2001). That tap water may contain FLA has been well documented by analyzing the different steps of water purification (Hoffmann and Michel 2001). Also our unpublished studies showed that water treatment procedures employed at the local municipal water supplier in the Stockholm region, Stockholm water, fail to eliminate *Acanthamoeba*, it is well established that whereas *Acanthamoeba* spp. are common in natural and tap water glob-

ally, presence of *Naegleria* spp. is restricted to areas with naturally warm water (John 1982). The increasing use of contact lenses and the difficulties involved in maintaining their sterility, makes the presence of *Acanthamoeba* in human environments a potential threat. There is accumulating information on the prevalence of FLA in areas with a hot climate (Lawande 1983, Ugonabo and Gugani 1989, Rivera *et al.* 1989, Bose *et al.* 1990, Hamado *et al.* 1993, Da Silva and Da Rosa 2003). However, little is known about the occurrence of *Acanthamoeba* keratitis in poor countries such as Nicaragua.

In thermal areas we found significantly more *Naegleria* than *Acanthamoeba*. Such environmental conditions occur in geothermal areas globally, but may occur also in the environment, which has been heated by other means. Raised temperatures during the hot summer months or warm water from power plants facilitate the growth of *N. fowleri* (Martinez and Visvesvara 1997). However, in the cyst form these protozoa are extremely resistant and may occur in dust (Lawande 1983, Da Silva and Da Rosa 2003).

Reports of fatal meningo-encephalitis due to *N. fowleri* in geothermally heated water in Australia, and in Bath, England, have imposed restrictions in the usage of natural warm water for recreational purposes. That the presence of *Naegleria* in our environment poses a real threat is reflected by research aiming at inactivation of amoebae in large volumes of fresh water (Vernhes *et al.* 2002).

In addition to their direct pathogenicity, free-living amoebae are receiving increasing attention as reservoirs and potential vehicles for the transmission of bacteria in the environment. Thus the information provided in this study may serve as base-line for future studies on the role of free-living amoebae e.g. in water-borne-disease outbreaks in the region. Among such potentially important enteropathogens are *Vibrio cholerae*, *E. coli* 0157, and *Helicobacter pylori* (King *et al.* 1988, Winiacka-Krusnell *et al.* 2001, Barker *et al.* 2002).

In summary, our study showed that *Acanthamoeba* and *Naegleria* are found

regularly in León municipality and that *Naegleria* is common in the geothermal area studied. The common presence of *N. lovaniensis*, which is an indicator species for *N. fowleri* suggests that recreational water contact in thermal area may pose a risk to public health.

ACKNOWLEDGMENTS

We thank students of the UNAN-León for participation in the collection of the water samples and Brenda Mora for laboratory work. This study was supported by the Swedish Agency for Research Cooperation with Developing Countries (SAREC).

RESUMEN

Las amebas de vida libre (AVL) son un grupo de organismos de distribución mundial. Entre las AVL hay parásitos facultativos en humanos y otros animales, los cuales pertenecen a los géneros *Acanthamoeba*, *Naegleria* y *Balamuthia* que causan infecciones severas en el sistema nervioso central. Sin embargo no se tiene ninguna información de Nicaragua. El objetivo de este estudio fue evaluar la presencia de amebas de vida libre, en diferentes fuentes de agua de la parte urbana y rural del Departamento de León, y áreas geotérmicas de Nicaragua. Estas amebas fueron identificadas usando criterios morfológicos, fisiológicos, histoquímicos y moleculares. En los resultados se encontró amebas de vida libre en el 43% del total de las muestras. En la municipalidad de León, se encontraron 21% de *Acanthamoeba* sp. y en las áreas geotérmicas un 2%. Las amebaflagelados tipo *Naegleria* fueron 10% y 19% respectivamente. Del grupo amebaflagelados fueron 24, de las cuales 12 se identificaron como *N. lovaniensis*. En estas muestras no se aisló *N. fowleri* (ameba patógena), sin embargo, la presencia de *N. lovaniensis* es un indicador de la presencia de *N. fowleri*, la cual puede ser un riesgo a la salud pública. Además, estas amebas pueden servir como vectores de bacterias enteropatógenas.

Palabras claves: amebas de vida libre, encefalitis amebiana, queratitis amebiana, *Acanthamoeba*, *Naegleria*, saneamiento de agua.

REFERENCES

- Bakardjiev, A., C. Glaser, F. Schuster & G.S. Visvesvara. 2002. Three-year-old girl with fever and coma. *Pediatr. Infect. Dis. J.* 21: 85-6.
- Barker, J., T.J. Humphrey & M.W.R. Brown. 1999. Survival of *Escherichia coli* 0157 in a soil protozoan: implications for disease. *FEMS Microbiol. Lett.* 173: 291-295.
- Behets, J., F. Seghi, P. Declerck, L. Verelst, L. Duvivier, A. Van Damme & F. Ollevier. 2003. Detection of *Naegleria* spp. and *Naegleria fowleri*: a comparison of flagellation tests, ELISA and PCR. *Water Sci. Technol.* 47: 117-122.
- Benson, R.L., L. Ansbacher, R.E. Hutchison & W. Rogers. 1985. Cerebrospinal fluid centrifuge analysis in primary amebic meningoencephalitis due to *Naegleria fowleri*. *Arch. Pathol. Lab. Med.* 109: 668-671.
- Bose, K., D. Ghosh, K. Ghosh, A. Bhattacharya & S. Das. 1990. Characterization of potentially pathogenic free-living amebae in sewage samples of Calcutta, India. *Braz. J. Med. Biol. Res.* 23: 1271-1278.
- Da Silva, M.A. & J.A. Da Rosa. 2003. Isolation of potentially pathogenic free-living amoebas in hospital dust. *Rev. Saude Publica*, 37: 242-246.
- De Jonckheere, J. 1980. Growth characteristics, cytopathic effect in cell culture, and virulence in mice of 36 type strains belonging to 19 different *Acanthamoeba* spp. *Appl. Environ. Microbiol.* 39: 681-685.
- De Jonckheere, J.F. & S. Brown. 1997. Primary amebic meningoencephalitis in a patient with AIDS: unusual protozoological findings. *Clin. Infect. Dis.* 25: 943-944.
- Falcone, S., R.M. Quencer & M.J. Post. 1994. Magnetic resonance imaging of unusual intracranial infections. *Top. Magn. Reson. Imaging* 6: 41-52.
- Grimm, D., W.F. Ludwig, B.C. Brandt, R. Michel, K.H. Schleifer, J. Hacker & M. Steinert. 2001. Development of 18S rRNA-targeted oligonucleotide probes for specific detection of *Hartmannella* and *Naegleria* in *Legionella*-positive environmental samples. *Syst. Appl. Microbiol.* 24: 76-82.
- Gyori, E. 2003. December 2002: 19-year old male with febrile illness after jet ski accident. *Brain Pathol.* 13: 237-239.
- Hamado, H., S. Aufy, I. el-Hayawan, M. Saleh & I. Nagaty. 1993. Study of free-living amebae in Egypt. *J. Egypt Soc. Parasitol.* 23: 631-637.
- Hecht, R.H., A.H. Cohen, J. Stoner & C. Irwin. 1972. Primary amebic meningoencephalitis in California. *Calif. Med.* 117: 69-73.

- Hoffmann, R. & R. Michel. 2001. Distribution of free-living amoebae (FLA) during preparation and supply of drinking water. *Int. J. Hyg. Environ. Health* 203: 215-219.
- John, D.T. 1982. Primary amebic meningoencephalitis and the biology of *Naegleria fowleri*. *Annu. Rev. Microbiol.* 36: 101-123.
- King, C.H., E.B. Jr. Shotts, R.E. Wooley & K.G. Porter. 1988. Survival of coliforms and bacterial pathogens within protozoa during chlorination. *Appl. Environ. Microbiol.* 54: 3023-3033.
- Lawande, R.V. 1983. Recovery of soil amoebae from the air during the harmattan in Zaria, Nigeria. *Ann. Trop. Med. Parasitol.* 77: 45-49.
- Linder, M., J. Winiacka-Krusnell & E. Linder. 2002. Use of recombinant cellulose-binding domains of *Trichoderma reesei* cellulase as a selective immunocytochemical marker for cellulose in protozoa. *Appl. Environ. Microbiol.* 68: 2503-2508.
- Marciano-Cabral, F., R. Puffenbarger & G.A. Cabral. 2000. The increasing importance of *Acanthamoeba* infections. *J. Eukaryot. Microbiol.* 47: 29-36.
- Martinez, A.J. & G.S. Visvesvara. 1997. Free-living, amphizoic and opportunistic amoebae. *Brain Pathol.* 7: 583-598.
- Rivera, F., F. Lares, E. Gallegos, E. Ramirez, P. Bonilla, A. Calderon, J.J. Martinez, S. Rodriguez & J. Alcocer. 1989. Pathogenic amoebae in natural thermal waters of three resorts of Hidalgo, Mexico. *Environ. Res.* 50: 289-295.
- Schuster, F.L. 2002. Cultivation of pathogenic and opportunistic free-living amoebae. *Clin. Microbiol. Rev.* 15: 342-354.
- Sheehan, K.B., J.A. Fagg, M.J. Ferris & J.M. Henson. 2003. PCR detection and analysis of the free-living amoeba *Naegleria* in hot springs in Yellowstone and Grand Teton National Parks. *Appl. Environ. Microbiol.* 69: 5914-5918.
- Stephany, J.D., G.S. Pearl & O.R. Gonzalez. 2004. Pathologic quiz case: headache in an 8 year old child. Primary amebic meningoencephalitis due to *Naegleria fowleri*. *Arch. Pathol. Lab. Med.* 128: 33-34.
- Stothard, D.R., J. Hay, J.M. Schroeder-Diedrich, D.V. Seal & T.J. Byers. 1999. Fluorescent oligonucleotide probes for clinical and environmental detection of *Acanthamoeba* and the T4 18S rRNA gene sequence type. *J. Clin. Microbiol.* 37: 2687-2693.
- Szenasi, Z., T. Endo, K. Yagita & E. Nagy. 1998. Isolation, identification and increasing importance of 'free-living' amoebae causing human disease. *J. Med. Microbiol.* 47: 5-16.
- Tsvetkova, N., M. Schild, S. Panaiotov, R. Kurdova-Mintcheva, B. Gottstein, J. Walochnik, H. Aspöck, M.S. Lucas & N. Müller. 2004. The identification of free-living environmental isolates of amoebae from Bulgaria. *Parasitol. Res.* 92: 405-413.
- Ugonabo, J. & H. Gugani. 1989. Nasal carriage of *Naegleria fowleri* and its environmental occurrence in Boro state, Nigeria. *J. Commun. Dis.* 21: 111-113.
- Visvesvara, G.S. 1993. Epidemiology of infections with free-living amoebae and laboratory diagnosis of microsporidiosis. *Mt. Sinai J. Med.* 60: 283-288.
- Vernhes, M.C., A. Benichou, P. Pernin, P.A. Cabanes & J. Teissie. 2002. Elimination of free-living amoebae in fresh water with pulsed electric fields. *Water Research* 36: 3429-3438.
- Winiacka-Krusnell, J. & E. Linder. 2001. Bacterial infections of free-living amoebae. *Res. Microbiol.* 152: 613-619.